# (19) World Intellectual Property Organization

International Bureau



# Rec'd PCT/PTO 07 SEP 2004

# 

(43) International Publication Date 18 March 2004 (18.03.2004)

PCT

(10) International Publication Number WO 2004/022778 A1

(51) International Patent Classification7: G01N 33/53

C12Q 1/68,

(AU). O'BRIEN, Philippa [GB/AU]; 73 Ocean Street, Woollahra, New South Wales 20258 (AU).

(21) International Application Number:

PCT/AU2003/001166

(74) Agent: F B RICE & CO.; 139 Rathdowne Street, Carlton,

(22) International Filing Date:

5 September 2003 (05.09.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2002951346

5 September 2002 (05.09.2002)

(71) Applicant (for all designated States except US): GARVAN INSTITUTE OF MEDICAL RESEARCH [AU/AU]; C/-St Vincent's Hospital, 384 victoria Street, Darlinghurst, New South Wales 2010 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SUTHERLAND, Robert [NZ/AU]; 20 Northcot Road, Lindfield, New South Wales 2070 (AU). HENSHALL, Susan [AU/AU]; 14 Mosman Street, Mosman, New South Wales 2068

- Victoria 3053 (AU). (81) Designated States (national): AE, AG, AL, AM, AT, AU,
- AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS OF DIAGNOSIS AND PROGNOSIS OF OVARIAN CANCER

(57) Abstract: The present invention provides novel genes and proteins for diagnosing ovarian cancer and/or a likelihood for survival, or recurrence of disease, wherein the expresson of the genes and proteins is up-regulated or down-regulated or associated with the occurrence or recurrence of a specific scanner sub-type. The ovarian cancer-associated genes and proteins of the invention are specifically exemplified by the genes and proteins set forth in Tables 1 to 3 and the Sequence Listing.

WO 2004/022778 PCT/AU2003/001166

1

#### METHODS OF DIAGNOSIS AND PROGNOSIS OF OVARIAN CANCER

#### Field of the invention

The present invention relates to the identification of nucleic acid and protein expression profiles and nucleic acids, products, and antibodies thereto that are involved in ovarian cancer; and to the use of such expression profiles and compositions in the diagnosis, prognosis and therapy of ovarian cancer. More particularly, this invention relates to novel genes that are expressed at elevated or reduced levels in malignant tissues and uses therefor in the diagnosis of cancer or malignant tumors in human subjects. This invention also relates to the use of nucleic acid or antibody probes to specifically detect ovarian cancer cells, such as, for example, in the ovarian surface epithelium, wherein over-expression or reduced expression of nucleic acids hybridizing to the probes is highly associated with the occurrence and/or recurrence of an ovarian tumor, and/or the likelihood of patient survival. The diagnostic and prognostic test of the present invention is particularly useful for the early detection of ovarian cancer or metastases thereof, or other cancers, and for monitoring the progress of disease, such as, for example, during remission or following surgery or chemotherapy. The present invention is also directed to methods of therapy wherein the activity of a protein encoded by a diagnostic/prognostic gene described herein is modulated.

20

5

10

15

#### Background of the invention

#### 1. General

As used herein the term "derived from" shall be taken to indicate that a specified integer are obtained from a particular source albeit not necessarily directly from that source.

25

Unless the context requires otherwise or specifically stated to the contrary, integers, steps, or elements of the invention recited herein as singular integers, steps or elements clearly encompass both singular and plural forms of the recited integers, steps or elements.

30

The embodiments of the invention described herein with respect to any single embodiment shall be taken to apply *mutatis mutandis* to any other embodiment of the invention described herein.

10

20

25

30

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated step or element or integer or group of steps or elements or integers but not the exclusion of any other step or element or integer or group of elements or integers.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations or any two or more of said steps or features.

The present invention is not to be limited in scope by the specific examples described herein. Functionally equivalent products, compositions and methods are clearly within the scope of the invention, as described herein.

The present invention is performed without undue experimentation using, unless otherwise indicated, conventional techniques of molecular biology, microbiology, virology, recombining DNA technology, peptide synthesis in solution, solid phase peptide synthesis, and immunology. Such procedures are described, for example, in the following texts that are incorporated herein by reference:

- Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratories, New York, Second Edition (1989), whole of Vols I, II, and III;
- DNA Cloning: A Practical Approach, Vols. I and II (D. N. Glover, ed., 1985), IRL Press, Oxford, whole of text;
- Oligonucleotide Synthesis: A Practical Approach (M. J. Gait, ed., 1984) IRL Press, Oxford, whole of text, and particularly the papers therein by Gait, pp1-22; Atkinson et al., pp35-81; Sproat et al., pp 83-115; and Wu et al., pp 135-151;
- 4. Nucleic Acid Hybridization: A Practical Approach (B. D. Hames & S. J. Higgins, eds., 1985) IRL Press, Oxford, whole of text;
- 5. Perbal, B., A Practical Guide to Molecular Cloning (1984);
- 6. Wünsch, E., ed. (1974) Synthese von Peptiden in Houben-Weyls Metoden der
  Organischen Chemie (Müler, E., ed.), vol. 15, 4th edn., Parts 1 and 2, Thieme,
  Stuttgart.

10

15

20

25

30

7. Handbook of Experimental Immunology, Vols. I-IV (D. M. Weir and C. C. Blackwell, eds., 1986, Blackwell Scientific Publications).

This specification contains nucleotide and amino acid sequence information prepared using Patentin Version 3.1, presented herein after the claims. Each nucleotide sequence is identified in the sequence listing by the numeric indicator <210> followed by the sequence identifier (e.g. <210>1, <210>2, <210>3, etc). The length and type of sequence (DNA, protein (PRT), etc), and source organism for each nucleotide sequence, are indicated by information provided in the numeric indicator fields <211>, <212> and <213>, respectively. Nucleotide sequences referred to in the specification are defined by the term "SEQ ID NO:", followed by the sequence identifier (eg. SEQ ID NO: 1 refers to the sequence in the sequence listing designated as <400>1).

The designation of nucleotide residues referred to herein are those recommended by the IUPAC-IUB Biochemical Nomenclature Commission, wherein A represents Adenine, C represents Cytosine, G represents Guanine, T represents thymine, Y represents a pyrimidine residue, R represents a purine residue, M represents Adenine or Cytosine, K represents Guanine or Thymine, S represents Guanine or Cytosine, W represents Adenine or Thymine, H represents a nucleotide other than Guanine, B represents a nucleotide other than Adenine, V represents a nucleotide other than Thymine, D represents a nucleotide other than Cytosine and N represents any nucleotide residue.

# 2. Description of the related art

Cancer is a multi-factorial disease and major cause of morbidity in humans and other animals, and deaths resulting from cancer in humans are increasing and expected to surpass deaths from heart disease in future. Carcinomas of the lung, prostate, breast, colon, pancreas, and ovary are major contributing factors to total cancer death in humans. For example, prostate cancer is the fourth most prevalent cancer and the second leading cause of cancer death in males. Similarly, cancer of the ovary is the second most common cancer of the female reproductive organs and the fourth most common cause of cancer death among females. With few exceptions, metastatic disease from carcinoma is fatal. Even if patients survive their primary cancers, recurrence or metastases are common.

It is widely recognized that simple and rapid tests for solid cancers or tumors have considerable clinical potential. Not only can such tests be used for the early diagnosis of

WO 2004/022778 PCT/AU2003/001166

4

cancer but they also allow the detection of tumor recurrence following surgery and chemotherapy. A number of cancer-specific blood tests have been developed which depend upon the detection of tumor-specific antigens in the circulation (Catalona, W.J., et al., 1991, "Measurement of prostate-specific antigen in serum as a screening test for prostate cancer", N. Engl. J. Med. 324, 1156-1161; Barrenetxea, G., et al., 1998, "Use of serum tumor markers for the diagnosis and follow-up of breast cancer", Oncology, 55, 447-449; Cairns, P., and Sidreansky, D., 1999, "Molecular methods for the diagnosis of cancer". Biochim. Biophys. Acta. 1423, C 11-C 18).

Ovarian cancer is the fourth most frequent cause of cancer death in females and in the United States, and accounts for approximately 13,000 deaths annually. Furthermore, ovarian cancer remains the number one killer of women with gynaecological malignant hyperplasia and the incidence is rising in industrialized countries. The etiology of the neoplastic transformation remains unknown although there is epidemiological evidence for an association with disordered endocrine function. The incidence of ovarian carcinoma is higher in nulliparous females and in those with early menopause.

Most ovarian cancers are thought to arise from the ovarian surface of epithelium (OSE). Epithelial ovarian cancer is seldom encountered in women less than 35 years of age. Its incidence increases sharply with advancing age and peaks at ages 75 to 80, with the median age being 60 years. The single most important known risk factor is a strong familial history of breast or ovarian cancer. To date, little is known about the structure and function of the OSE cells. It is known that the OSE is highly dynamic tissue that undergoes morphogenic changes, and has proliferative properties sufficient to cover the ovulatory site following ovulation. Morphological and histochemical studies suggest that the OSE has secretory, endocytotic and transport functions which are hormonally-controlled (Blaustein and Lee, *Oncol. 8*, 34-43, 1979; Nicosia and Johnson, *Int. J. Gynecol. Pathol.*, 3, 249-260, 1983; Papadaki and Beilby, *J. Cell Sci. 8*, 445-464, 1971; Anderson *et al.*, *J. Morphol.*, 150, 135-164, 1976).

30

35

10

15

20

25

Ovarian cancers are not readily detectable by diagnostic techniques (Siemens *et al.*, *J. Cell. Physiol.*, 134: 347-356, 1988). In fact, the diagnosis of carcinoma of the ovary is generally only possible when the disease has progressed to a late stage of development. Approximately 75% of women diagnosed with ovarian cancer are already at an advanced stage (III and IV) of the disease at their initial diagnosis. During the past 20 years, neither diagnosis nor five year survival rates have greatly improved for these patients. This is

substantially due to the high percentage of high-stage initial detection of the disease. There is therefore a need to develop new markers that improve early diagnosis and thereby reduce the percentage of high-stage initial diagnoses.

A number of proteinaceous ovarian tumor markers were evaluated several years ago, 5 however these were found to be non-specific, and determined to be of low value as markers for primary ovarian cancer (Kudlacek et al., Gyn. Onc. 35, 323-329, 1989; Rustin et al., J. Clin. Onc., 7, 1667-1671, 1989; Sevelda et al., Am. J. Obstet. Gynecol., 161, 1213-1216, 1989; Omar et al., Tumor Biol., 10, 316-323, 1989). monoclonal antibodies were also shown to react with ovarian tumor associated antigens, 10 however they were not specific for ovarian cancer and merely recognize determinants associated with high molecular weight mucin-like glycoproteins (Kenemans et al., Eur. J. Obstet. Gynecol. Repod. Biol. 29, 207-218, 1989; McDuffy, Ann. Clin. Biochem., 26, 379-387, 1989). More recently, oncogenes associated with ovarian cancers have been identified, including HER-2/neu (c-erbB-2) which is over-expressed in one-third of ovarian 15 cancers (USSN 6,075,122 by Cheever et al, issued June 13, 2000), the fms oncogene, and abnormalities in the p53 gene, which are seen in about half of ovarian cancers.

Whilst previously identified markers for carcinomas of the ovary have facilitated efforts to diagnose and treat these serious diseases, there is a clear need for the identification of additional markers and therapeutic targets. The identification of tumor markers that are amenable to the early-stage detection of localized tumors is critical for more effective management of carcinomas of the ovary.

### 25 Summary of the invention

20

30

35

In work leading up to the present invention, the inventors sought to identify nucleic acid markers that were diagnostic of ovarian cancers generally, or diagnostic of specific ovarian cancers such as, for example, serous ovarian cancer (SOC), mucinous ovarian cancer (MOC), non-invasive (borderline ovarian cancer or low malignant potential ovarian cancer), mixed phenotype ovarian cancer, endometrioid ovarian cancer (EnOC) and clear cell ovarian cancer (CICA), papillary serous ovarian cancer, Brenner cell or undifferentiated adenocarcinoma, by virtue of their modulated expression in cancer tissues derived from a patient cohort compared to their expression in healthy or non-cancerous cells and tissues. Additionally, the inventors sought to determine whether any correlation exists between the expression of any particular gene in a subject having ovarian cancer and the survival, or likelihood for survivial, of the subject during the

10

15

20

25

30

35

medium to long term (i.e. in the period between about 1-2 years from primary diagnosis, or longer). The inventors also sought to to determine whether any correlation exists between the expression of any particular gene in a subject following treatment for ovarian cancer and the recurrence, or likelihood for recurrence, of ovarian cancer in the subject during the medium to long term (i.e. in the period between about 1-2 years from primary diagnosis, or longer).

As exemplified herein, the inventors identified a number of genes whose expression is altered (up-regulated or down-regulated) in individuals with ovarian cancer compared to healthy individuals., eg., subjects who do not have ovarian cancer. The particular genes are identified in Tables 1 and 2. Preferably, the genes are selected from the group of candidate genes set forth in Table 3.

The list of genes and proteins exemplified herein by Table 1 were identified by a statistical analysis as outlined in the examples which gave a P-value, eg., by comparison of expression to the expression of that gene in normal ovaries.

Accordingly, one aspect of the present invention provides a method of detecting an ovarian cancer-associated transcript in a biological sample, the method comprising contacting the biological sample with a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Table 1 or 2 or 3. Preferably the percentage identity to a sequence disclosed in any one of Tables 1-3 is at least about 85% or 90% or 95%, and still more preferably at least about 98% or 99%.

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

(i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;

20

- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- 10 (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
- (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- 25 (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27,

20

25

30

- 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
- (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- Even more preferably, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
  - (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 5, 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 5, 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 5, 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 6, 8, 10, 12, 14, 16, 18, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 47, 49, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
- 35 (v) a sequence that is complementary to (i) or (ii) or (iii) or (iv).

As used herein, the term "modified level" includes an enhanced, increased or elevated level of an integer being assayed, or alternatively, a reduced or decreased level of an integer being assayed.

- In one embodiment an elevated, enhanced or increased level of expression of the nucleic acid is detected. In accordance with this embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an enhanced level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
- 25 (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200; and
- (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iii) or (iv).

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an enhanced level of hybridization of the probe for the subject

being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 10, 12, 14, 16, 18, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 47, 49, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

In an alternative preferred embodiment, a reduced level of a diagnostic marker is indicative of ovarian cancer. In accordance with this embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a reduced level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

10

- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, AI745249 and AI694200;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200;
- (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a reduced level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
  - a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, and 5;
- 25 (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, and 5;
  - (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 3, and 5;
- 30 (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, and 6; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- Preferably, the ovarian cancer that is diagnosed according to the present invention is an epithelial ovarian cancer, such as, for example, serous ovarian cancer, non-invasive

**WO** 2004/022778

ovarian cancer, mixed phenotpye ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer, clear cell ovarian cancer, papillary serous ovarian cancer, Brenner cell or undifferentiated adenocarcinoma. As will be apparent from the preferred embodiments described below, certain of the genes represented in Table 1, Table 2 and Table 3 are expressed at modified levels in subjects having serous or mucinous ovarian cancers. Data presented in Figures 1-4 also exemplify novel diagnostics and prognostics for serous ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer and clear cell ovarian cancer.

As exemplified herein by Table 2, the present inventors have identified those genes having an elevated or reduced average ratio of expression of specific genes between ovarian cancer patients vs non-ovarian cancer patients, wherein a high ratio in Table 2 indicates an enhanced expression in an ovarian cancer patients and wherein a negative ratio indicates that a reduced expression in an ovarian cancer patient.

15

20

5

In an alternative preferred embodiment, the present invention provides a method of diagnosing a serous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a serous ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- 25 (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441, X51630, and AB018305;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441, X51630, And AB018305;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441, X51630, And AB018305; and

WO 2004/022778 PCT/AU2003/001166

13

(v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

In a further alternative preferred embodiment, the present invention provides a method of diagnosing a mucinous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a mucinous ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, Al073913, Al928445, NM\_022454, W40460, AA132961 and AF111856;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, AI073913, AI928445, NM\_022454, W40460, AA132961 and AF111856;
- (iii) a sequence that is at least about 80% identical to (i) or (ii);

5

10

15

20

25

30

35

- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, AI073913, AI928445, NM 022454, W40460, AA132961 and AF111856; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

In a preferred embodiment, the present invention provides a method of diagnosing a mucinous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an enhanced level of hybridization of the probe for

the subject being tested compared to the hybridization obtained for a control subject not

having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- (i) a sequence comprising at least about 20 contiguous nucleotides from SEQ ID NO: 57 or 59 or 61:
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from SEQ ID NO: 57 or 59 or 61;
- (iii) a sequence that is at least about 80% identical to SEQ ID NO: 57 or 59 or 61;
- (iv) a sequence that encodes the amino acid sequence set forth in SEQ ID NO: 58 or 60 or 62; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

Those skilled in the art will be aware that as a carcinoma progresses, metastases occur in organs and tissues outside the site of the primary tumor. For example, in the case of ovarian cancer, metastases commonly appear in a tissue selected from the group consisting of omentum, abdominal fluid, lymph nodes, lung, liver, brain, and bone. Accordingly, the term "ovarian cancer" as used herein shall be taken to include an early or developed tumor of the ovary, such as, for example, any one or more of a number of cancers of epithelial origin, such as serous, mucinous, endometrioid, clear cell, papillary serous, Brenner cell or undifferentiated adenocarcinoma, non-invasive ovarian cancer such as borderline ovarian cancer or low-malignant potential ovarian cancer, or a mixed phenotype ovarian cancer, and optionally, any metastases outside the ovary that occurs in a subject having a primary tumor of the ovary.

25

30

20

5

10

As used herein, the term "diagnosis", and variants thereof, such as, but not limited to "diagnose", "diagnosed" or "diagnosing" shall not be limited to a primary diagnosis of a clinical state, however should be taken to include any primary diagnosis or prognosis of a clinical state. For example, the "diagnostic assay" formats described herein are equally relevant to assessing the remission of a patient, or monitoring disease recurrence, or tumor recurrence, such as following surgery or chemotherapy, or determining the appearance of metastases of a primary tumor. All such uses of the assays described herein are encompassed by the present invention.

Both classical hybridization and amplification formats, and combinations thereof, are encompassed by the invention. In one embodiment, the hybridization comprises

**WO 2004/022778** 

5

PCT/AU2003/001166

15

performing a nucleic acid hybridization reaction between a labeled probe and a second nucleic acid in the biological sample from the subject being tested, and detecting the label. In another embodiment, the hybridization comprising performing a nucleic acid amplification reaction eg., polymerase chain reaction (PCR), wherein the probe consists of a nucleic acid primer and nucleic acid copies of the nucleic acid in the biological sample are amplified. As will be known to the skilled artisan, amplification may proceed classical nucleic acid hybridization detection systems, to enhance specificity of detection, particularly in the case of less abundant mRNA species in the sample.

In a preferred embodiment, the polynucleotide is immobilised on a solid surface.

The present invention clearly encompasses nucleic acid-based methods and proteinbased methods for diagnosing cancer in humans and other mammals.

Accordingly, in a related embodiment, the present invention provides a method of detecting an ovarian cancer-associated polypeptide in a biological sample the method comprising contacting the biological sample with an antibody that binds specifically to an ovarian cancer-associated polypeptide in the biological sample, the polypeptide being encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3.

Preferably the percentage identity to a sequence disclosed in any one of Tables 1-3 is at least about 85% or 90% or 95%, and still more preferably at least about 98% or 99%.

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a modified level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34,

WO 2004/022778 PCT/AU2003/001166

36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a modified level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 6, 8, 10, 12, 14, 16, 18, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 47, 49, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.

In one embodiment an elevated, enhanced or increased level of expression of the antigen-antibody complex is detected. In accordance with this embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200.

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for

WO 2004/022778 PCT/AU2003/001166

a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.

5

10

15

20

25

30

In an alternative preferred embodiment, a reduced level of a diagnostic marker is indicative of ovarian cancer. In accordance with this embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a reduced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200.

In a preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a reduced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, and 6.

Preferably, the ovarian cancer that is diagnosed according to the present invention is an epithelial ovarian cancer, such as, for example, serous ovarian cancer or mucinous ovarian cancer.

10

15

25

30

35

In an alternative preferred embodiment, the present invention provides a method of diagnosing a serous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a modified level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a serous ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441, X51630, And AB018305.

In a further alternative preferred embodiment, the present invention provides a method of diagnosing a mucinous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a reduced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a mucinous ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, Al073913, Al928445, NM\_022454, W40460, AA132961 and AF111856.

In a preferred embodiment, the present invention provides a method of diagnosing a mucinous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a mucinous ovarian cancer, and wherein said antibody binds to a

polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to SEQ ID NO: 58 or 60 or 62.

In a further related embodiment, the present invention provides a method of detecting an ovarian cancer-associated antibody in a biological sample the method comprising contacting the biological sample with a polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3, wherein the polypeptide specifically binds to the ovarian cancer-associated antibody.

Preferably, in the above methods, the biological sample is contacted with a plurality of the polynucleotides, polypeptides or antibodies referred to above.

15 In a particularly preferred embodiment, the present invention provides an antibody-based mulptiplex assay for determing the likelihood of survivial of a subject from an ovarian cancer. In one embodiment, the invention provides a method of determining the likelihood of survival of a subject suffering from a serous ovarian cancer, said method comprising contacting a biological sample from said subject being tested with at least two 20 antibodies for a time and under conditions sufficient for antigen-antibody complexes to form and then detecting the complexes wherein an enhanced level of the antigenantibody complexes for the subject being tested compared to the amount of the antigenantibody complexes formed for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein one antibody 25 binds to an sFRP polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 72 and wherein one antibody binds to a SOCS3 polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 74.

The present invention is not to be limited by the source or nature of the biological sample. In one embodiment, the biological sample is from a patient undergoing a therapeutic regimen to treat ovarian cancer. In an alternative preferred embodiment, the biological sample is from a patient suspected of having ovarian cancer.

In addition to providing up-regulated and down-regulated genes, the list of genes and proteins exemplified herein by Table 1 were identified by a statistical analysis as outlined in the examples which gave a P-value, eg., by comparison of expression to

clinicopathological parameters for disease recurrence or patient survival. Accordingly, the present invention is particularly useful for prognostic applications, in particular for assessing the medium-to-long term survival of a subject having an ovarian cancer, or alternatively or in addition, for assessing the likelihood of disease recurrence.

5

Accordingly, a further aspect of the present invention provides a method of monitoring the efficacy of a therapeutic treatment of ovarian cancer, the method comprising:

- (i) providing a biological sample from a patient undergoing the therapeutic treatment; and
- (ii) determining the level of a ovarian cancer-associated transcript in the biological sample by contacting the biological sample with a polynucleotide that selectively hybridizes to a sequence having at least about 80% identity to a sequence as shown in any one of Tables 1-3, thereby monitoring the efficacy of the therapy.

15

10

Preferably the method further comprises comparing the level of the ovarian cancer-associated transcript to a level of the ovarian cancer-associated transcript in a biological sample from the patient prior to, or earlier in, the therapeutic treatment.

- In a related embodiment, the present invention provides a method of monitoring the efficacy of a therapeutic treatment of ovarian cancer, the method comprising:
  - (i) providing a biological sample from a patient undergoing the therapeutic treatment; and

(ii) determining the level of a ovarian cancer-associated antibody in the biological sample by contacting the biological sample with a polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at

least 80% identical to a sequence as shown in Tables 1-3, wherein the polypeptide specifically binds to the ovarian cancer-associated antibody,

thereby monitoring the efficacy of the therapy.

30

25

Preferably the method further comprises comparing the level of the ovarian cancer-associated antibody to a level of the ovarian cancer-associated antibody in a biological sample from the patient prior to, or earlier in, the therapeutic treatment.

In a further related embodiment, the present invention provides a method of monitoring the efficacy of a therapeutic treatment of ovarian cancer, the method comprising:

- (i) providing a biological sample from a patient undergoing the therapeutic treatment; and
- (ii) determining the level of a ovarian cancer-associated polypeptide in the biological sample by contacting the biological sample with an antibody, wherein the antibody specifically binds to a polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3, thereby monitoring the efficacy of the therapy.

10

5

Preferably the method further comprises comparing the level of the ovarian cancer-associated polypeptide to a level of the ovarian cancer-associated polypeptide in a biological sample from the patient prior to, or earlier in, the therapeutic treatment.

- It will also be apparent from the following preferred embodiments, that the expression of certain genes listed in Table 1 and Table 3 is statistically correlated with survival and death of patients having ovarian cancer, wherein a low P value indicates an enhanced likelihood that a patient having altered expression of the gene will die from the cancer.
- Accordingly, in one embodiment, the present invention provides a method of determining the likelihood of survival of a subject suffering from an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic
  acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, Al970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, AI796870, X02761, AW968613, AW972565, AF045229, AW953853, U52426, F06700, AI798863, H52761, BE546947, AU076643, U20536, AA581602,

AJ245210, X65965, AI806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, AI640160, U11862, AW295980, X59135, BE466173, AI354722, M90464, AA829286, AI333771, BE465867, NM\_014992, BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM\_002776, BE261944, NM\_006379, AI002238, X81789, NM\_002122, AB001914, AA311919, AI381750, AA292998, BE439580, AI677897, N72403, BE003054, AL035588, AI080491, AW770994, H24177, AF146761, NM\_001955, AI680737, AI752666, AA505445, BE246649, and NM\_003955;

- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 10 and having an Accession Number selected from the group consisting of: NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, AI970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, AI796870, X02761, 15 AW968613, AW972565, AF045229, AW953853, U52426, F06700, AI798863, H52761, BE546947, AU076643, U20536, AA581602, AJ245210, X65965, Al806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135, BE466173, Al354722, M90464, AA829286, Al333771, BE465867, NM\_014992, BE616902, AA430373, 20 R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM\_002776, BE261944, NM\_006379, Al002238, X81789, NM\_002122, AB001914, AA311919, AI381750, AA292998, BE439580, AI677897, N72403, BE003054, AL035588, Al080491, AW770994, H24177, AF146761, NM\_001955, 25 Al680737, Al752666, AA505445, BE246649, and NM 003955;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, Al970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, Al796870, X02761, AW968613, AW972565, AF045229, AW953853, U52426, F06700, Al798863, H52761, BE546947, AU076643, U20536, AA581602, AJ245210, X65965, Al806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135, BE466173, Al354722,

M90464, AA829286, AI333771, BE465867, NM\_014992, BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM\_002776, BE261944, NM\_006379, AI002238, X81789, NM\_002122, AB001914, AA311919, AI381750, AA292998, BE439580, AI677897, N72403, BE003054, AL035588, AI080491, AW770994, H24177, AF146761, NM\_001955, AI680737, AI752666, AA505445, BE246649, and NM\_003955; and

(v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

5

30

35

- In a preferred embodiment, the present invention provides a method of determining the likelihood of survival of a subject suffering from an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- 25 (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
  - (v) a sequence that is complementary to (i) or (ii) or (iii) or (iv).

In an alternative preferred embodiment, the present invention provides a method of determining the likelihood of survival of a subject suffering from an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody

10

15

20

25

30

35

complex formed for a control subject not having ovarian cancer indicates that the subject being tested has has a poor probability of survival, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, AI970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, Al796870, X02761, AW968613, AW972565. AF045229. AW953853, U52426, F06700, AI798863, H52761, BE546947, AU076643, U20536, AA581602, AJ245210, X65965, AI806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135, BE466173, Al354722, M90464, AA829286, Al333771, BE465867, NM\_014992, BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM\_002776, BE261944, NM\_006379, Al002238, X81789, NM\_002122, AB001914, AA311919, Al381750, AA292998, BE439580, Al677897, N72403, BE003054, AL035588, Al080491, AW770994, H24177, AF146761, NM\_001955, Al680737, Al752666, AA505445, BE246649, and NM\_003955.

In an alternative preferred embodiment, the present invention provides a method of determining the likelihood of survival of a subject suffering from an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has has a poor probability of survival, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.

In a particularly preferred embodiment, the present invention provides a marker for determining the likelihood of a subject surviving from serous cancer. In accordance with this embodiment of the invention, there is provided a method of determining the likelihood of survival of a subject suffering from a serous ovarian cancer, said method

20

25

30

35

comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 71 or 73;
- 10 (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 71 or 73;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii) and encoding an sFRP protein or a SOCS3 protein;
- 15 (iv) a sequence that encodes a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 72 or 74; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

In an alternative preferred embodiment, the present invention provides a method of determining the likelihood of survival of a subject suffering from a serous ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said antibody binds to an sFRP polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 72 or a SOCS3 polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 74 or.

It will also be apparent from the following preferred embodiments, that the expression of certain genes listed in Table 1 and Table 3 is statistically correlated with recurrence of ovarian cancer, wherein a low P value indicates an enhanced likelihood that a patient having altered expression of the gene will experience recurrence of the disease.

In yet another preferred embodiment, the present invention provides a method of determining the likelihood that a subject will suffer from a recurrence of an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a high probability of recurrence, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- a sequence comprising at least about 20 contiguous nucleotides from a nucleic 10 (i) acid set forth in Table 1 and having an Accession Number selected from the group consisting of: M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694, L08239, BE397649, NM\_012317, NM\_000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, 15 NM\_002514, Al863735, NM\_005397, W26391, H15474, U51166, AA243499, AW408807, Al738719, AB040888, BE313077, Al677897, C14898, Al821730, AF007393, H65423, N46243, AA095971, U20350, NM\_005756, D19589, AW957446, AW294647, BE159718, AI888490, AA022569, BE147740, AI798863, BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM\_005512, . 20 AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763, AK000596, AI970797, AW519204, Z42387, AF145713, AA972412, AK001564, AW959861, BE313555, W25005, Al193356, AF111106, Al130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, Al497778, Al745379, U51712, AW375974, 25 NM\_000636, AA130986, AA216363, AA628980, AA811657, AF251237. AA897108, AB040888, AF212225, Al089575, Al282028, Al368826, Al718702, AL109791, AW090198, AW296454, AW445034, AK002039, Al827248, AW452948, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, 30 M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051,
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions
   to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1
   and having an Accession Number selected from the group consisting of: M86849,

10

15

20

AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694, L08239, BE397649, NM 012317, NM 000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM\_002514, Al863735, NM: 005397, W26391, H15474, U51166, AA243499, AW408807. AI738719, AB040888, BE313077, AI677897, C14898, AI821730, AF007393, H65423, N46243, AA095971, U20350, NM 005756, D19589, AW957446, AW294647, BE159718, Al888490, AA022569, BE147740, Al798863, BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM 005512, AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763, AK000596, AI970797, AW519204, Z42387, AF145713, AA972412, AK001564, AW959861, BE313555, W25005, Al193356, AF111106, Al130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, Al497778, Al745379, U51712, AW375974, AF251237, NM 000636, AA130986, AA216363, AA628980, AA811657, AA897108, AB040888, AF212225, Al089575, Al282028, Al368826, Al718702, Al827248, AK002039, AL109791, AW090198, AW296454, AW445034, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051:

- (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in 25 Table 1 and having an Accession Number selected from the group consisting of:M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694, L08239, BE397649, NM\_012317, NM\_000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM 002514, Al863735, NM 005397, W26391, H15474, U51166, AA243499, 30 AW408807, Al738719, AB040888, BE313077, Al677897, C14898, Al821730, AF007393. H65423. N46243. AA095971, U20350, NM\_005756, D19589, AW957446, AW294647, BE159718, AI888490, AA022569, BE147740, AI798863, BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM\_005512, AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, 35 AA188763, AK000596, AI970797, AW519204, Z42387, AF145713, AA972412,

10

15

20

25

30

35

AK001564, AW959861, BE313555, W25005, AI193356, AF111106, AI130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, AI497778, AI745379, U51712, AW375974, AF251237, NM\_000636, AA130986, AA216363, AA628980, AA811657, AA897108, AB040888, AF212225, AI089575, AI282028, AI368826, AI718702, AI827248, AK002039, AL109791, AW090198, AW296454, AW445034, AW452948, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051; and

(v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

In an alternative preferred embodiment, the present invention provides a method of determining the likelihood that a subject will suffer from a recurrence of an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a high probability of recurrence, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694, L08239, BE397649, NM 012317, NM 000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM 002514, AI863735, NM 005397, W26391, H15474, U51166, AA243499, AW408807, AI738719, AB040888, BE313077, Al677897, C14898, Al821730, AF007393, H65423, N46243, AA095971, U20350, NM 005756, D19589, AW957446, AW294647, BE159718, AI888490, AA022569, BE147740, AI798863, BE464341, AL080235, AI557212, X75208, AA628980, BE242587, NM 005512, AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763, AK000596, Al970797, AW519204, Z42387, AF145713, AA972412, AK001564, AW959861, BE313555, W25005, AI193356,

AF111106, AI130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, AI497778, AI745379, U51712, AW375974, AF251237, NM\_000636, AA130986, AA216363, AA628980, AA811657, AA897108, AB040888, AF212225, AI089575, AI282028, AI368826, AI718702, AI827248, AK002039, AL109791, AW090198, AW296454, AW445034, AW452948, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051.

10

15

5

The recurrence of ovarian cancer is a clinical recurrence as determined by the presence of one or more clinical symptoms of an ovarian cancer, such as, for example, a metastases, or alternatively, as determined in a biochemical test, immunological test or serological test such as, for example, a cross-reactivity in a biological sample to a CA125 antibody.

Preferably, the recurrence is capable of being detected at least about 2 years from treatment, more preferably about 2-3 years from treatment, and even more preferably about 4 or 5 or 10 years from treatment.

20

25

30

35

Preferably, in the above diagnostic and/or prognostic methods, the biological sample is contacted with a plurality of the nucleic acids and/or polypeptides and/or antibodies referred to above. In a particularly preferred embodiment, mulpiplex assays are performed to detect enhanced expression at least of sFRP4 and SOC3 at the protein level (eg., using antigen-based or antibody-based assays) or at the mRNA level (eg., by detecting elevated levels of mRNA transcripts).

A further embodiment of the present invention provides a method of diagnosing epithelial ovarian cancer by detecting aberrant methylation of a promoter that regulates expression of a tumor suppressor gene eg., MCC. In particular, the present invention contemplates the detection of hypermethylation of the promoter of a tumor suppressor gene. Without being bound by any theory or mode of action, such hypermethylation leads to gene inactivation, thereby reducing expression fo the tumor suppressor gene and permitting oncogenesis. In one preferred embodiment, the present invention provides a method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising determining aberrant methylation in a promoter sequence that regulates

expression of a tumor suppressor gene in a biological sample from said subject compared to the methylation of the promoter in nucleic acid obtained for a control subject not having ovarian cancer wherein said aberrant methylation indicates that the subject being tested has an ovarian ovarian cancer.

5

10

15

20

25

30

35

In a further aspect, the present invention provides a method for identifying a compound that modulates an ovarian cancer-associated polypeptide, the method comprising:

- (i) contacting the compound with a ovarian cancer-associated polypeptide, the polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3; and
- (ii) determining the functional effect of the compound upon the polypeptide.

The functional effect may, for example, be a physical effect or a chemical effect. In one embodiment, the functional effect is determined by measuring ligand binding to the polypeptide. In a particular embodiment, the polypeptide is expressed in a eukaryotic host cell or cell membrane. Preferably the polypeptide is recombinant.

In another aspect, the present invention provides a method of inhibiting proliferation of a ovarian tumour cell, which method comprises contacting said cell with a compound identified using the method *supra* for identifying a compound that modulates an ovarian cancer-associated polypeptide.

In a further aspect, the present invention provides a method of inhibiting proliferation of a ovarian cancer-associated cell to treat ovarian cancer in a patient, the method comprising the step of administering to the patient a therapeutically effective amount of a compound identified using the method *supra* for identifying a compound that modulates an ovarian cancer-associated polypeptide.

In a further aspect, the present invention provides a drug screening assay comprising:

- (i) administering a test compound to a mammal having ovarian cancer or a cell isolated therefrom;
- (ii) comparing the level of gene expression of a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3 in a treated cell or mammal with the level of gene expression of the polynucleotide in a control cell or mammal, wherein a test compound that

WO 2004/022778 PCT/AU2003/001166

31

modulates the level of expression of the polynucleotide is a candidate for the treatment of ovarian cancer.

Typically, the control is a mammal with ovarian cancer or a cell therefrom that has not been treated with the test compound. Alternatively, the control is a normal cell or mammal.

The present invention also provides a method for treating a mammal having ovarian cancer comprising administering a compound identified the drug screening method supra.

In a further aspect, the present invention provides a pharmaceutical composition for use in treating a mammal having ovarian cancer, the composition comprising a compound identified the screening method *supra* for identifying a compound that modulates an ovarian cancer-associated polypeptide, or alternatively, using the drug screening method *supra*, and a physiologically acceptable carrier or diluent.

In a further aspect, the present invention provides an assay device, preferably for use in the diagnosis or prognosis of ovarian cancer, said device comprising a plurality of polynucleotides immobilized to a solid phase, wherein each of said polnucleotides consists of a gene as listed in any one of Tables 1-3. Preferably, the solid phase is a substantially planar chip.

In a related embodiment, the present invention provides an assay device, preferably for use in the diagnosis or prognosis of ovarian cancer, said device comprising a plurality of different antibodies immobilized to a solid phase, wherein each of said antibodies binds to a polypeptide listed in Tables 1-3. Preferably, the solid phase is a substantially planar chip.

30 Preferably, the assay device *supra* is used in a method of diagnosis or prognosis as described herein.

Alternatively, the assay device is used to identify modulatory compounds of the expression of one or more genes/proteins listed in any one of Tables 1-3.

5

10

15

20

25

In a further aspect, the present invention provides a non-human transgenic animal which is transgenic by virtue of comprising a gene set forth in any one of Tables 1-3 and, in particular, to the use of any such transgenic animal in the performance of a diagnostic or prognostic method of the invention as transgenic "knock-out" animals that have disrupted expression of a gene as set forth in any one of Tables 1-3.

In a further aspect, the present invention provides an isolated polynucleotide selected from the group consisting of:

- (a) polynucleotides comprising a nucleotide sequence as shown in Tables 1-3, or the complement thereof;
- (b) polynucleotides comprising a nucleotide sequence capable of selectively hybridizing to a nucleotide sequence as shown in Tables 1-3;
- (c) polynucleotides comprising a nucleotide sequence capable of selectively hybridizing to the complement of a nucleotide sequence as shown in Tables 1-3; and
- (d) polynucleotides comprising a polynucleotide sequence which is degenerate as a result of the genetic code to the polynucleotides defined in (a), (b) or (c).

Preferred polynucleotides comprise a polynucleotide sequence as shown in Tables 1-3 or a sequence having at least 80% homology thereto.

Preferably, the isolated polynucleotide is used for the diagnosis or prognosis of ovarian cancer, more preferably by a method as described herein. In a particularly preferred embodiment, the present invention provides for the use of a polynucleotide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.

The present invention also provides a nucleic acid vector comprising a polynucleotide of the invention. In one embodiment, the polynucleotide is operably linked to a regulatory control sequence capable of directing expression of the polynucleotide in a host cell. In a particularly preferred embodiment, the present invention provides for the use of a vector comprising a polynucleotide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.

5

10

15

20

25

30

WO 2004/022778 PCT/AU2003/001166

33

The present invention further provides a host cell comprising a vector as described in the preceding paragraph. In a particularly preferred embodiment, the present invention provides for the use of a host cell comprising an introduced polynucleotide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.

In a further aspect, the present invention provides an isolated polypeptide which is encoded by a gene set forth in any one of Tables 1-3. The present invention also provides an isolated polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3. In a particularly preferred embodiment, the present invention provides for the use of an isolated polypeptide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.

In a further aspect the present invention provides an antibody that binds specifically a polypeptide listed in Tables 1-3. In a particularly preferred embodiment, the present invention provides for the use of an antibody that binds to an isolated polypeptide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.

20

25

15

5

10

#### Brief description of the Drawings

Figure 1 is a photographic representation showing expression of genes as identified by immunohistochemical staining of fixed normal (i.e. non-cancerous or healthy) tissues (panel A) or ovarian cancer tissue (panel B). The inset in panel A shows inclusion cysts. The expression levels of the following genes listed in Table 1 or Table 3 were determined: Claudin-3 (SEQ ID NO: 15); EP-CAM (Accession No. NM\_002354); and SOX17 (SEQ ID NO: 17). Positive controls CA125, MUC-1 and E-Cadherin were also included for comparison.

10

15

20

Figure 2 is a graphical representation showing the correlation between expression of different genes in serous ovarian cancer (SOC), mucinous ovarian cancer (MOC), endometroid ovarian cancer (EnOC) and clear cell ovarian cancer (CICA). Genes indicated on the x-axis in each case are as in the legend to Figure 1.

Figure 3 is a copy of a photographic representation showing immunohistochemical staining of ovary tissue from a normal healthy subject (normal ovary), a subject diagnosed with mucinous ovarian cancer (MOC) and a subject diagnosed with serous ovarian cancer (SOC), following staining with probes that are specific for L1-Cadherin (top row), meprin alpha (middle row) or galectin-4 (lower row). Magnification is indicated as 20-40X.

Figure 4a is a copy of a photographic representation showing immunohistochemical staining of samples from a normal healthy subject (normal) or primary serous ovarian tumor (SOC), following staining with probes that are specific for sFRP4 (top row), or SOCS3 (lower row). Magnification is indicated as 20X.

Figure 4b is a copy of a graphical representation showing a Kaplan-Meier survival curve correlating sFRP4 expression to patient survival over the medium term (i.e., from about 12 months to about 48 months) to long term (more than about 48 months), indicating that high expression of sFRP4 is associated with poor survival in patients (n=127) having SOC (p=0.0056).

## 25 <u>Detailed description of the preferred embodiments</u>

Ovarian cancer-associated sequences:

Ovarian cancer-associated sequences can include both nucleic acid (i.e., "ovarian cancer-associated genes") and protein (i.e., "ovarian cancer-associated proteins").

As used herein, the term "ovarian cancer-associated protein" shall be taken to mean any protein that has an expression pattern correlated to an ovarian cancer, the recurrence of an ovarian cancer or the survival of a subject suffering from ovarian cancer.

Similarly, the term "ovarian cancer-associated gene" shall be taken to mean any nucleic acid encoding an ovarian cancer-associated protein or nucleic acid having an expression profile that is correlated to an ovarian cancer, the recurrence of an ovarian cancer or the survival of a subject suffering from ovarian cancer.

5

10

15

20

25

30

35

As will be appreciated by those in the art and is more fully outlined below, ovarian cancer-associated genes are useful in a variety of applications, including diagnostic applications, which will detect naturally occurring nucleic acids, as well as screening applications; e.g., biochips comprising nucleic acid probes or PCR microtitre plates with selected probes to the ovarian cancer sequences are generated.

For identifying ovarian cancer-associated sequences, the ovarian cancer screen typically includes comparing genes identified in different tissues, e.g., normal and cancerous tissues, or tumour tissue samples from patients who have metastatic disease vs. non metastatic tissue. Other suitable tissue comparisons include comparing ovarian cancer samples with metastatic cancer samples from other cancers, such as lung, breast, gastrointestinal cancers, ovarian, etc. Samples of different stages of ovarian cancer, e.g., survivor tissue, drug resistant states, and tissue undergoing metastasis, are applied to biochips comprising nucleic acid probes. The samples are first microdissected, if applicable, and treated as is known in the art for the preparation of mRNA. Suitable biochips are commercially available, e.g. from Affymetrix. Gene expression profiles as described herein are generated and the data analyzed.

In one embodiment, the genes showing changes in expression as between normal and disease states are compared to genes expressed in other normal tissues, preferably normal ovarian, but also including, and not limited to lung, heart, brain, liver, breast, kidney, muscle, colon, small intestine, large intestine, spleen, bone and placenta. In a preferred embodiment, those genes identified during the ovarian cancer screen that are expressed in any significant amount in other tissues are removed from the profile, although in some embodiments, this is not necessary. That is, when screening for drugs, it is usually preferable that the target be disease specific, to minimise possible side effects.

In a preferred embodiment, ovarian cancer-associated sequences are those that are upregulated in ovarian cancer; that is, the expression of these genes is modified (upregulated or down-regulated) in ovarian cancer tissue as compared to non-cancerous tissue (see Table 1).

"Up-regulation" as used herein means at least about a two-fold change, preferably at least about a three fold change, with at least about five-fold or higher being preferred. All Unigene cluster identification numbers and accession numbers herein are for the GenBank sequence database and the sequences of the accession numbers are hereby expressly incorporated by reference. Sequences are also available in other databases, e.g., European Molecular Biology Laboratory (EMBL) and DNA Database of Japan (DDBJ).

"Down-regulation" as used herein often means at least about a 1.5-fold change more preferably a two-fold change, preferably at least about a three fold change, with at least about five-fold or higher being most preferred.

15

20

25

30

35

10

Particularly preferred sequences are those referred to in Tables 1 or 3 that have a P value of less than 0.05, more preferably a P value of less than about 0.01.

Similarly, preferred sequences are those referred to in Table 2 as having an absolute ratio of expression between ovarian patients and normal patients of at least about  $\pm 5.0$ , more preferably at least about  $\pm 6.0$ , even more preferrably at least about  $\pm 7.0$  or at least about  $\pm 8.0$  or at least about  $\pm 9.0$  or at least about  $\pm 10.0$ .

Detection of ovarian cancer sequences for diagnostic/prognostic applications

In one aspect, the RNA expression levels of genes are determined for different cellular states in the ovarian cancer phenotype. Expression levels of genes in normal tissue (i.e., not undergoing ovarian cancer) and in ovarian cancer tissue (and in some cases, for varying severities of ovarian cancer that relate to prognosis, as outlined below) are evaluated to provide expression profiles. An expression profile of a particular cell state or point of development is essentially a "fingerprint" of the state. While two states may have any particular gene similarly expressed, the evaluation of a number of genes simultaneously allows the generation of a gene expression profile that is reflective of the state of the cell. By comparing expression profiles of cells in different states, information regarding which genes are important (including both up- and down-regulation of genes) in each of these states is obtained. Then, diagnosis are performed or confirmed to

37

5

10

15

20

determine whether a tissue sample has the gene expression profile of normal or cancerous tissue. This will provide for molecular diagnosis of related conditions.

"Differential expression," or grammatical equivalents as used herein, refers to qualitative or quantitative differences in the temporal and/or cellular gene expression patterns within and among cells and tissue. Thus, a differentially expressed gene can qualitatively have its expression altered, including an activation or inactivation, in, e.g., normal versus ovarian cancer tissue. Genes are turned on or turned off in a particular state, relative to another state thus permitting comparison of two or more states. A qualitatively regulated gene will exhibit an expression pattern within a state or cell type which is detectable by standard techniques. Some genes will be expressed in one state or cell type, but not in both. Alternatively, the difference in expression are quantitative, e.g., in that expression is increased or decreased; i.e., gene expression is either upregulated, resulting in an increased amount of transcript, or downregulated, resulting in a decreased amount of transcript. The degree to which expression differs need only be large enough to quantify via standard characterization techniques as outlined below, such as by use of Affymetrix GeneChipTM expression arrays, Lockhart, Nature Biotechnology 14:1675-1680 (1996), hereby expressly incorporated by reference. Other techniques include, but are not limited to, quantitative reverse transcriptase PCR, northern analysis and RNase protection. As outlined above, preferably the change in expression (i.e., upregulation or downregulation) is at least about 50%, more preferably at least about 100%, more preferably at least about 150%, more preferably at least about 200%, with from 300 to at least 1000% being especially preferred.

Evaluation are at the gene transcript, or the protein level. The amount of gene expression are monitored using nucleic acid probes to the DNA or RNA equivalent of the gene transcript, and the quantification of gene expression levels, or, alternatively, the final gene product itself (protein) are monitored, e.g., with antibodies to the ovarian cancer-associated protein and standard immunoassays (ELISAs, etc.) or other techniques, including mass spectroscopy assays, 2D gel electrophoresis assays, etc. Proteins corresponding to ovarian cancer genes, i.e., those identified as being important in a ovarian cancer phenotype, are evaluated in a ovarian cancer diagnostic test.

In a preferred embodiment, gene expression monitoring is performed on a plurality of genes. Multiple protein expression monitoring are performed as well. Similarly, these assays are performed on an individual basis as well.

In this embodiment, the ovarian cancer nucleic acid probes are attached to biochips as outlined herein for the detection and quantification of ovarian cancer sequences in a particular cell. The assays are further described below in the example. PCR techniques are used to provide greater sensitivity.

In a preferred embodiment nucleic acids encoding the ovarian cancer-associated protein are detected. Although DNA or RNA encoding the ovarian cancer-associated protein are detected, of particular interest are methods wherein an mRNA encoding a ovarian cancer-associated protein is detected. Probes to detect mRNA are nucleotide/deoxynucleotide probe that is complementary to and hybridizes with the mRNA and includes, but is not limited to, oligonucleotides, cDNA or RNA. Probes also should contain a detectable label, as defined herein. In one method the mRNA is detected after immobilizing the nucleic acid to be examined on a solid support such as nylon membranes and hybridizing the probe with the sample. Following washing to remove the non-specifically bound probe, the label is detected. In another method detection of the mRNA is performed in situ. In this method permeabilized cells or tissue samples are contacted with a detectably labeled nucleic acid probe for sufficient time to allow the probe to hybridize with the target mRNA. Following washing to remove the non-specifically bound probe, the label is detected. For example a digoxygenin labeled riboprobe (RNA probe) that is complementary to the mRNA encoding a ovarian cancerassociated protein is detected by binding the digoxygenin with an anti-digoxygenin secondary antibody and developed with nitro blue tetrazolium and 5-bromo-4-chloro-3indoyl phosphate.

25

30

5

10

15

20

In a preferred embodiment, various proteins from the three classes of proteins as described herein (secreted, transmembrane or intracellular proteins) are used in diagnostic assays. The ovarian cancer-associated proteins, antibodies, nucleic acids, modified proteins and cells containing ovarian cancer sequences are used in diagnostic assays. This are performed on an individual gene or corresponding polypeptide level. In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes and/or corresponding polypeptides.

35 As described and defined herein, ovarian cancer-associated proteins, including intracellular, transmembrane or secreted proteins, find use as markers of ovarian cancer.

Detection of these proteins in putative ovarian cancer tissue allows for detection or diagnosis of ovarian cancer. In one embodiment, antibodies are used to detect ovarian cancer-associated proteins. A preferred method separates proteins from a sample by electrophoresis on a gel (typically a denaturing and reducing protein gel, but are another type of gel, including isoelectric focusing gels and the like). Following separation of proteins, the ovarian cancer-associated protein is detected, e.g., by immunoblotting with antibodies raised against the ovarian cancer-associated protein. Methods of immunoblotting are well known to those of ordinary skill in the art.

10

15

20

25

30

In another preferred method, antibodies to the ovarian cancer-associated protein find use in *in situ* imaging techniques, e.g., in histology (e.g., *Methods in Cell Biology: Antibodies in Cell* Biology, volume 37 (Asai, ed. 1993)). In this method cells are contacted with from one to many antibodies to the ovarian cancer-associated protein(s). Following washing to remove non-specific antibody binding, the presence of the antibody or antibodies is detected. In one embodiment the antibody is detected by incubating with a secondary antibody that contains a detectable label. In another method the primary antibody to the ovarian cancer-associated proteins) contains a detectable label, e.g. an enzyme marker that can act on a substrate. In another preferred embodiment each one of multiple primary antibodies contains a distinct and detectable label. This method finds particular use in simultaneous screening for a plurality of ovarian cancer-associated proteins. As will be appreciated by one of ordinary skill in the art, many other histological imaging techniques are also provided by the invention.

in a preferred embodiment the label is detected in a fluorometer which has the ability to detect and distinguish emissions of different wavelengths. In addition, a fluorescence activated cell sorter (FACS) are used in the method. In another preferred embodiment, antibodies find use in diagnosing ovarian cancer from blood, serum, plasma, stool, and other samples. Such samples, therefore, are useful as samples to be probed or tested for the presence of ovarian cancer-associated proteins. Antibodies are used to detect a ovarian cancer-associated protein by previously described immunoassay techniques including ELISA, immunoblotting (western blotting), immunoprecipitation, BIACORE technology and the like. Conversely, the presence of antibodies may indicate an immune response against an endogenous ovarian cancer-associated protein.

In a preferred embodiment, in situ hybridization of labeled ovarian cancer nucleic acid probes to tissue arrays is done. For example, arrays of tissue samples, including ovarian

. 40

cancer tissue and/or normal tissue, are made. In situ hybridization (see, e.g., Ausubel, supra) is then performed. When comparing the fingerprints between an individual and a standard, the skilled artisan can make a diagnosis, a prognosis, or a prediction based on the findings. It is further understood that the genes which indicate the diagnosis may differ from those which indicate the prognosis and molecular profiling of the condition of the cells may lead to distinctions between responsive or refractory conditions or are predictive of outcomes.

5

10

15

20

25

30

35

In a preferred embodiment, the ovarian cancer-associated proteins, antibodies, nucleic acids, modified proteins and cells containing ovarian cancer sequences are used in prognosis assays. As above, gene expression profiles are generated that correlate to ovarian cancer, in terms of long term prognosis. Again, this are done on either a protein or gene level, with the use of genes being preferred. As above, ovarian cancer probes are attached to biochips for the detection and quantification of ovarian cancer sequences in a tissue or patient. The assays proceed as outlined above for diagnosis. PCR method may provide more sensitive and accurate quantification.

Characteristics of ovarian cancer-associated proteins and genes encoding same

Ovarian cancer-associated proteins of the present invention are classified as secreted proteins, transmembrane proteins or intracellular proteins. In one embodiment, the ovarian cancer-associated protein is an intracellular protein. Intracellular proteins are found in the cytoplasm and/or in the nucleus. Intracellular proteins are involved in all aspects of cellular function and replication (including, e.g., signaling pathways); aberrant expression of such proteins often results in unregulated or disregulated cellular processes (see, e.g., Molecular Biology of the Cell (Alberts, ed., 3rd ed., 1994). For example, many intracellular proteins have enzymatic activity such as protein kinase activity, protein phosphatase activity, protease activity, nucleotide cyclase activity, polymerase activity and the like. Intracellular proteins also serve as docking proteins that are involved in organizing complexes of proteins, or targeting proteins to various subcellular localizations, and are involved in maintaining the structural integrity of organelles.

An increasingly appreciated concept in characterising proteins is the presence in the proteins of one or more motifs for which defined functions have been attributed. In addition to the highly conserved sequences found in the enzymatic domain of proteins, highly conserved sequences have been identified in proteins that are involved in

41

protein-protein interaction. For example, Src-homology-2 (SH2) domains bind tyrosine-phosphorylated targets in a sequence dependent manner. PTB domains, which are distinct from SH2 domains, also bind tyrosine phosphorylated targets, SH3 domains bind to proline-rich targets. In addition, PH domains, tetratricopeptide repeats and WD domains to name only a few, have been shown to mediate protein-protein interactions. Some of these may also be involved in binding to phospholipids or other second messengers. As will be appreciated by one of ordinary skill in the art, these motifs are identified on the basis of primary sequence; thus, an analysis of the sequence of proteins may provide insight into both the enzymatic potential of the molecule and/or molecules with which the protein may associate. One useful database is Pfam (protein families), which is a large collection of multiple sequence alignments and hidden Markov models covering many common protein domains. Versions are available via the internet from Washington University in St. Louis, the Sanger Center in England, and the Karolinska Institute in Sweden (see, e.g., Bateman et al., 2000, Nuc. Acids Res. 28: 263-266; Sonnhammer et al., 1997, Proteins 28: 405-420; Bateman et al., 1999, Nuc. Acids Res. 27:260-262; and Sonnhammer et al., 1998, Nuc. Acids Res. 26: 320-322.

10

15

20

25

30

35

In another embodiment, the ovarian cancer sequences are transmembrane proteins. Transmembrane proteins are molecules that span a phospholipid bilayer of a cell. They may have an intracellular domain, an extracellular domain, or both. The intracellular domains of such proteins may have a number of functions including those already described for intracellular proteins. For example, the intracellular domain may have enzymatic activity and/or may serve as a binding site for additional proteins. Frequently the intracellular domain of transmembrane proteins serves both roles. For example certain receptor tyrosine kinases have both protein kinase activity and SH2 domains. In addition, autophosphorylation of tyrosines on the receptor molecule itself, creates binding sites for additional SH2 domain containing proteins.

Transmembrane proteins may contain from one to many transmembrane domains. For example, receptor tyrosine kinases, certain cytokine receptors, receptor guanylyl cyclases and receptor serine/threonine protein kinases contain a single transmembrane domain. However, various other proteins including channels and adenylyl cyclases contain numerous transmembrane domains. Many important cell surface receptors such as G protein coupled receptors (GPCRs) are classified as "seven transmembrane domain" proteins, as they contain 7 membrane spanning regions. Characteristics of transmembrane domains include approximately 20 consecutive hydrophobic amino acids

42

that are followed by charged amino acids. Therefore, upon analysis of the amino acid sequence of a particular protein, the localization and number of transmembrane domains within the protein are predicted (see, e.g. PSORT web site http://psort.nibb.ac.jp/). Important transmembrane protein receptors include, but are not limited to the insulin receptor, insulin-like growth factor receptor, human growth hormone receptor, glucose transporters, transferrin receptor, epidermal growth factor receptor, low density lipoprotein receptor, epidermal growth factor receptor, leptin receptor, interleukin receptors, e.g. IL-1 receptor, IL-2 receptor,

The extracellular domains of transmembrane proteins are diverse; however, conserved 10 motifs are found repeatedly among various extracellular domains. Conserved structure and/or functions have been ascribed to different extracellular motifs. Many extracellular domains are involved in binding to other molecules. In one aspect, extracellular domains are found on receptors. Factors that bind the receptor domain include circulating ligands, 15 which are peptides, proteins, or small molecules such as adenosine and the like. For example, growth factors such as EGF, FGF and PDGF are circulating growth factors that bind to their cognate receptors to initiate a variety of cellular responses. Other factors include cytokines, mitogenic factors, neurotrophic factors and the like. Extracellular domains also bind to cell-associated molecules. In this respect, they mediate cell-cell 20 interactions., Cell-associated ligands are tethered to the cell, e.g., via a glycosylphosphatidylinositol (GPI) anchor, or may themselves be transmembrane proteins. Extracellular domains also associate with the extracellular matrix and contribute to the maintenance of the cell structure.

Ovarian cancer-associated proteins that are transmembrane are particularly preferred in the present invention as they are readily accessible targets for immunotherapeutics, as are described herein. In addition, as outlined below, transmembrane proteins are also useful in imaging modalities. Antibodies are used to label such readily accessible proteins *in situ*. Alternatively, antibodies can also label intracellular proteins, in which case samples are typically permeablized to provide access to intracellular proteins.

It will also be appreciated by those in the art that a transmembrane protein are made soluble by removing transmembrane sequences, e.g., through recombinant methods. Furthermore, transmembrane proteins that have been made soluble are made to be secreted through recombinant means by adding an appropriate signal sequence.

35

In another embodiment, the ovarian cancer-associated proteins are secreted proteins; the secretion of which are either constitutive or regulated. These proteins have a signal peptide or signal sequence that targets the molecule to the secretory pathway. Secreted proteins are involved in numerous physiological events; by virtue of their circulating nature, they serve to transmit signals to various other cell types. The secreted protein may function in an autocrine manner (acting on the cell that secreted the factor), a paracrine manner (acting on cells in close proximity to the cell that secreted the factor) or an endocrine manner (acting on cells at a distance). Thus secreted molecules find use in modulating or altering numerous aspects of physiology. Ovarian cancer-associated proteins that are secreted proteins are particularly preferred in the present invention as they serve as good targets for diagnostic markers, e.g., for blood, plasma, serum, or stool tests.

## Mammalian subjects

5

10

15

20

25

30

35

The present invention provides nucleic acid and protein sequences that are differentially expressed in ovarian cancer, herein termed "ovarian cancer sequences." As outlined below, ovarian cancer sequences include those that are up-regulated (i.e., expressed at a higher level) in ovarian cancer, as well as those that are down-regulated (i.e., expressed at a lower level). In a preferred embodiment, the ovarian cancer sequences are from humans; however, as will be appreciated by those in the art, ovarian cancer sequences from other organisms are useful in animal models of disease and drug evaluation; thus, other ovarian cancer sequences are provided, from vertebrates, including mammals, including rodents (rats, mice, hamsters, guinea pigs, etc.), primates, farm animals (including sheep, goats, pigs, cows, horses, etc.) and pets, e.g., (dogs, cats, etc.).

#### Assay control samples

It will be apparent from the preceding discussion that many of the diagnostic methods provided by the present invention involve a degree of quantification to determine, on the one hand, the over-expression or reduced-expression of a diagnostic/prognostic marker in tissue that is suspected of comprising a cancer cell. Such quantification can be readily provided by the inclusion of appropriate control samples in the assays described below, derived from healthy or normal individuals. Alternatively, if internal controls are not included in each assay conducted, the control may be derived from an established data set that has been generated from healthy or normal individuals.

In the present context, the term "healthy individual" shall be taken to mean an individual who is known not to suffer from ovarian cancer, such knowledge being derived from clinical data on the individual, including, but not limited to, a different cancer assay to that described herein. As the present invention is particularly useful for the early detection of ovarian cancer, it is preferred that the healthy individual is asymptomatic with respect to the early symptoms associated with ovarian cancer. Although early detection using well-known procedures is difficult, reduced urinary frequency, rectal pressure, and abdominal bloating and swelling, are associated with the disease in its early stages, and, as a consequence, healthy individuals should not have any of these clinical symptoms. Clearly, subjects suffering from later symptoms associated with ovarian cancer, such as, for example, metastases in the omentum, abdominal fluid, lymph nodes, lung, liver, brain, or bone, and subjects suffering from spinal cord compression, elevated calcium level, chronic pain, or pleural effusion, should also be avoided from the "healthy individual" data set.

15

20

25

30

35

10

The term "normal individual" shall be taken to mean an individual having a normal level of expression of a cancer-associate gene or cancer-associated protein in a particular sample derived from said individual. As will be known to those skilled in the art, data obtained from a sufficiently large sample of the population will normalize, allowing the generation of a data set for determining the average level of a particular parameter. Accordingly, the level of expression of a cancer-associate gene or cancer-associated protein can be determined for any population of individuals, and for any sample derived from said individual, for subsequent comparison to levels determined for a sample being assayed. Where such normalized data sets are relied upon, internal controls are preferably included in each assay conducted to control for variation.

In one embodiment, the present invention provides a method for detecting a cancer cell in a subject, said method comprising:

- (i) determining the level of mRNA encoding a cancer-associated protein expressed in a test sample from said subject; and
- comparing the level of mRNA determined at (i) to the level of mRNA encoding a cancer-associated protein expressed in a comparable sample from a healthy or normal individual,

wherein a level of mRNA at (i) that is modified in the test sample relative to the comparable sample from the normal or healthy individual is indicative of the presence of a cancer cell in said subject.

Alternatively, or in addition, the controll may comprise a cancer-associated sequence that is known to be expressed at a particular level in an ovarian cancer, eg., CA125, MUC-1 or E-Cadherin, amongast others.

5

10

## Biological samples

Preferred biological samples in which the assays of the invention are performed include bodily fluids, ovarian tissue and cells, and those tissues known to comprise cancer cells arising from a metastasis of an ovarian cancer, such as, for example, in carcinomas of the lung, prostate, breast, colon, pancreas, placenta, or omentum, and in cells of brain anaplastic oligodendrogliomas.

Bodily fluids shall be taken to include whole blood, serum, peripheral blood mononuclear cells (PBMC), or buffy coat fraction.

. 15

In the present context, the term "cancer cell" includes any biological specimen or sample comprising a cancer cell irrespective of its degree of isolation or purity, such as, for example, tissues, organs, cell lines, bodily fluids, or histology specimens that comprise a cell in the early stages of transformation or having been transformed.

20

25

30

35

As the present invention is particularly useful for the early detection and prognosis of cancer ofe rthe medium to long term, the definition of "cancer cell" is not to be limited by the stage of a cancer in the subject from which said cancer cell is derived (ie. whether or not the patient is in remission or undergoing disease recurrence or whether or not the cancer is a primary tumor or the consequence of metastases). Nor is the term "cancer cell" to be limited by the stage of the cell cycle of said cancer cell.

Preferably, the sample comprises ovarian tissue, prostate tissue, kidney tissue, uterine tissue, placenta, a cervical specimen, omentum, rectal tissue, brain tissue, bone tissue, lung tissue, lymphatic tissue, urine, semen, blood, abdominal fluid, or serum, or a cell preparation or nucleic acid preparation derived therefrom. More preferably, the sample comprises serum or abdominal fluid, or a tissue selected from the group consisting of: ovary, lymph, lung, liver, brain, placenta, brain, omentum, and prostate. Even more preferably, the sample comprises serum or abdominal fluid, ovary (eg. OSE), or lymph node tissue. The sample can be prepared on a solid matrix for histological analyses, or alternatively, in a suitable solution such as, for example, an extraction buffer or

46

suspension buffer, and the present invention clearly extends to the testing of biological solutions thus prepared.

Polynucleotide probes and amplification primers

Polynucleotide probes are derived from or comprise the nucleic acid sequences whose nucleotide sequences are provided by reference to the public database accession numbers given in Tables 1-3 (referred to herein as the nucleotide sequences shown in Tables 1-3), and sequences homologues thereto as well as variants, derivatives and fragments thereof.

10

15

20

25

30

35

5

Whilst the probes may comprise double-stranded or single-stranded nucleic acid, single-stranded probes are preferred because they do not require melting prior to use in hybridizations. On the other hand, longer probes are also preferred because they can be used at higher hybridization stringency than shorter probes and may produce lower background hybridization than shorter probes.

So far as shorter probes are concerned, single-stranded, chemically-synthesized oligonucleotide probes are particularly preferred by the present invention. To reduce the noise associated with the use of such probes during hybridization, the nucleotide sequence of the probe is carefully selected to maximize the Tm at which hybridizations can be performed, reduce non-specific hybridization, and to reduce self-hybridization. Such considerations may be particularly important for applications involving high throughput screening using microarray technology. In general, this means that the nucleotide sequence of an oligonucleotide probe is selected such that it is unique to the target RNA or protein-encoding sequence, has a low propensity to form secondary structure, low self-complementary, and is not highly A/T-rich.

The only requirement for the probes is that they cross-hybridize to nucleic acid encoding the target diagnostic protein or the complementary nucleotide sequence thereto and are sufficiently unique in sequence to generate high signal:noise ratios under specified hybridization conditions. As will be known to those skilled in the art, long nucleic acid probes are preferred because they tend to generate higher signal:noise ratios than shorter probes and/or the duplexes formed between longer molecules have higher melting temperatures (i.e. Tm values) than duplexes involving short probes. Accordingly, full-length DNA or RNA probes are contemplated by the present invention, as are specific probes comprising the sequence of the 3'-untranslated region or complementary thereto.

WO 2004/022778

PCT/AU2003/001166

47

In a particularly preferred embodiment, the nucleotide sequence of an oligonucleotide probe has no detectable nucleotide sequence identity to a nucleotide sequence in a BLAST search (Altschul *et al.*, *J. Mol. Biol. 215*, 403-410, 1990) or other database search, other than a sequence selected from the group consisting of: (a) a sequence encoding a polypeptide listed in any one of Tables 1-3; (b) the 5'-untranslated region of a sequence encoding a polypeptide listed in any one of Tables 1-3; (c) a 3'-untranslated region of a sequence encoding a polypeptide listed in any one of Tables 1-3; and (d) an exon region of a sequence encoding a polypeptide listed in any one of Tables 1-3.

10

15

20

25

30

35

Additionally, the self-complementarity of a nucleotide sequence can be determined by aligning the sequence with its reverse complement, wherein detectable regions of identity are indicative of potential self-complementarity. As will be known to those skilled in the art, such sequences may not necessarily form secondary structures during hybridization reaction, and, as a consequence, successfully identify a target nucleotide sequence. It is also known to those skilled in the art that, even where a sequence does form secondary structures during hybridization reactions, reaction conditions can be modified to reduce the adverse consequences of such structure formation. Accordingly, a potential for self-complementarity should not necessarily exclude a particular candidate oligonucleotide from selection. In cases where it is difficult to determine nucleotide sequences having no potential self-complementarity, the uniqueness of the sequence should outweigh a consideration of its potential for secondary structure formation.

Recommended pre-requisites for selecting oligonucleotide probes, particularly with respect to probes suitable for microarray technology, are described in detail by Lockhart et al., "Expression monitoring by hybridization to high-density oligonucleotide arrays", Nature Biotech. 14, 1675-1680, 1996.

The nucleic acid probe may comprise a nucleotide sequence that is within the coding strand of a gene listed in any one of Tables 1-3. Such "sense" probes are useful for detecting RNA by amplification procedures, such as, for example, polymerase chain reaction (PCR), and more preferably, quantitative PCR or reverse transcription polymerase chain reaction (RT-PCR). Alternatively, "sense" probes may be expressed to produce polypeptides or immunologically active derivatives thereof that are useful for detecting the expressed protein in samples.

30

35

The nucleotide sequences referred to in Tables 1-3 and homologues thereof, typically encode polypeptides. It will be understood by a skilled person that numerous different polynucleotides can encode the same polypeptide as a result of the degeneracy of the genetic code. In addition, it is to be understood that skilled persons may, using routine techniques, make nucleotide substitutions that do not affect the polypeptide sequence encoded by the polynucleotides of the invention to reflect the codon usage of any particular host organism in which the polypeptides of the invention are to be expressed.

Polynucleotides may comprise DNA or RNA. They are single-stranded or double-stranded. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothicate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the polynucleotides described herein are modified by any method available in the art. Such modifications are carried out in order to enhance the *in vivo* activity or life span of the diagnostic/prognostic polynucleotides.

The terms "variant" or "derivative" in relation to the nucleotide sequences of the present invention include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the sequence provided that the resultant nucleotide sequence codes for a polypeptide having biological activity, preferably having substantially the same activity as the polypeptide sequences presented in the sequence listings.

With respect to sequence homology, preferably there is at least 75%, more preferably at least 85%, more preferably at least 90% homology to a sequence shown in Tables 1-3 herein over a region of at least 20, preferably at least 25 or 30, for instance at least 40, 60, 100, 500, 1000 or more contiguous nucleotides. More preferably there is at least 95%, more preferably at least 98%, homology. In one embodiment, homologues are naturally occurring sequences, such as orthologues, tissue-specific isoforms and allelic variants.

Homology comparisons are conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences.

5

10

15

20

25

30

35

Percentage (%) homology are calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each nucleotide in one sequence directly compared with the corresponding nucleotide in the other sequence, one base at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short number of bases (for example less than 50 contiguous nucleotides).

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following nucleotides to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting "gaps" in the sequence alignment to try to maximise local homology.

However, these more complex methods assign "gap penalties" to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. "Affine gap costs" are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons.

In determining whether or not two amino acid sequences fall within the stated defined percentage identity limits, those skilled in the art will be aware that it is necessary to conduct a side-by-side comparison of amino acid sequences. In such comparisons or alignments, differences will arise in the positioning of non-identical amino acid residues depending upon the algorithm used to perform the alignment. In the present context, references to percentage identities and similarities between two or more amino acid sequences shall be taken to refer to the number of identical and similar residues respectively, between said sequences as determined using any standard algorithm known to those skilled in the art. In particular, amino acid identities and similarities are calculated using the GAP program of the Computer Genetics Group, Inc., University

WO 2004/022778

Research Park, Madison, Wisconsin, United States of America (Devereaux et al, Nucl. Acids Res. 12, 387-395,1984), which utilizes the algorithm of Needleman and Wunsch J. Mol. Biol. 48, 443-453, 1970, or alternatively, the CLUSTAL W algorithm of Thompson et al., Nucl. Acids Res. 22, 4673-4680, 1994, for multiple alignments, to maximize the number of identical/similar amino acids and to minimize the number and/or length of sequence gaps in the alignment.

A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux *et al.*, 1984, Nucleic Acids Research 12:387). The default scoring matrix has a match value of 10 for each identical nucleotide and -9 for each mismatch. The default gap creation penalty is -50 and the default gap extension penalty is -3 for each nucleotide.

Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.*, 1999 *ibid* – Chapter 18), FASTA (Atschul *et al.*, 1990, J. Mol. Biol., 403-410) and the GENEWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel *et al.*, 1999 *ibid*, pages 7-58 to 7-60). However it is preferred to use the GCG Bestfit program.

20

30

35

5

10

15

Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

A preferred sequence comparison program is the GCG Wisconsin Bestfit program described above.

The present invention also encompasses the use of nucleotide sequences that are capable of hybridizing selectively to the sequences presented herein, or any variant, fragment or derivative thereof, or to the complement of any of the above. Nucleotide sequences are preferably at least 15 nucleotides in length, more preferably at least 20, 30, 40 or 50 nucleotides in length.

The term "hybridization" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction technologies.

Polynucleotides capable of selectively hybridizing to the nucleotide sequences presented herein, or to their complement, will be generally at least 70%, preferably at least 80 or 90% and more preferably at least 95% or 98% homologous to the corresponding nucleotide sequences referred to in Tables 1-3 over a region of at least 20, preferably at least 25 or 30, for instance at least 40, 60, 100, 500, 1000 or more contiguous nucleotides.

The term "selectively hybridizable" means that the polynucleotide used as a probe is used under conditions where a target polynucleotide is found to hybridize to the probe at a level significantly above background. The background hybridization may occur because of other polynucleotides present, for example, in the cDNA or genomic DNA library being screening. In this event, background implies a level of signal generated by interaction between the probe and a non-specific DNA member of the library which is less than 10 fold, preferably less than 100 fold as intense as the specific interaction observed with the target DNA. The intensity of interaction are measured, for example, by radiolabelling the probe, e.g. with <sup>32</sup>P.

Hybridization conditions are based on the melting temperature (Tm) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA), and confer a defined "stringency" as explained below.

For the purposes of defining the level of stringency, a high stringency hybridization is achieved using a hybridization buffer and/or a wash solution comprising the following:

- (i) a salt concentration that is equivalent to 0.1xSSC-0.2xSSC buffer or lower salt concentration;
  - (ii) a detergent concentration equivalent to 0.1% (w/v) SDS or higher; and
  - (iii) an incubation temperature of 55°C or higher.

Conditions for specifically hybridizing nucleic acid, and conditions for washing to remove non-specific hybridizing nucleic acid, are well understood by those skilled in the art. For the purposes of further clarification only, reference to the parameters affecting hybridization between nucleic acid molecules is found in Ausubel *et al.* (Current Protocols in Molecular Biology, Wiley Interscience, ISBN 047150338, 1992), which is herein incorporated by reference.

5

10

15

20

25

30

5

15

20

25

30

35

Maximum stringency typically occurs at about Tm-5°C (5°C below the Tm of the probe); high stringency at about 5°C to 10°C below Tm; intermediate stringency at about 10°C to 20°C below Tm; and low stringency at about 20°C to 25°C below Tm. As will be understood by those of skill in the art, a maximum stringency hybridization are used to identify or detect identical polynucleotide sequences while an intermediate (or low) stringency hybridization are used to identify or detect similar or related polynucleotide sequences.

In a preferred aspect, the present invention covers nucleotide sequences that can hybridize to the nucleotide sequence of the present invention under stringent conditions (e.g. 65°C and 0.1xSSC {1xSSC = 0.15 M NaCl, 0.015 M Na<sub>3</sub> Citrate pH 7.0}).

Where the diagnostic/prognostic polynucleotide is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the present invention. Where the polynucleotide is single-stranded, it is to be understood that the complementary sequence of that polynucleotide is also included within the scope of the present invention.

Polynucleotides which are not 100% homologous to the sequences of the present invention but are useful in perfoming the diagnostic and/or prognostic assays of the invertion by virtue of their ability to selectively hybridize to the target gene transcript, or to encode an immunologically cross-reactive protein to the target protein, are obtained in a number of ways, such as, for example by probing DNA libraries made from a range of individuals, for example individuals from different populations. In particular, given that that changes in the expression of diagnostic/prognostic cancer-associated genes correlate with ovarian cancer, characterisation of variant sequences in individuals suffering from ovarian cancer is used to identify variations in the sequences of ovarian-cancer associated genes (and proteins) that are predictive of and/or causative of ovarian cancer.

Accordingly the present invention provides methods of identifying sequence variants that are associated with ovarian cancer which methods comprise determining all or part of the nucleotide sequence of a gene referred to in Tables 1-3, derived from an individual suffering from ovarian cancer and comparing the sequence to that of the corresponding wild-type gene.

In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in mammalian cells (e.g. rat, mouse, bovine and primate cells), are obtained and such

53

homologues and fragments thereof in general will be capable of selectively hybridizing to the sequences shown in the sequence listing herein. Such sequences are obtained by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of the sequences referred to in Tables 1-3 under conditions of medium to high stringency. Similar considerations apply to obtaining species homologues and allelic variants of the nucleotide sequences referred to in Tables 1-3.

Variants and strain/species homologues may also be obtained using degenerate PCR which will use primers designed to target sequences within the variants and homologues encoding conserved amino acid sequences within the sequences of the present invention. Conserved sequences are predicted, for example, by aligning the amino acid sequences from several variants/homologues. Sequence alignments are performed using computer software known in the art. For example the GCG Wisconsin PileUp program is widely used.

15

20

25

30

5

10

The primers used in degenerate PCR will contain one or more degenerate positions and will be used at stringency conditions lower than those used for cloning sequences with single sequence primers against known sequences.

Alternatively, such polynucleotides are obtained by site-directed mutagenesis of characterised sequences, such as the sequences referred to in Tables 1-3. This are useful where for example silent codon changes are required to sequences to optimise codon preferences for a particular host cell in which the polynucleotide sequences are being expressed. Other sequence changes are desired in order to introduce restriction enzyme recognition sites, or to alter the property or function of the polypeptides encoded by the polynucleotides.

Polynucleotides comprising a diagnostic/prognostic cancer-associated gene are used to produce a primer by standard derivatization means, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a detectable label by conventional means using radioactive or non-radioactive labels, or the polynucleotides are cloned into vectors. Such primers, probes and other fragments will be at least 15, preferably at least 20, for example at least 25, 30 or 40 nucleotides in length. Preferred fragments are less than 5000, 2000, 1000, 500 or 200 nucleotides in length.

Polynucleotides such as a DNA polynucleotides and probes according to the invention are produced by recombinant or synthetic means, including cloning by standard techniques.

In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

5

10

15

20

25

30

35

Longer polynucleotides will generally be produced using recombinant means, for example using PCR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking a region of the sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers are designed to contain suitable restriction enzyme recognition sites so that the amplified DNA are cloned into a suitable cloning vector

Polynucleotide probes or primers preferably carry a detectable label. Suitable labels include radioisotopes such as <sup>32</sup>P or <sup>35</sup>S, enzyme labels, or other protein labels such as biotin. Such labels are added to polynucleotides or primers and are detected using by techniques known in the art.

Polynucleotide probes or primers labeled or unlabeled are also used by a person skilled in the art in nucleic acid-based tests for detecting or sequencing diagnostic/prognostic cancer-associated gene.

Such tests for detecting generally comprise bringing a biological sample containing DNA or RNA into contact with a probe comprising a polynucleotide probe or primer under at least low stringency hybridization conditions and detecting any duplex formed between the probe/primer and nucleic acid in the sample. Such detection are achieved using techniques such as PCR or by immobilising the probe on a solid support, removing nucleic acid in the sample which is not hybridized to the probe, and then detecting nucleic acid which has hybridized to the probe. Alternatively, the sample nucleic acid are immobilised on a solid support, and the amount of probe bound to such a support are detected. Suitable assay methods of this and other formats are found in for example W089/03891 and W090/13667.

WO 2004/022778

Tests for sequencing nucleotides include bringing a biological sample containing target DNA or RNA into contact with a probe comprising a polynucleotide probe or primer under at least low stringency hybridization conditions and determining the sequence by, for example the Sanger dideoxy chain termination method (see Sambrook et al.).

Such a method generally comprises elongating, in the presence of suitable reagents, the primer by synthesis of a strand complementary to the target DNA or RNA and selectively terminating the elongation reaction at one or more of an A, C, G or T/U residue; allowing strand elongation and termination reaction to occur; separating out according to size the elongated products to determine the sequence of the nucleotides at which selective termination has occurred. Suitable reagents include a DNA polymerase enzyme, the deoxynucleotides dATP, dCTP, dGTP and dTTP, a buffer and ATP. Dideoxynucleotides are used for selective termination.

15

10

5

Tests for detecting or sequencing nucleotides in a biological sample are used as part of the methods of the invention for detecting ovarian cancer-associated transcripts and monitoring the efficacy of treatment of patients suffering from ovarian cancer as described in more detail herein.

20

25

30

35

The probes/primers may conveniently be packaged in the form of a test kit in a suitable container. In such kits the probe are bound to a solid support where the assay format for which the kit is designed requires such binding. The kit may also contain suitable reagents for treating the sample to be probed, hybridizing the probe to nucleic acid in the sample, control reagents, instructions, and the like.

Preferably, a kit of the invention comprises primers/probes suitable for selectively detecting a plurality of sequences, more preferably for selectively detecting a plurality of sequences that are listed in one or more of Tables 1-3 as having a P value of less than 0.05, more preferably a P value of less than 0.01. Similarly, a kit of the invention preferably comprises primers suitable for selectively detecting a plurality of sequences referred to in Table 1 or 2 or 3.

### Nucleic acid-based assay formats

As discussed in detail below, the status of expression of a cancer-associated gene in patient samples may be analyzed by a variety protocols that are well known in the art

including *in situ* hybridization, northern blotting techniques, RT-PCR analysis (such as, for example, performed on laser capture microdissected samples), and microarray technology, such as, for example, using tissue microarrays probed with nucleic acid probes, or nucleic acid microarrays (ie. RNA microarrays or amplified DNA microarrays) microarrays probed with nucleic acid probes. All such assay formats are encompassed by the present invention.

5

10

15

20

25

30

35

For high throughput screening of large numbers of samples, such as, for example, public health screening of subjects, particularly human subjects, having a higher risk of developing cancer, microarray technology is a preferred assay format.

In accordance with such high throughput formats, techniques for producing immobilised arrays of DNA molecules have been described in the art. Generally, most prior art methods describe how to synthesise single-stranded nucleic acid molecule arrays, using for example masking techniques to build up various permutations of sequences at the various discrete positions on the solid substrate. U.S. Patent No. 5,837,832, the contents of which are incorporated herein by reference, describes an improved method for producing DNA arrays immobilised to silicon substrates based on very large scale integration technology. In particular, U.S. Patent No. 5,837,832 describes a strategy called "tiling" to synthesize specific sets of probes at spatially-defined locations on a substrate which are used to produced the immobilised DNA arrays. U.S. Patent No. 5,837,832 also provides references for earlier techniques that may also be used.

Thus DNA are synthesised *in situ* on the surface of the substrate. However, DNA may also be printed directly onto the substrate using for example robotic devices equipped with either pins or piezo electric devices.

The plurality of polynucleotide sequences are typically immobilised onto or in discrete regions of a solid substrate. The substrate are porous to allow immobilisation within the substrate or substantially non-porous, in which case the library sequences are typically immobilised on the surface of the substrate. The solid substrate are made of any material to which polypeptides can bind, either directly or indirectly. Examples of suitable solid substrates include flat glass, silicon wafers, mica, ceramics and organic polymers such as plastics, including polystyrene and polymethacrylate. It may also be possible to use semi-permeable membranes such as nitrocellulose or nylon membranes, which are widely available. The semi-permeable membranes are mounted on a more robust solid

surface such as glass. The surfaces may optionally be coated with a layer of metal, such as gold, platinum or other transition metal. A particular example of a suitable solid substrate is the commercially available BIACore<sup>TM</sup> chip (Pharmacia Biosensors).

5 Preferably, the solid substrate is generally a material having a rigid or semi-rigid surface. In preferred embodiments, at least one surface of the substrate will be substantially flat, although in some embodiments it are desirable to physically separate synthesis regions for different polymers with, for example, raised regions or etched trenches. It is also preferred that the solid substrate is suitable for the high density application of DNA 10 sequences in discrete areas of typically from 50 to 100 µm, giving a density of 10000 to 40000 cm<sup>-2</sup>.

The solid substrate is conveniently divided up into sections. This are achieved by techniques such as photoetching, or by the application of hydrophobic inks, for example teflon-based inks (Cel-line, USA).

15

30

35

Discrete positions, in which each different member of the array is located may have any convenient shape, e.g., circular, rectangular, elliptical, wedge-shaped, etc.

20 Attachment of the polynucleotide sequences to the substrate are by covalent or noncovalent means. The plurality of polynucleotide sequences are attached to the substrate via a layer of molecules to which the sequences bind. For example, the sequences are labelled with biotin and the substrate coated with avidin and/or streptavidin. convenient feature of using biotinylated sequences is that the efficiency of coupling to the 25 solid substrate are determined easily. Since the library sequences may bind only poorly to some solid substrates, it is often necessary to provide a chemical interface between the solid substrate (such as in the case of glass) and the sequences. Examples of suitable chemical interfaces include hexaethylene glycol. Another example is the use of polylysine coated glass, the polylysine then being chemically modified using standard procedures to introduce an affinity ligand. Other methods for attaching molecules to the surfaces of solid substrate by the use of coupling agents are known in the art, see for example WO98/49557.

The complete DNA array is typically read at the same time by charged coupled device (CCD) camera or confocal imaging system. Alternatively, the DNA array are placed for detection in a suitable apparatus that can move in an x-y direction, such as a plate

58

reader. In this way, the change in characteristics for each discrete position are measured automatically by computer controlled movement of the array to place each discrete element in turn in line with the detection means.

The detection means are capable of interrogating each position in the library array optically or electrically. Examples of suitable detection means include CCD cameras or confocal imaging systems.

In a preferred embodiment, the level of expression of the cancer-associated gene in the test sample is determined by hybridizing a probe/primer to RNA in the test sample under at least low stringency hybridization conditions and detecting the hybridization using a detection means.

Similarly, the level of mRNA in the comparable sample from the healthy or normal individual is preferably determined by hybridizing a probe/primer to RNA in said comparable sample under at least low stringency hybridization conditions and detecting the hybridization using a detection means.

For the purposes of defining the level of stringency to be used in these diagnostic assays, a low stringency is defined herein as being a hybridization and/or a wash carried out in 6xSSC buffer, 0.1% (w/v) SDS at 28°C, or equivalent conditions. A moderate stringency is defined herein as being a hybridization and/or washing carried out in 2xSSC buffer, 0.1% (w/v) SDS at a temperature in the range 45°C to 65°C, or equivalent conditions. A high stringency is defined herein as being a hybridization and/or wash carried out in 0.1xSSC buffer, 0.1% (w/v) SDS, or lower salt concentration, and at a temperature of at least 65°C, or equivalent conditions. Reference herein to a particular level of stringency encompasses equivalent conditions using wash/hybridization solutions other than SSC known to those skilled in the art.

Generally, the stringency is increased by reducing the concentration of SSC buffer, and/or increasing the concentration of SDS and/or increasing the temperature of the hybridization and/or wash. Those skilled in the art will be aware that the conditions for hybridization and/or wash may vary depending upon the nature of the hybridization matrix used to support the sample RNA, or the type of hybridization probe used.

15

20

25

59

In general, the sample or the probe is immobilized on a solid matrix or surface (e.g., nitrocellulose). For high throughput screening, the sample or probe will generally comprise an array of nucleic acids on glass or other solid matrix, such as, for example, as described in WO 96/17958. Techniques for producing high density arrays are described, for example, by Fodor *et al.*, Science 767-773, 1991, and in U.S. Pat. No. 5,143,854. Typical protocols for other assay formats can be found, for example in Current Protocols In Molecular Biology, Unit 2 (Northern Blotting), Unit 4 (Southern Blotting), and Unit 18 (PCR Analysis), Frederick M. Ausubul *et al.* (ed)., 1995.

The detection means according to this aspect of the invention may be any nucleic acidbased detection means such as, for example, nucleic acid hybridization or amplification reaction (eg. PCR), a nucleic acid sequence-based amplification (NASBA) system, inverse polymerase chain reaction (iPCR), *in situ* polymerase chain reaction, or reverse transcription polymerase chain reaction (RT-PCR), amongst others.

15

20

The probe can be labelled with a reporter molecule capable of producing an identifiable signal (e.g., a radioisotope such as <sup>32</sup>P or <sup>35</sup>S, or a fluorescent or biotinylated molecule). According to this embodiment, those skilled in the art will be aware that the detection of sald reporter molecule provides for identification of the probe and that, following the hybridization reaction, the detection of the corresponding nucleotide sequences in the sample is facilitated. Additional probes can be used to confirm the assay results obtained using a single probe.

Wherein the detection means is an amplification reaction such as, for example, a polymerase chain reaction or a nucleic acid sequence-based amplification (NASBA) system or a variant thereof, one or more nucleic acid probes molecules of at least about 20 contiguous nucleotides in length is hybridized to mRNA encoding a cancer-associated protein, or alternatively, hybridized to cDNA or cRNA produced from said mRNA, and nucleic acid copies of the template are enzymically-amplified.

30

25

Those skilled in the art will be aware that there must be a sufficiently high percentage of nucleotide sequence identity between the probes and the RNA sequences in the sample template molecule for hybridization to occur. As stated previously, the stringency conditions can be selected to promote hybridization.

In one format, PCR provides for the hybridization of non-complementary probes to different strands of a double-stranded nucleic acid template molecule (ie. a DNA/RNA, RNA/RNA or DNA/DNA template), such that the hybridized probes are positioned to facilitate the 5'-to 3' synthesis of nucleic acid in the intervening region, under the control of a thermostable DNA polymerase enzyme. In accordance with this embodiment, one sense probe and one antisense probe as described herein would be used to amplify DNA from the hybrid RNA/DNA template or cDNA.

In the present context, the cDNA would generally be produced by reverse transcription of mRNA present in the sample being tested (ie. RT-PCR). RT-PCR is particularly useful when it is desirable to determine expression of a cancer-associated gene. It is also known to those skilled in the art to use mRNA/DNA hybrid molecules as a template for such amplification reactions, and, as a consequence, first strand cDNA synthesis is all that is required to be performed prior to the amplification reaction.

15

30

10

5

Variations of the embodiments described herein are described in detail by McPherson *et al.*, PCR: A Practical Approach. (series eds, D. Rickwood and B.D. Hames), IRL Press Limited, Oxford. pp1-253, 1991.

The amplification reaction detection means described *supra* can be further coupled to a classical hybridization reaction detection means to further enhance sensitivity and specificity of the inventive method, such as by hybridizing the amplified DNA with a probe which is different from any of the probes used in the amplification reaction.

Similarly, the hybridization reaction detection means described *supra* can be further coupled to a second hybridization step employing a probe which is different from the probe used in the first hybridization reaction.

The comparison to be performed in accordance with the present invention may be a visual comparison of the signal generated by the probe, or alternatively, a comparison of data integrated from the signal, such as, for example, data that have been corrected or normalized to allow for variation between samples. Such comparisons can be readily performed by those skilled in the art.

# **Polypeptides**

5

25

30

35

Cancer-associated polypeptides are encoded by cancer-associated genes. It will be understood that such polypeptides include those polypeptide and fragments thereof that are homologous to the polypeptides encoded by the nucleotide sequences referred to in Tables 1-3, which are obtained from any source, for example related viral/bacterial proteins, cellular homologues and synthetic peptides, as well as variants or derivatives thereof.

Thus, the present invention encompasses the use of variants, homologues or derivatives of the cancer-associated proteins descirbed in the accompanying Tables. In one embodiment, homologues are naturally occurring sequences, such as orthologues, tissue-specific isoforms and allelic variants.

In the context of the present invention, a homologous sequence is taken to include an amino acid sequence which is at least 60, 70, 80 or 90% identical, preferably at least 95 or 98% identical at the amino acid level over at least 20, 40, 60 or 80 amino acids with a sequence encoded by a nucleotide sequence referred to in any one of Tables 1-3. In particular, homology should typically be considered with respect to those regions of the sequence known to be essential for specific biological functions rather than non-essential neighbouring sequences.

Although amino acid homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

Homology comparisons are carried out as described above for nucleotide sequences with the appropriate modifications for amino acid sequences. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

It should also be noted that where computer algorithms are used to align amino acid sequences, although the final % homology are measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example

of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). It is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

62

The terms "variant" or "derivative" in relation to the amino acid sequences of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acids from or to the sequence providing the resultant amino acid sequence preferably has biological activity, preferably having at least 25 to 50% of the activity as the polypeptides referred to in the sequence listings, more preferably at least substantially the same activity. Particular details of biological activity for each polypeptide are given in Tables 1-3.

Thus, the polypeptides referred to in Tables 1-3 and homologues thereof, are modified for use in the present invention. Typically, modifications are made that maintain the activity of the sequence. Thus, in one embodiment, amino acid substitutions are made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions provided that the modified sequence retains at least about 25 to 50% of, or substantially the same activity. However, in an alternative preferred embodiment, modifications to the amino acid sequences of a cancer-associated protein are made intentionally to reduce the biological activity of the polypeptide. For example truncated polypeptides that remain capable of binding to target molecules but lack functional effector domains are useful as inhibitors of the biological activity of the full length molecule.

25

35

5

10

In general, preferably less than 20%, 10% or 5% of the amino acid residues of a variant or derivative are altered as compared with the corresponding region of the polypeptides referred to in Tables 1-3.

Amino acid substitutions may include the use of non-naturally occurring analogues, for example to increase blood plasma half-life of a therapeutically administered polypeptide (see below for further details on the production of peptide derivatives for use in therapy).

Conservative substitutions are made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column are substituted for each other:

ALIDUATIO	I Nam malan	
ALIPHATIC	Non-polar	GAP
		ILV
	Polar - uncharged	CSTM
		NQ
	Polar - charged	DE
		KR
AROMATIC		HFWY

Cancer-associated proteins also include fragments of the above mentioned full length polypeptides and variants thereof, including fragments of the sequences referred to in Tables 1-3 and homologues thereof. Preferred fragments include those which include an epitope. Suitable fragments will be at least about 6 or 8, e.g. at least 10, 12, 15 or 20 amino acids in length. They may also be less than 200, 100 or 50 amino acids in length. Polypeptide fragments may contain one or more (e.g. 2, 3, 5, or 10) substitutions, deletions or insertions, including conserved substitutions. Where substitutions, deletion and/or insertions have been made, for example by means of recombinant technology, preferably less than 20%, 10% or 5% of the amino acid residues depicted in the sequence listings are altered.

Cancer-associated proteins are preferably in a substantially isolated form. It will be understood that the protein are mixed with carriers or diluents which will not interfere with the intended purpose of the protein and still be regarded as substantially isolated. A cancer-associated protein of the invention may also be in a substantially purified form, in which case it will generally comprise the protein in a preparation in which more than 90%, e.g. 95%, 98% or 99% pure as determined by SDS/PAGE or other art-recognized means for assessing protein purity.

### Protein Production

10

15

20

25

For producing full-length polypeptides or immunologically active derivatives thereof by recombinant means, a protein-encoding region comprising at least about 15 contiguous nucleotides of the protein-encoding region of a nucleic acid referred to in any one of Tables 1-3 is placed in operable connection with a promoter or other regulatory sequence capable of regulating expression in a cell-free system or cellular system.

WO 2004/022778

10

15

20

25

30

35

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences of a classical genomic gene, including the TATA box which is required for accurate transcription initiation, with or without a CCAAT box sequence and additional regulatory elements (i.e., upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. In the present context, the term "promoter" is also used to describe a recombinant, synthetic or fusion molecule, or derivative which confers, activates or enhances the expression of a nucleic acid molecule to which it is operably connected, and which encodes the polypeptide or peptide fragment. Preferred promoters can contain additional copies of one or more specific regulatory elements to further enhance expression and/or to alter the spatial expression and/or temporal expression of the said nucleic acid molecule.

Placing a nucleic acid molecule under the regulatory control of, i.e., "in operable connection with", a promoter sequence means positioning said molecule such that expression is controlled by the promoter sequence. Promoters are generally positioned 5' (upstream) to the coding sequence that they control. To construct heterologous promoter/structural gene combinations, it is generally preferred to position the promoter at a distance from the gene transcription start site that is approximately the same as the distance between that promoter and the gene it controls in its natural setting, i.e., the gene from which the promoter is derived. Furthermore, the regulatory elements comprising a promoter are usually positioned within 2 kb of the start site of transcription of the gene. As is known in the art, some variation in this distance can be accommodated without loss of promoter function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the positioning of the element in its natural setting, i.e., the genes from which it is derived. Again, as is known in the art, some variation in this distance can also occur.

The prerequisite for producing intact polypeptides and peptides in bacteria such as E. coli is the use of a strong promoter with an effective ribosome binding site. Typical promoters suitable for expression in bacterial cells such as E. coli include, but are not limited to, the lacz promoter, temperature-sensitive  $\lambda_L$  or  $\lambda_R$  promoters, T7 promoter or the IPTG-inducible tac promoter. A number of other vector systems for expressing the nucleic acid molecule of the invention in E. coli are well-known in the art and are described, for example, in Ausubel et al (ln: Current Protocols in Molecular Biology. Wiley Interscience, ISBN 047150338, 1987) or Sambrook et al (ln: Molecular cloning. A

WO 2004/022778 PCT/AU2003/001166

laboratory manual, second edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989). Numerous plasmids with suitable promoter sequences for expression in bacteria and efficient ribosome binding sites have been described, such as for example, pKC30 (λ<sub>L</sub>: Shimatake and Rosenberg, *Nature 292*, 128, 1981); pKK173-3 (*tac*: Amann and Brosius, *Gene 40*, 183, 1985), pET-3 (T7: Studier and Moffat, *J. Mol. Biol. 189*, 113, 1986); the pBAD/TOPO or pBAD/Thio-TOPO series of vectors containing an arabinose-inducible promoter (Invitrogen, Carlsbad, CA), the latter of which is designed to also produce fusion proteins with thioredoxin to enhance solubility of the expressed protein; the pFLEX series of expression vectors (Pfizer Inc., CT, USA); or the pQE series of expression vectors (Qiagen, CA), amongst others.

Typical promoters suitable for expression in viruses of eukaryotic cells and eukaryotic cells include the SV40 late promoter, SV40 early promoter and cytomegalovirus (CMV) promoter, CMV IE (cytomegalovirus immediate early) promoter amongst others. Preferred vectors for expression in mammalian cells (eg. 293, COS, CHO, 293T cells) include, but are not limited to, the pcDNA vector suite supplied by Invitrogen, in particular pcDNA 3.1 myc-His-tag comprising the CMV promoter and encoding a C-terminal 6xHis and MYC tag; and the retrovirus vector pSRαtkneo (Muller *et al.*, *Mol. Cell. Biol.*, *11*, 1785, 1991). The vector pcDNA 3.1 myc-His (Invitrogen) is particularly preferred for expressing a secreted form of a protein in 293T cells, wherein the expressed peptide or protein can be purified free of conspecific proteins, using standard affinity techniques that employ a Nickel column to bind the protein via the His tag.

A wide range of additional host/vector systems suitable for expressing polypeptides or immunological derivatives thereof are available publicly, and described, for example, in Sambrook *et al* (*In:* Molecular cloning. A laboratory manual, second edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989).

Means for introducing the isolated nucleic acid molecule or a gene construct comprising same into a cell for expression are well-known to those skilled in the art. The technique used for a given organism depends on the known successful techniques. Means for introducing recombinant DNA into animal cells include microinjection, transfection mediated by DEAE-dextran, transfection mediated by liposomes such as by using lipofectamine (Gibco, MD, USA) and/or cellfectin (Gibco, MD, USA), PEG-mediated DNA uptake, electroporation and microparticle bombardment such as by using DNA-coated tungsten or gold particles (Agracetus Inc., WI, USA) amongst others.

5

10

15

20

25

For producing mutants, nucleotide insertion derivatives of the protein-encoding region are produced by making 5' and 3' terminal fusions, or by making intra-sequence insertions of single or multiple nucleotides or nucleotide analogues. Insertion nucleotide sequence variants are produced by introducing one or more nucleotides or nucleotide analogues into a predetermined site in the nucleotide sequence of said sequence, although random insertion is also possible with suitable screening of the resulting product being performed. Deletion variants are produced by removing one or more nucleotides from the nucleotide sequence. Substitutional nucleotide variants are produced by substituting at least one nucleotide in the sequence with a different nucleotide or a nucleotide analogue in its place, with the immunologically active derivative encoded therefor having an identical amino acid sequence, or only a limited number of amino acid modifications that do not alter its antigenicity compared to the base peptide or its ability to bind antibodies prepared against the base peptide. Such mutant derivatives will preferably have at least 80% identity with the base amino acid sequence from which they are derived.

Preferred immunologically active derivatives of a full-length polypeptide encoded by a gene referred to in any one of Tables 1-3 will comprise at least about 5-10 contiguous amino acids of the full-length amino acid sequence, more preferably at least about 10-20 contiguous amino acids in length, and even more preferably 20-30 contiguous amino acids in length.

For the purposes of producing derivatives using standard peptide synthesis techniques, such as, for example, Fmoc chemistry, a length not exceeding about 30-50 amino acids in length is preferred, as longer peptides are difficult to produce at high efficiency. Longer peptide fragments are readily achieved using recombinant DNA techniques wherein the peptide is expressed in a cell-free or cellular expression system comprising nucleic acid encoding the desired peptide fragment.

30

It will be apparent to the skilled artisan that any sufficiently antigenic region of at least about 5-10 amino acid residues can be used to prepare antibodies that bind generally to the polypeptides listed in Tables 1-3.

An expressed protein or synthetic peptide is preferably produced as a recombinant fusion protein, such as for example, to aid in extraction and purification. To produce a fusion

polypeptide, the open reading frames are covalently linked in the same reading frame, such as, for example, using standard cloning procedures as described by Ausubel *et al.* (Current Protocols in Molecular Biology, Wiley Interscience, ISBN 047150338, 1992), and expressed under control of a promoter. Examples of fusion protein partners include glutathione-S-transferase (GST), FLAG, hexahistidine, GAL4 (DNA binding and/or transcriptional activation domains) and  $\beta$ -galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. Preferably the fusion protein will not hinder the immune function of the target protein.

10

5

In a particularly preferred embodiment, polypeptides are produced substantially free of conspecific proteins. Such purity can be assessed by standard procedures, such as, for example, SDS/polyacrylamide gel electrophoresis, 2-dimensional gene electrophoresis, chromatography, amino acid composition analysis, or amino acid sequence analysis.

15

20

25

30

35

To produce isolated polypeptides or fragments, eg., for antibody production, standard protein purification techniques may be employed. For example, gel filtration, ion exchange chromatography, reverse phase chromatography, or affinity chromatography, or a combination of any one or more said procedures, may be used. High pressure and low pressure procedures can also be employed, such as, for example, FPLC, or HPLC. To isolate the full-length proteins or peptide fragments comprising more than about 50-100 amino acids in length, it is particularly preferred to express the polypeptide in a suitable cellular expression system in combination with a suitable affinity tag, such as a 6xHis tag, and to purify the polypeptide using an affinity step that bonds it via the tag (supra). Optionally, the tag may then be cleaved from the expressed polypeptide.

Alternatively, for short immunologically active derivatives of a full-length polypeptide, preferably those peptide fragments comprising less than about 50 amino acids in length, chemical synthesis techniques are conveniently used. As will be known to those skilled in the art, such techniques may also produce contaminating peptides that are shorter than the desired peptide, in which case the desired peptide is conveniently purified using reverse phase and/or ion exchange chromatography procedures at high pressure (ie. HPLC or FPLC).

WO 2004/022778

PCT/AU2003/001166

68

The invention also provides monoclonal or polyclonal antibodies that bind specifically to polypeptides of the invention or fragments thereof. Thus, the present invention further provides a process for the production of monoclonal or polyclonal antibodies to polypeptides of the invention.

5

10

15

The phrase "binds specifically" to a polypeptide means that the binding of the antibody to the protein or peptide is determinative of the presence of the protein, in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and more typically more than 10 to 100 times background. Typically, antibodies of the invention bind to a protein of interest with a Kd of at least about 0.1 mM, more usually at least about 1  $\mu$ M, preferably at least about 0.1  $\mu$ M, and most preferably at least, 0.01  $\mu$ M.

Reference herein to antibody or antibodies includes whole polyclonal and monoclonal antibodies, and parts thereof, either alone or conjugated with other moieties. Antibody parts include Fab and F(ab)<sub>2</sub> fragments and single chain antibodies. The antibodies may be made *in vivo* in suitable laboratory animals, or, in the case of engineered antibodies (Single Chain Antibodies or SCABS, etc) using recombinant DNA techniques *in vitro*.

20

25

30

35

In accordance with this aspect of the invention, the antibodies may be produced for the purposes of immunizing the subject, in which case high titer or neutralizing antibodies that bind to a B cell epitope will be especially preferred. Suitable subjects for immunization will, of course, depend upon the immunizing antigen or antigenic B cell epitope. It is contemplated that the present invention will be broadly applicable to the immunization of a wide range of animals, such as, for example, farm animals (e.g. horses, cattle, sheep, pigs, goats, chickens, ducks, turkeys, and the like), laboratory animals (e.g. rats, mice, guinea pigs, rabbits), domestic animals (cats, dogs, birds and the like), feral or wild exotic animals (e.g. possums, cats, pigs, buffalo, wild dogs and the like) and humans.

Alternatively, the antibodies may be for commercial or diagnostic purposes, in which case the subject to whom the diagnostic/prognostic protein or immunogenic fragment or epitope thereof is administered will most likely be a laboratory or farm animal. A wide range of animal species are used for the production of antisera. Typically the animal used for production of antisera is a rabbit, a mouse, rat, hamster, guinea pig, goat,

5

10

15

20

25

30

35

sheep, pig, dog, horse, or chicken. Because of the relatively large blood volume of rabbits, a rabbit is a preferred choice for production of polyclonal antibodies. However, as will be known to those skilled in the art, larger amounts of immunogen are required to obtain high antibodies from large animals as opposed to smaller animals such as mice. In such cases, it will be desirable to isolate the antibody from the immunized animal.

Preferably, the antibody is a high titer antibody. By "high titer" means a sufficiently high titer to be suitable for use in diagnostic or therapeutic applications. As will be known in the art, there is some variation in what might be considered "high titer". For most applications a titer of at least about 10<sup>3</sup>-10<sup>4</sup> is preferred. More preferably, the antibody titer will be in the range from about 10<sup>4</sup> to about 10<sup>5</sup>, even more preferably in the range from about 10<sup>6</sup>.

More preferably, in the case of B cell epitopes from pathogens, viruses or bacteria, the antibody is a neutralizing antibody (i.e. it is capable of neutralizing the infectivity of the organism fro which the B cell epitope is derived).

To generate antibodies, the diagnostic/prognostic protein or immunogenic fragment or epitope thereof, optionally formulated with any suitable or desired carrier, adjuvant, BRM, or pharmaceutically acceptable excipient, is conveniently administered in the form of an injectable composition. Injection may be intranasal, intramuscular, sub-cutaneous, intravenous, intradermal, intraperitoneal, or by other known route. For intravenous injection, it is desirable to include one or more fluid and nutrient replenishers. Means for preparing and characterizing antibodies are well known in the art. (See, e.g., ANTIBODIES: A LABORATORY MANUAL, Cold Spring Harbor Laboratory, 1988, incorporated herein by reference).

The efficacy of the diagnostic/prognostic protein or immunogenic fragment or epitope thereof in producing an antibody is established by injecting an animal, for example, a mouse, rat, rabbit, guinea pig, dog, horse, cow, goat or pig, with a formulation comprising the diagnostic/prognostic protein or immunogenic fragment or epitope thereof, and then monitoring the immune response to the B cell epitope, as described in the Examples. Both primary and secondary immune responses are monitored. The antibody titer is determined using any conventional immunoassay, such as, for example, ELISA, or radio immunoassay.

. .70

The production of polyclonal antibodies may be monitored by sampling blood of the immunized animal at various points following immunization. A second, booster injection, may be given, if required to achieve a desired antibody titer. The process of boosting and titering is repeated until a suitable titer is achieved. When a desired level of immunogenicity is obtained, the immunized animal is bled and the serum isolated and stored, and/or the animal is used to generate monoclonal antibodies (Mabs).

For the production of monoclonal antibodies (Mabs) any one of a number of well-known techniques may be used, such as, for example, the procedure exemplified in US Patent No. 4,196,265, incorporated herein by reference.

For example, a suitable animal will be immunized with an effective amount of the diagnostic/prognostic protein or immunogenic fragment or epitope thereof under conditions sufficient to stimulate antibody producing cells. Rodents such as mice and rats are preferred animals, however, the use of rabbit, sheep, or frog cells is also possible. The use of rats may provide certain advantages, but mice are preferred, with the BALB/c mouse being most preferred as the most routinely used animal and one that generally gives a higher percentage of stable fusions.

Following immunization, somatic cells with the potential for producing antibodies, specifically B lymphocytes (B cells), are selected for use in the MAb generating protocol. These cells may be obtained from biopsied spleens, tonsils or lymph nodes, or from a peripheral blood sample. Spleen cells and peripheral blood cells are preferred, the former because they are a rich source of antibody-producing cells that are in the dividing plasmablast stage, and the latter because peripheral blood is easily accessible. Often, a panel of animals will have been immunized and the spleen of animal with the highest antibody titer removed. Spleen lymphocytes are obtained by homogenizing the spleen with a syringe. Typically, a spleen from an immunized mouse contains approximately  $5 \times 10^7$  to  $2 \times 10^8$  lymphocytes.

30

35

5

10

15

20

25

The B cells from the immunized animal are then fused with cells of an immortal myeloma cell, generally derived from the same species as the animal that was immunized with the diagnostic/prognostic protein or immunogenic fragment or epitope thereof. Myeloma cell lines suited for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency and enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the

desired fused cells, or hybridomas. Any one of a number of myeloma cells may be used and these are known to those of skill in the art (e.g. murine P3-X63/Ag8, X63-Ag8.653, NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XX0; or rat R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210; and U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6). A preferred murine myeloma cell is the NS-1 myeloma cell line (also termed P3-NS-1-Ag4-1), which is readily available from the NIGMS Human Genetic Mutant Cell Repository under Accession No. GM3573. Alternatively, a murine myeloma SP2/0 non-producer cell line that is 8-azaguanine-resistant is used.

To generate hybrids of antibody-producing spleen or lymph node cells and myeloma cells, somatic cells are mixed with myeloma cells in a proportion between about 20:1 to about 1:1, respectively, in the presence of an agent or agents (chemical or electrical) that promote the fusion of cell membranes. Fusion methods using Sendai virus have been described by Kohler and Milstein, *Nature* 256, 495-497, 1975; and Kohler and Milstein, *Eur. J. Immunol.* 6, 511-519, 1976. Methods using polyethylene glycol (PEG), such as 37% (v/v) PEG, are described in detail by Gefter *et al., Somatic Cell Genet.* 3, 231-236, 1977. The use of electrically induced fusion methods is also appropriate.

Hybrids are amplified by culture in a selective medium comprising an agent that blocks the *de novo* synthesis of nucleotides in the tissue culture media. Exemplary and preferred agents are aminopterin, methotrexate and azaserine. Aminopterin and methotrexate block de novo synthesis of both purines and pyrimidines, whereas azaserine blocks only purine synthesis. Where aminopterin or methotrexate is used, the media is supplemented with hypoxanthine and thymidine as a source of nucleotides (HAT medium). Where azaserine is used, the media is supplemented with hypoxanthine.

The preferred selection medium is HAT, because only those hybridomas capable of operating nucleotide salvage pathways are able to survive in HAT medium, whereas myeloma cells are defective in key enzymes of the salvage pathway, (e.g., hypoxanthine phosphoribosyl transferase or HPRT), and they cannot survive. B cells can operate this salvage pathway, but they have a limited life span in culture and generally die within about two weeks. Accordingly, the only cells that can survive in the selective media are those hybrids formed from myeloma and B cells.

72

The amplified hybridomas are subjected to a functional selection for antibody specificity and/or titer, such as, for example, by immunoassay (e.g. radioimmunoassay, enzyme immunoassay, cytotoxicity assay, plaque assay, dot immunobinding assay, and the like).

5 The selected hybridomas are serially diluted and cloned into individual antibodyproducing cell lines, which clones can then be propagated indefinitely to provide MAbs. The cell lines may be exploited for MAb production in two basic ways. A sample of the hybridoma is injected, usually in the peritoneal cavity, into a histocompatible animal of the type that was used to provide the somatic and myeloma cells for the original fusion. 10 The injected animal develops tumors secreting the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can then be tapped to provide MAbs in high concentration. The individual cell lines could also be cultured in vitro, where the MAbs are naturally secreted into the culture medium from which they are readily obtained in high concentrations. MAbs produced by either means may be further purified, if desired, using filtration, 15 centrifugation and various chromatographic methods such as HPLC or affinity chromatography.

Monoclonal antibodies of the present invention also include anti-idiotypic antibodies produced by methods well-known in the art. Monoclonal antibodies according to the present invention also may be monoclonal heteroconjugates, (i.e., hybrids of two or more antibody molecules). In another embodiment, monoclonal antibodies according to the invention are chimeric monoclonal antibodies. In one approach, the chimeric monoclonal antibody is engineered by cloning recombinant DNA containing the promoter, leader, and variable-region sequences from a mouse anti-PSA producing cell and the constant-region exons from a human antibody gene. The antibody encoded by such a recombinant gene is a mouse-human chimera. Its antibody specificity is determined by the variable region derived from mouse sequences. Its isotype, which is determined by the constant region, is derived from human DNA.

30

35

20

25

In another embodiment, the monoclonal antibody according to the present invention is a "humanized" monoclonal antibody, produced by any one of a number of techniques well-known in the art. That is, mouse complementary determining regions ("CDRs") are transferred from heavy and light V-chains of the mouse Ig into a human V-domain, followed by the replacement of some human residues in the framework regions of their

murine counterparts. "Humanized" monoclonal antibodies in accordance with this invention are especially suitable for use *in vivo* in diagnostic and therapeutic methods.

As stated above, the monoclonal antibodies and fragments thereof according to this invention are multiplied according to *in vitro* and *in vivo* methods well-known in the art. Multiplication *in vitro* is carried out in suitable culture media such as Dulbecco's modified Eagle medium or RPMI 1640 medium, optionally replenished by a mammalian serum such as fetal calf serum or trace elements and growth-sustaining supplements, e.g., feeder cells, such as normal mouse peritoneal exudate cells, spleen cells, bone marrow macrophages or the like. In vitro production provides relatively pure antibody preparations and allows scale-up to give large amounts of the desired antibodies. Techniques for large scale hybridoma cultivation under tissue culture conditions are known in the art and include homogenous suspension culture, (e.g., in an airlift reactor or in a continuous stirrer reactor or immobilized or entrapped cell culture).

15

20

25

10

5

Large amounts of the monoclonal antibody of the present invention also may be obtained by multiplying hybridoma cells *in vivo*. Cell clones are injected into mammals which are histocompatible with the parent cells, (e.g., syngeneic mice, to cause growth of antibody-producing tumors. Optionally, the animals are primed with a hydrocarbon, especially oils such as Pristane (tetramethylpentadecane) prior to injection.

In accordance with the present invention, fragments of the monoclonal antibody of the invention are obtained from monoclonal antibodies produced as described above, by methods which include digestion with enzymes such as pepsin or papain and/or cleavage of disulfide bonds by chemical reduction. Alternatively, monoclonal antibody fragments encompassed by the present invention are synthesized using an automated peptide synthesizer, or they may be produced manually using techniques well known in the art.

The monoclonal conjugates of the present invention are prepared by methods known in the art, e.g., by reacting a monoclonal antibody prepared as described above with, for instance, an enzyme in the presence of a coupling agent such as glutaraldehyde or periodate. Conjugates with fluorescein markers are prepared in the presence of these coupling agents, or by reaction with an isothiocyanate. Conjugates with metal chelates are similarly produced. Other moieties to which antibodies may be conjugated include

radionuclides such as, for example, <sup>3</sup>H, <sup>125</sup>I, .<sup>32</sup>P, .<sup>35</sup>S, <sup>14</sup>C, <sup>51</sup>Cr, <sup>35</sup>Cl, <sup>57</sup>Co, <sup>58</sup>Co, <sup>59</sup>Fe, <sup>75</sup>Se, and <sup>152</sup>Eu.

Radioactively labeled monoclonal antibodies of the present invention are produced according to well-known methods in the art. For instance, monoclonal antibodies are iodinated by contact with sodium or potassium iodide and a chemical oxidizing agent such as sodium hypochlorite, or an enzymatic oxidizing agent, such as lactoperoxidase. Monoclonal antibodies according to the invention may be labeled with technetium<sup>99</sup> by ligand exchange process, for example, by reducing pertechnetate with stannous solution, chelating the reduced technetium onto a Sephadex column and applying the antibody to this column or by direct labeling techniques, (e.g., by incubating pertechnate, a reducing agent such as SNCl<sub>2</sub>, a buffer solution such as sodium-potassium phthalate solution, and the antibody).

Any immunoassay may be used to monitor antibody production by the diagnostic/prognostic protein or immunogenic fragment or epitope thereof. Immunoassays, in their most simple and direct sense, are binding assays. Certain preferred immunoassays are the various types of enzyme linked immunosorbent assays (ELISAs) and radioimmunoassays (RIA) known in the art. Immunohistochemical detection using tissue sections is also particularly useful. However, it will be readily appreciated that detection is not limited to such techniques, and Western blotting, dot blotting, FACS analyses, and the like may also be used.

Most preferably, the assay will be capable of generating quantitative results.

25

30

35

5

10

For example, antibodies are tested in simple competition assays. A known antibody preparation that binds to the B cell epitope and the test antibody are incubated with an antigen composition comprising the B cell epitope, preferably in the context of the native antigen. "Antigen composition" as used herein means any composition that contains some version of the B cell epitope in an accessible form. Antigen-coated wells of an ELISA plate are particularly preferred. In one embodiment, one would pre-mix the known antibodies with varying amounts of the test antibodies (e.g., 1:1, 1:10 and 1:100) for a period of time prior to applying to the antigen composition. If one of the known antibodies is labeled, direct detection of the label bound to the antigen is possible; comparison to an unmixed sample assay will determine competition by the test antibody and, hence, cross-

5

10

15

20

30

35

reactivity. Alternatively, using secondary antibodies specific for either the known or test antibody, one will be able to determine competition.

An antibody that binds to the antigen composition will be able to effectively compete for binding of the known antibody and thus will significantly reduce binding of the latter. The reactivity of the known antibodies in the absence of any test antibody is the control. A significant reduction in reactivity in the presence of a test antibody is indicative of a test antibody that binds to the B cell epitope (i.e., it cross-reacts with the known antibody).

In one exemplary ELISA, the antibodies against the diagnostic/prognostic protein or immunogenic fragment or B cell epitope are immobilized onto a selected surface exhibiting protein affinity, such as a well in a polystyrene microtiter plate. Then, a composition containing a peptide comprising the B cell epitope is added to the wells. After binding and washing to remove non-specifically bound immune complexes, the bound epitope may be detected. Detection is generally achieved by the addition of a second antibody that is known to bind to the B cell epitope and is linked to a detectable label. This type of ELISA is a simple "sandwich ELISA". Detection may also be achieved by the addition of said second antibody, followed by the addition of a third antibody that has binding affinity for the second antibody, with the third antibody being linked to a detectable label.

Antibodies of the invention may be bound to a solid support and/or packaged into kits in a suitable container along with suitable reagents, controls, instructions and the like.

## 25 Immunoassay formats

In one embodiment, a cancer-associated protein or an immunogenic fragment or epitope thereof is detected in a patient sample, wherein the level of the protein or immunogenic fragment or epitope in the sample is indicative of ovarian cancer or disease recurrence or an indicator of poor survival. Preferably, the method comprises contacting a biological sample derived from the subject with an antibody capable of binding to a cancer-associated protein or an immunogenic fragment or epitope thereof, and detecting the formation of an antigen-antibody complex.

In another embodiment, an antibody against a cancer-associated protein or epitope thereof is detected in a patient sample, wherein the level of the antibody in the sample is indicative of ovarian cancer or disease recurrence or an indicator of poor survival.

Preferably, the method comprises contacting a biological sample derived from the subject with a cancer-associated protein or an antigenic fragment eg., a B cell epitope or other immunogenic fragment thereof, and detecting the formation of an antigen-antibody complex.

5

10

15

20

The diagnostic assays of the invention are useful for determining the progression of ovarian cancer or a metastasis thereof in a subject. In accordance with these prognostic applications of the invention, the level of a cancer-associated protein or an immunogenic fragment or epitope thereof in a biological sample is correlated with the disease state eg., as determined by clinical symptoms or biochemical tests (eg., CA125 levels).

Accordingly, a further embodiment of the invention provides a method for detecting a cancer cell in a subject, said method comprising:

- determining the level of a cancer-associate protein in a test sample from said subject; and
- (ii) comparing the level determined at (i) to the level of said cancer-associated protein in a comparable sample from a healthy or normal individual,

wherein a level of said cancer-associate protein at (i) that is modified in the test sample relative to the comparable sample from the normal or healthy individual is indicative of the presence of a cancer cell in said subject.

In one embodiment of the diagnostic/prognostic methods described herein, the biological sample is obtained previously from the subject. In accordance with such an embodiment, the prognostic or diagnostic method is performed ex vivo.

25

In yet another embodiment, the subject diagnostic/prognostic methods further comprise processing the sample from the subject to produce a derivative or extract that comprises the analyte.

Preferred detection systems contemplated herein include any known assay for detecting proteins or antibodies in a biological sample isolated from a human subject, such as, for example, SDS/PAGE, isoelectric focussing, 2-dimensional gel electrophoresis comprising SDS/PAGE and isoelectric focussing, an immunoassay, a detection based system using an antibody or non-antibody ligand of the protein, such as, for example, a small molecule (e.g. a chemical compound, agonist, antagonist, allosteric modulator, competitive inhibitor, or non-competitive inhibitor, of the protein). In accordance with

5

25

30

35

these embodiments, the antibody or small molecule may be used in any standard solid phase or solution phase assay format amenable to the detection of proteins. Optical or fluorescent detection, such as, for example, using mass spectrometry, MALDI-TOF, biosensor technology, evanescent fiber optics, or fluorescence resonance energy transfer, is clearly encompassed by the present invention. Assay systems suitable for use in high throughput screening of mass samples, particularly a high throughput spectroscopy resonance method (e.g. MALDI-TOF, electrospray MS or nano-electrospray MS), are particularly contemplated.

Immunoassay formats are particularly preferred, eg., selected from the group consisting of, an immunoblot, a Western blot, a dot blot, an enzyme linked immunosorbent assay (ELISA), radioimmunoassay (RIA), enzyme immunoassay. Modified immunoassays utilizing fluorescence resonance energy transfer (FRET), isotope-coded affinity tags (ICAT), matrix-assisted laser desorption/ionization time of flight (MALDI-TOF), electrospray ionization (ESI), biosensor technology, evanescent fiber-optics technology or protein chip technology are also useful.

Preferably, the assay is a semi-quantitative assay or quantitative assay.

Standard solid phase ELISA formats are particularly useful in determining the concentration of a protein or antibody from a variety of patient samples.

In one form such as an assay involves immobilising a biological sample comprising antibodies against the cancer-associated protein or epitope, or alternatively an ovarian cancer-associated protein or an immunogenic fragment thereof, onto a solid matrix, such as, for example a polystyrene or polycarbonate microwell or dipstick, a membrane, or a glass support (e.g. a glass slide).

In the case of an antigen-based assay, an antibody that specifically binds an ovarian cancer-associated protein is brought into direct contact with the immobilised biological sample, and forms a direct bond with any of its target protein present in said sample. For an antibody-based assay, an immobilized ovarian cancer-associated protein or an immunogenic fragment or epitope thereof is contacted with the sample. The added antibody or protein in solution is generally labelled with a detectable reporter molecule, such as for example, a fluorescent label (e.g. FITC or Texas Red) or an enzyme (e.g. horseradish peroxidase (HRP)), alkaline phosphatase (AP) or β-galactosidase.

Alternatively, or in addition, a second labelled antibody can be used that binds to the first antibody or to the isolated/recombinant antigen. Following washing to remove any unbound antibody or antigen, as appropriate, the label is detected either directly, in the case of a fluorescent label, or through the addition of a substrate, such as for example hydrogen peroxide, TMB, or toluidine, or 5-bromo-4-chloro-3-indol-beta-D-

78

PCT/AU2003/001166

Such ELISA based systems are particularly suitable for quantification of the amount of a protein or antibody in a sample, such as, for example, by calibrating the detection system against known amounts of a standard.

In another form, an ELISA consists of immobilizing an antibody that specifically binds an ovarian cancer-associated protein on a solid matrix, such as, for example, a membrane, a polystyrene or polycarbonate microwell, a polystyrene or polycarbonate dipstick or a glass support. A patient sample is then brought into physical relation with said antibody, and the antigen in the sample is bound or 'captured'. The bound protein can then be detected using a labelled antibody. For example if the protein is captured from a human sample, an anti-human antibody is used to detect the captured protein. Alternatively, a third labelled antibody can be used that binds the second (detecting) antibody.

20

25

30

35

5

10

15

galaotopyranoside (x-gal).

It will be apparent to the skilled person that the assay formats described herein are amenable to high throughput formats, such as, for example automation of screening processes, or a microarray format as described in Mendoza *et al*, <u>Biotechniques</u> 27(4): 778-788, 1999. Furthermore, variations of the above described assay will be apparent to those skilled in the art, such as, for example, a competitive ELISA.

Alternatively, the presence of antibodies against the cancer-associate protein, or alternatively an oarian cancer-associated protein or an immunogenic fragment thereof, is detected using a radioimmunoassay (RIA). The basic principle of the assay is the use of a radiolabelled antibody or antigen to detect antibody antigen interactions. For example, an antibody that specifically binds to an ovarian cancer-associated protein can be bound to a solid support and a biological sample brought into direct contact with said antibody. To detect the bound antigen, an isolated and/or recombinant form of the antigen is radiolabelled is brought into contact with the same antibody. Following washing the amount of bound radioactivity is detected. As any antigen in the biological sample inhibits binding of the radiolabelled antigen the amount of radioactivity detected is inversely

79

proportional to the amount of antigen in the sample. Such an assay may be quantitated by using a standard curve using increasing known concentrations of the isolated antigen.

As will be apparent to the skilled artisan, such an assay may be modified to use any reporter molecule, such as, for example, an enzyme or a fluorescent molecule, in place of a radioactive label.

5

10

15

25

30

35

Western blotting is also useful for detecting an ovarian cancer-associated protein or an immunogenic fragment thereof. In such an assay protein from a biological sample is separated using sodium dodecyl sulphate (SDS) polyacrylamide gel electrophoresis (SDS-PAGE) using techniques well known in the art and described in, for example, Scopes (*In:* Protein Purification: Principles and Practice, Third Edition, Springer Verlag, 1994). Separated proteins are then transferred to a solid support, such as, for example, a membrane or more specifically PVDF membrane, using methods well known in the art, for example, electrotransfer. This membrane may then be blocked and probed with a labelled antibody or ligand that specifically binds an ovarian cancer-associated protein. Alternatively, a labelled secondary, or even tertiary, antibody or ligand can be used to detect the binding of a specific primary antibody.

High-throughput methods for detecting the presence or absence of antibodies, or alternatively ovarian cancer-associated protein or an immunogenic fragment thereof are particularly preferred.

In one embodiment, MALDI-TOF is used for the rapid identification of a protein. Accordingly, there is no need to detect the proteins of interest using an antibody or ligand that specifically binds to the protein of interest. Rather, proteins from a biological sample are separated using gel electrophoresis using methods well known in the art and those proteins at approximately the correct molecular weight and/or isoelectric point are analysed using MALDI-TOF to determine the presence or absence of a protein of interest.

Alternatively, MALDI or ESI or a combination of approaches is used to determine the concentration of a particular protein in a biological sample, such as, for example sputum. Such proteins are preferably well characterised previously with regard to parameters such as molecular weight and isoelectric point.

5

10

15

30

35

Biosensor devices generally employ an electrode surface in combination with current or impedance measuring elements to be integrated into a device in combination with the assay substrate (such as that described in U.S. Patent No. 5,567,301). An antibody or ligand that specifically binds to a protein of interest is preferably incorporated onto the surface of a biosensor device and a biological sample isolated from a patient (for example sputum that has been solubilised using the methods described herein) contacted to said device. A change in the detected current or impedance by the biosensor device indicates protein binding to said antibody or ligand. Some forms of biosensors known in the art also rely on surface plasmon resonance to detect protein interactions, whereby a change in the surface plasmon resonance surface of reflection is indicative of a protein binding to a ligand or antibody (U.S. Patent No. 5,485,277 and 5,492,840).

Biosensors are of particular use in high throughput analysis due to the ease of adapting such systems to micro- or nano-scales. Furthermore, such systems are conveniently adapted to incorporate several detection reagents, allowing for multiplexing of diagnostic reagents in a single biosensor unit. This permits the simultaneous detection of several epitopes in a small amount of body fluids.

Evanescent biosensors are also preferred as they do not require the pretreatment of a biological sample prior to detection of a protein of interest. An evanescent biosensor generally relies upon light of a predetermined wavelength interacting with a fluorescent molecule, such as for example, a fluorescent antibody attached near the probe's surface, to emit fluorescence at a different wavelength upon binding of the diagnostic protein to the antibody or ligand.

To produce protein chips, the proteins, peptides, polypeptides, antibodies or ligands that are able to bind specific antibodies or proteins of interest are bound to a solid support such as for example glass, polycarbonate, polytetrafluoroethylene, polystyrene, silicon oxide, metal or silicon nitride. This immobilization is either direct (e.g. by covalent linkage, such as, for example, Schiff's base formation, disulfide linkage, or amide or urea bond formation) or indirect. Methods of generating a protein chip are known in the art and are described in for example U.S. Patent Application No. 20020136821, 20020192654, 20020102617 and U.S. Patent No. 6,391,625. In order to bind a protein to a solid support it is often necessary to treat the solid support so as to create chemically reactive groups on the surface, such as, for example, with an aldehyde-containing silane reagent.

Alternatively, an antibody or ligand may be captured on a microfabricated polyacrylamide gel pad and accelerated into the gel using microelectrophoresis as described in, Arenkov et al. Anal. Biochem. 278:123-131, 2000.

A protein chip is preferably generated such that several proteins, ligands or antibodies are arrayed on said chip. This format permits the simultaneous screening for the presence of several proteins in a sample.

Alternatively, a protein chip may comprise only one protein, ligand or antibody, and be used to screen one or more patient samples for the presence of one polypeptide of interest. Such a chip may also be used to simultaneously screen an array of patient samples for a polypeptide of interest.

Preferably, a sample to be analysed using a protein chip is attached to a reporter molecule, such as, for example, a fluorescent molecule, a radioactive molecule, an enzyme, or an antibody that is detectable using methods well known in the art. Accordingly, by contacting a protein chip with a labelled sample and subsequent washing to remove any unbound proteins the presence of a bound protein is detected using methods well known in the art, such as, for example using a DNA microarray reader.

20

25

30

35

10

15

Alternatively, biomolecular interaction analysis-mass spectrometry (BIA-MS) is used to rapidly detect and characterise a protein present in complex biological samples at the low- to sub-fmole level (Nelson et al. Electrophoresis 21: 1155-1163, 2000). One technique useful in the analysis of a protein chip is surface enhanced laser desorption/ionization-time of flight-mass spectrometry (SELDI-TOF-MS) technology to characterise a protein bound to the protein chip. Alternatively, the protein chip is analysed using ESI as described in U.S. Patent Application 20020139751.

As will be apparent to the skilled artisan, protein chips are particularly amenable to multiplexing of detection reagents. Accordingly, several antibodies or ligands each able to specifically bind a different peptide or protein may be bound to different regions of said protein chip. Analysis of a biological sample using said chip then permits the detecting of multiple proteins of interest, or multiple B cell epitopes of the ovarian cancer-associated protein. Multiplexing of diagnostic and prognostic markers is particularly contemplated in the present invention.

In a further embodiment, the samples are analysed using ICAT, essentially as described in US Patent Application No. 20020076739. This system relies upon the labelling of a protein sample from one source (i.e. a healthy individual) with a reagent and the labelling of a protein sample from another source (i.e. a tuberculosis patient) with a second reagent that is chemically identical to the first reagent, but differs in mass due to isotope composition. It is preferable that the first and second reagents also comprise a biotin molecule. Equal concentrations of the two samples are then mixed, and peptides recovered by avidin affinity chromatography. Samples are then analysed using mass spectrometry. Any difference in peak heights between the heavy and light peptide ions directly correlates with a difference in protein abundance in a biological sample. The identity of such proteins may then be determined using a method well known in the art, such as, for example MALDI-TOF, or ESI.

As will be apparent to those skilled in the art a diagnostic or prognostic assay described herein may be a multiplexed assay. As used herein the term "multiplex", shall be understood not only to mean the detection of two or more diagnostic or prognostic markers in a single sample simultaneously, but also to encompass consecutive detection of two or more diagnostic or prognostic markers in a single sample, simultaneous detection of two or more diagnostic or prognostic markers in distinct but matched samples, and consecutive detection of two or more diagnostic or prognostic markers in distinct but matched samples. As used herein the term "matched samples" shall be understood to mean two or more samples derived from the same initial biological sample, or two or more biological samples isolated at the same point in time.

Accordingly, a multiplexed assay may comprise an assay that detects several antibodies and/or epitopes in the same reaction and simultaneously, or alternatively, it may detect other one or more antigens/antibodies in addition to one or more antibodies and/or epitopes. As will be apparent to the skilled artisan, if such an assay is antibody or ligand based, both of these antibodies must function under the same conditions.

30

35

5

10

15

20

25

## Diagnostic assay kits

A further aspect of the present invention provides a kit for detecting *M. tuberculosis* infection in a biological sample. In one embodiment, the kit comprises:

- (i) one or more isolated antibodies that bind to an ovarian cancer-associated protein or an immunogenic fragment or epitope thereof; and
- (ii) means for detecting the formation of an antigen-antibody complex.

In an alternative embodiment, the kit comprises:

- (i) an isolated or recombinant ovarian cancer-associated protein or an immunogenic fragment or epitope thereof; and
- (ii) means for detecting the formation of an antigen-antibody complex.

Optionally, the kit further comprises means for the detection of the binding of an antibody, fragment thereof or a ligand to an ovarian cancer-associated protein. Such means include a reporter molecule such as, for example, an enzyme (such as horseradish peroxidase or alkaline phosphatase), a substrate, a cofactor, an inhibitor, a dye, a radionucleotide, a luminescent group, a fluorescent group, biotin or a colloidal particle, such as colloidal gold or selenium. Preferably such a reporter molecule is directly linked to the antibody or ligand.

In yet another embodiment, a kit may additionally comprise a reference sample. Such a reference sample.

In another embodiment, a reference sample comprises a peptide that is detected by an antibody or a ligand. Preferably, the peptide is of known concentration. Such a peptide is of particular use as a standard. Accordingly various known concentrations of such a peptide may be detected using a prognostic or diagnostic assay described herein.

In yet another embodiment, a kit comprises means for protein isolation (Scopes (In: Protein Purification: Principles and Practice, Third Edition, Springer Verlag, 1994).

25

30

35

20

5

10

## **Bioinformatics**

The ability to identify genes that are over or under expressed in ovarian cancer can additionally provide high-resolution, high-sensitivity datasets which are used in the areas of diagnostics, therapeutics, drug development, pharmacogenetics, protein structure, biosensor development, and other related areas. For example, the expression profiles are used in diagnostic or prognostic evaluation of patients with ovarian cancer. Or as another example, subcellular toxicological information are generated to better direct drug structure and activity correlation (see Anderson, *Pharmaceutical Proteomics: Targets, Mechanism, and Function,* paper presented at the IBC Proteomics conference, Coronado, CA (June 11-12, 1998)). Subcellular toxicological information can also be utilized in a biological sensor device to predict the likely toxicological effect of chemical

5

10

15

20

25

30

35

exposures and likely tolerable exposure thresholds (see U.S. Patent No. 5,811,231). Similar advantages accrue from datasets relevant to other biomolecules and bioactive agents (e.g., nucleic acids, saccharides, lipids, drugs, and the like).

Thus, in another embodiment, the present invention provides a database that includes at least one set of assay data. The data contained in the database is acquired, e.g., using array analysis either singly or in a library format. The database are in substantially any form in which data are maintained and transmitted, but is preferably an electronic database. The electronic database of the invention are maintained on any electronic device allowing for the storage of and access to the database, such as a personal computer, but is preferably distributed on a wide area network, such as the World Wide Web.

The focus of the present section on databases that include peptide sequence data is for clarity of illustration only. It will be apparent to those of skill in the art that similar databases are assembled for any assay data acquired using an assay of the invention.

The compositions and methods for identifying and/or quantitating the relative and/or absolute abundance of a variety of molecular and macromolecular species from a biological sample undergoing ovarian cancer, i.e., the identification of ovarian cancer-associated sequences described herein, provide an abundance of information, which are correlated with pathological conditions, predisposition to disease, drug testing, therapeutic monitoring, gene-disease causal linkages, identification of correlates of immunity and physiological status, among others. Although the data generated from the assays of the invention is suited for manual review and analysis, in a preferred embodiment, prior data processing using high-speed computers is utilized.

An array of methods for indexing and retrieving biomolecular information is known in the art. For example, U.S. Patents 6,023,659 and 5,966,712 disclose a relational database system for storing biomolecular sequence information in a manner that allows sequences to be catalogued and searched according to one or more protein function hierarchies. U.S. Patent 5,953,727 discloses a relational database having sequence records containing information in a format that allows a collection of partial-length DNA sequences to be catalogued and searched according to association with one or more sequencing projects for obtaining full-length sequences from the collection of partial length sequences. U.S. Patent 5,706,498 discloses a gene database retrieval system for

5

10

30

35

making a retrieval of a gene sequence similar to a sequence data item in a gene database based on the degree of similarity between a key sequence and a target sequence. U.S. Patent 5,538,897 discloses a method using mass spectroscopy fragmentation patterns of peptides to identify amino acid sequences in computer databases by comparison of predicted mass spectra with experimentally-derived mass spectra using a closeness-of-fit measure. U.S. Patent 5,926,818 discloses a multi-dimensional database comprising a functionality for multi-dimensional data analysis described as on-line analytical processing (OLAP), which entails the consolidation of projected and actual data according to more than one consolidation path or dimension. U.S. Patent 5,295,261 reports a hybrid database structure in which the fields of each database record are divided into two classes, navigational and informational data, with navigational fields stored in a hierarchical topological map which are viewed as a tree structure or as the merger of two or more such tree structures.

See also Mount et al., Bioinformatics (2001); Biological Sequence Analysis: Probabilistic Models of Proteins and Nucleic Acids (Durbin et al., eds., 1999); Bioiraformatics: A Practical Guide to the Analysis of Genes and Proteins (Baxevanis & Oeullette eds., 1998)); Rashidi & Buehler, Bioinformatics: Basic Applications in Biological Science and Medicine (1999); Introduction to Computational Molecular Biology (Setubal et al., eds 1997); Bioinformatics: Methods and Protocols (Misener & Krawetz, eds, 2000); Bioinformatics: Sequence, Structure, and Databanks: A Practical Approach (Higgins & Taylor, eds., 2000); Brown, Bioinfor7natics: A Biologist's Guide to Biocomputing and the Internet (2001); Han & Kamber, Data Mining: Concepts and Techniques (2000); and Waterman, Introduction to Computational Biology: Maps, Sequences, and Genomes (1995).

The present invention provides a computer database comprising a computer and software for storing in computer-retrievable form assay data records cross-tabulated, e.g., with data specifying the source of the target-containing sample from which each sequence specificity record was obtained.

In an exemplary embodiment, at least one of the sources of target-containing sample is from a control tissue sample known to be free of pathological disorders. In a variation, at least one of the sources is a known pathological tissue specimen, e.g., a neoplastic lesion or another tissue specimen to be analyzed for prostate cancer. In another variation, the assay records cross-tabulate one or more of the following parameters for

each target species in a sample: (1) a unique identification code, which can include, e.g., a target molecular structure and/or characteristic separation coordinate (e.g., electrophoretic coordinates); (2) sample source; and (3) absolute and/or relative quantity of the target species present in the sample.

The invention also provides for the storage and retrieval of a collection of target data in a computer data storage apparatus, which can include magnetic disks, optical disks, magneto-optical disks, DRAM, SRAM, SGRAM, SDRAM, RDRAM, DDR RAM, magnetic bubble memory devices, and other data storage devices, including CPU registers and on-CPU data storage arrays. Typically, the target data records are stored as a bit pattern in an array of magnetic domains on a magnetizable medium or as an array of charge states or transistor gate states, such as an array of cells in a DRAM device (e.g., each cell comprised of a transistor and a charge storage area, which are on the transistor). In one embodiment, the invention provides such storage devices, and computer systems built therewith, comprising a bit pattern encoding a protein expression fingerprint record comprising unique identifiers for at least 10 target data records cross-tabulated with target source.

When the target is a peptide or nucleic acid, the invention preferably provides a method for identifying related peptide or nucleic acid sequences, comprising performing a computerised comparison between a peptide or nucleic acid sequence assay record stored in or retrieved from a computer storage device or database and at least one other sequence. The comparison can include a sequence analysis or comparison algorithm or computer program embodiment thereof (e.g., BLAST, FASTA, TFASTA, GAP, BESTFIT — see above) and/or the comparison are of the relative amount of a peptide or nucleic acid sequence in a pool of sequences determined from a polypeptide or nucleic acid sample of a specimen.

The invention also preferably provides a magnetic disk, such as an IBM-compatible (DOS, Windows, Windows95/.98/2000, Windows NT, OS/2) or other format (e.g., Linux, SunOS, Solaris, AIX, SCO Unix, VMS, MV, Macintosh, etc.) floppy diskette or hard (fixed, Winchester) disk drive, comprising a bit pattern encoding data from an assay of the invention in a file format suitable for retrieval and processing in a computerized sequence analysis, comparison, or relative quantitation method.

5

10

25

30.

35

The invention also provides a network, comprising a plurality of computing devices linked via a data link, such as an Ethernet cable (coax or IOBaseT), telephone line, ISDN line, wireless network, optical fiber, or other suitable signal transmission medium, whereby at least one network device (e.g., computer, disk array, etc.) comprises a pattern of magnetic domains (e.g., magnetic disk) and/or charge domains (e.g., an array of DRAM cells) composing a bit pattern encoding data acquired from an assay of the invention.

The invention also provides a method for transmitting assay data that includes generating an electronic signal on an electronic communications device, such as a modern, ISDN terminal adapter, DSL, cable modern, ATM switch, or the like, wherein the signal includes (in native or encrypted format) a bit pattern encoding data from an assay or a database comprising a plurality of assay results obtained by the method of the invention.

In a preferred embodiment, the invention provides a computer system for comparing a query target to a database containing an array of data structures, such as an assay result obtained by the method of the invention, and ranking database targets based on the degree of identity and gap weight to the target data. A central processor is preferably initialized to load and execute the computer program for alignment and/or comparison of the assay results. Data for a query target is entered into the central processor via an I/O device. Execution of the computer program results in the central processor retrieving the assay data from the data file, which comprises a binary description of an assay result.

The target data or record and the computer program are transferred to secondary memory, which is typically random access memory (e.g., DRAM, SRAM, SGRAM, or SDRAM). Targets are ranked according to the degree of correspondence between a selected assay characteristic (e.g., binding to a selected affinity moiety) and the same characteristic of the query target and results are output via an I/O device. For example, a central processor are a conventional computer (e.g., Intel Pentium, PowerPC, Alpha, PA-8000, SPARC, MIPS 4400, MIPS 10000, VAX, etc.); a program are a commercial or public domain molecular biology software package (e.g., UWGCG Sequence Analysis Software, Darwin); a data file are an optical or magnetic disk, a data server, a memory device (e.g., DRAM, SRAM, SGRAM, SDRAM, EPROM, bubble memory, flash memory, etc.); an I/O device are a terminal comprising a video display and a keyboard, a modem, an ISDN terminal adapter, an Ethernet port, a punched card reader, a magnetic strip reader, or other suitable I/O device.

The invention also preferably provides the use of a computer system, such as that described above, which comprises: (1) a computer; (2) a stored bit pattern encoding a collection of peptide sequence specificity records obtained by the methods of the invention, which are stored in the computer; (3) a comparison target, such as a query target; and (4) a program for alignment and comparison, typically with rank-ordering of comparison results on the basis of computed similarity values.

Transgenic Animals Expressing Ovarian Cancer-associated proteins and "Knock-Out" Animals

The present invention also contemplates transgenic animals which are transgenic by virtue of comprising a polynucleotide of the invention, i.e. animals transformed with a cancer-associated gene of the invention. Suitable animals are generally from the phylum chordata. Chordates includes vertebrate groups such as mammals, birds, reptiles and amphibians. Particular examples of mammals include non-human primates, cats, dogs, ungulates such as cows, goats, pigs, sheep and horses and rodents such as mice, rats, gerbils and hamsters. Transgenic animals within the meaning of the present invention are non-human animals and the production of transgenic humans is specifically excluded.

20

5

10

15

Techniques for producing transgenic animals are well known in the art. A useful general textbook on this subject is Houdebine, Transgenic animals – Generation and Use (Harwood Academic, 1997) – an extensive review of the techniques used to generate transgenic animals from fish to mice and cows.

25

30

35

Advances in technologies for embryo micromanipulation now permit introduction of heterologous DNA into, for example, fertilized mammalian ova. For instance, totipotent or pluripotent stem cells are transformed by microinjection, calcium phosphate mediated precipitation, liposome fusion, retroviral infection or other means, the transformed cells are then introduced into the embryo, and the embryo then develops into a transgenic animal. In a highly preferred method, developing embryos are infected with a retrovirus containing the desired DNA, and transgenic animals produced from the infected embryo. In a most preferred method, however, the appropriate DNAs are coinjected into the pronucleus or cytoplasm of embryos, preferably at the single cell stage, and the embryos allowed to develop into mature transgenic animals. Those techniques as well known. See reviews of standard laboratory procedures for microinjection of heterologous DNAs into

89

mammalian fertilized ova, including Hogan *et al.*, Manipulating the Mouse Embryo, (Cold Spring Harbor Press 1986); Krimpenfort *et al.*, Bio/Technology 9:844 (1991); Palmiter *et al.*, Cell, 41: 343 (1985); Kraemer *et al.*, Genetic manipulation of the Mammalian Embryo, (Cold Spring Harbor Laboratory Press 1985); Hammer *et al.*, Nature, 315: 680 (1985); Wagner *et al.*, U.S. Pat. No. 5,175,385; Krimpenfort *et al.*, U.S. Pat. No. 5,175,384, the respective contents of which are incorporated herein by reference

Another method used to produce a transgenic animal involves microinjecting a nucleic acid into pro-nuclear stage eggs by standard methods. Injected eggs are then cultured before transfer into the oviducts of pseudopregnant recipients.

Transgenic animals may also be produced by nuclear transfer technology as described in Schnieke, A.E. *et al.*, 1997, Science, 278: 2130 and Cibelli, J.B. *et al.*, 1998, Science, 280: 1256. Using this method, fibroblasts from donor animals are stably transfected with a plasmid incorporating the coding sequences for a binding domain or binding partner of interest under the control of regulatory. Stable transfectants are then fused to enucleated oocytes, cultured and transferred into female recipients.

Analysis of animals which may contain transgenic sequences would typically be performed by either PCR or Southern blot analysis following standard methods.

By way of a specific example for the construction of transgenic mammals, such as cows, nucleotide constructs comprising a sequence encoding a binding domain fused to GFP are microinjected using, for example, the technique described in U.S. Pat. No. 4,873,191, into oocytes which are obtained from ovaries freshly removed from the mammal. The oocytes are aspirated from the follicles and allowed to settle before fertilization with thawed frozen sperm capacitated with heparin and prefractionated by Percoll gradient to isolate the motile fraction.

The fertilized oocytes are centrifuged, for example, for eight minutes at 15,000 g to visualize the pronuclei for injection and then cultured from the zygote to morula or blastocyst stage in oviduct tissue-conditioned medium. This medium is prepared by using luminal tissues scraped from oviducts and diluted in culture medium. The zygotes must be placed in the culture medium within two hours following microinjection.

5

10

15

25

Oestrous is then synchronized in the intended recipient mammals, such as cattle, by administering coprostanol. Oestrous is produced within two days and the embryos are transferred to the recipients 5-7 days after estrous. Successful transfer are evaluated in the offspring by Southern blot.

90

PCT/AU2003/001166

5

10

15

20

25

Alternatively, the desired constructs are introduced into embryonic stem cells (ES cells) and the cells cultured to ensure modification by the transgene. The modified cells are then injected into the blastula embryonic stage and the blastulas replaced into pseudopregnant hosts. The resulting offspring are chimeric with respect to the ES and host cells, and nonchimeric strains which exclusively comprise the ES progeny are obtained using conventional cross-breeding. This technique is described, for example, in WO91/10741.

In another embodiment, transgenic animals of the present invention are transgenic "knock-out" animals where a specific gene corresponding to a polynucleotide referred to in Tables 1-3 has been rendered non-functional by homologous recombination. The generation of "knock-out" animals is similar to the production of other transgenic animals except that the polynucleotide constructs are designed to integrate into the endogenous genes and disrupt the function of the endogenous sequences. The generation of "knock-out" animals is known in the art, including the design of suitable constructs that will recombine at the appropriate site in the genome.

In one embodiment, the heterologous sequence which it is desired to recombine into the genome of a target animal comprises a functional sequence but under the control of an inducible promoter so that expression of the gene are regulated by administration of an endogenous molecule. This are advantageous where disruption of the gene is embryonic-lethal.

"Knock-out" animals are used as animal models for the study of gene function.

30

35

## Therapeutic peptides

In accordance with this embodiment, ovarian cancer-associated proteins of the present invention are administered therapeutically to patients for a time and under conditions sufficient to ameliorate the growth of a tumor in the subject or to prevent tumor recurrence.

It is preferred to use peptides that do not consisting solely of naturally-occurring amino acids but which have been modified, for example to reduce immunogenicity, to increase circulatory half-life in the body of the patient, to enhance bioavailability and/or to enhance efficacy and/or specificity.

5

A number of approaches have been used to modify peptides for therapeutic application. One approach is to link the peptides or proteins to a variety of polymers, such as polyethylene glycol (PEG) and polypropylene glycol (PPG) – see for example U.S. Patent Nos. 5,091,176, 5,214,131 and US 5,264,209.

10

30

35

Replacement of naturally-occurring amino acids with a variety of uncoded or modified amino acids such as D-amino acids and N-methyl amino acids may also be used to modify peptides

Another approach is to use bifunctional crosslinkers, such as N-succinimidyl 3-(2 pyridyldithio) propionate, succinimidyl 6-[3-(2 pyridyldithio) propionamido] hexanoate, and sulfosuccinimidyl 6-[3-(2 pyridyldithio) propionamido]hexanoate (see US Patent 5,580,853).

It are desirable to use derivatives of the ovarian cancer-associated proteins of the invention which are conformationally constrained. Conformational constraint refers to the stability and preferred conformation of the three-dimensional shape assumed by a peptide. Conformational constraints include local constraints, involving restricting the conformational mobility of a single residue in a peptide; regional constraints, involving restricting the conformational mobility of a group of residues, which residues may form some secondary structural unit; and global constraints, involving the entire peptide structure.

The active conformation of the peptide are stabilized by a covalent modification, such as cyclization or by incorporation of gamma-lactam or other types of bridges. For example, side chains are cyclized to the backbone so as create a L-gamma-lactam moiety on each side of the interaction site. See, generally, Hruby et al., "Applications of Synthetic Peptides," in Synthetic Peptides: A User's Guide: 259-345 (W. H. Freeman & Co. 1992). Cyclization also are achieved, for example, by formation of cystine bridges, coupling of amino and carboxy terminal groups of respective terminal amino acids, or coupling of the amino group of a Lys residue or a related homolog with a carboxy group of Asp, Glu or a

5

10

15

20

25

30

related homolog. Coupling of the .alpha-amino group of a polypeptide with the epsilon-amino group of a lysine residue, using iodoacetic anhydride, are also undertaken. See Wood and Wetzel, 1992, Int'l J. Peptide Protein Res. 39: 533-39.

Another approach described in US 5,891,418 is to include a metal-ion complexing backbone in the peptide structure. Typically, the preferred metal-peptide backbone is based on the requisite number of particular coordinating groups required by the coordination sphere of a given complexing metal ion. In general, most of the metal ions that may prove useful have a coordination number of four to six. The nature of the coordinating groups in the peptide chain includes nitrogen atoms with amine, amide, imidazole, or guanidino functionalities; sulfur atoms of thiols or disulfides; and oxygen atoms of hydroxy, phenolic, carbonyl, or carboxyl functionalities. In addition, the peptide chain or individual amino acids are chemically altered to include a coordinating group, such as for example oxime, hydrazino, sulfhydryl, phosphate, cyano, pyridino, piperidino, or morpholino. The peptide construct are either linear or cyclic, however a linear construct is typically preferred. One example of a small linear peptide is Gly-Gly-Gly which has four nitrogens (an N<sub>4</sub> complexation system) in the back bone that can complex to a metal ion with a coordination number of four.

A further technique for improving the properties of therapeutic peptides is to use non-peptide peptidomimetics. A wide variety of useful techniques are used to elucidating the precise structure of a peptide. These techniques include amino acid sequencing, x-ray crystallography, mass spectroscopy, nuclear magnetic resonance spectroscopy, computer-assisted molecular modeling, peptide mapping, and combinations thereof. Structural analysis of a peptide generally provides a large body of data which comprise the amino acid sequence of the peptide as well as the three-dimensional positioning of its atomic components. From this information, non-peptide peptidomimetics are designed that have the required chemical functionalities for therapeutic activity but are more stable, for example less susceptible to biological degradation. An example of this approach is provided in US 5,811,512.

Techniques for chemically synthesising therapeutic peptides of the invention are described in the above references and also reviewed by Borgia and Fields, 2000, TibTech 18: 243-251 and described in detail in the references contained therein.

The ovarian cancer proteins, nucleic acids, and antibodies as described herein are used in drug screening assays to identify candidate compounds for use in treating ovarian cancer. The ovarian cancer-associated proteins, antibodies, nucleic acids, modified proteins and cells containing ovarian cancer sequences are used in drug screening assays or by evaluating the effect of drug candidates on a "gene expression profile" or expression profile of polypeptides. In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent (e.g., Zlokarnik, et al., 1998, Science 279: 84-88); Heid, 1996, Genome Res 6: 986-94).

In a preferred embodiment, the ovarian cancer-associated proteins, antibodies, nucleic acids, modified proteins and cells containing the native or modified ovarian cancer-associated proteins are used in screening assays. That is, the present invention provides methods for screening for compounds/agents which modulate the ovarian cancer phenotype or an identified physiological function of a ovarian cancer-associated protein. As above, this are done on an individual gene level or by evaluating the effect of drug candidates on a "gene expression profile". In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent, see Zlokarnik, supra.

Having identified the differentially expressed genes herein, a variety of assays are executed. In a preferred embodiment, assays are run on an individual gene or protein level. That is, having identified a particular gene as up regulated in ovarian cancer, test compounds are screened for the ability to modulate gene expression or for binding to the ovarian cancer-associated protein. "Modulation" thus includes both an increase and a decrease in gene expression. The preferred amount of modulation will depend on the original change of the gene expression in normal versus tissue undergoing ovarian cancer, with changes of at least 10%, preferably 50%, more preferably 100-300%, and in some embodiments 300-1000% or greater. Thus, if a gene exhibits a 4-fold increase in ovarian cancer tissue compared to normal tissue, a decrease of about four-fold is often desired; similarly, a 10-fold decrease in ovarian cancer tissue compared to normal tissue often provides a target value of a 10-fold increase in expression to be induced by the test compound.

The amount of gene expression are monitored using nucleic acid probes and the quantification of gene expression levels, or, alternatively, the gene product itself are monitored, e.g., through the use of antibodies to the ovarian cancer-associated protein and standard immunoassays. Proteomics and separation techniques may also allow quantification of expression.

In a preferred embodiment, gene expression or protein monitoring of a number of entities, i.e., an expression profile, is monitored simultaneously. Such profiles will typically involve a plurality of those entities described herein.

10

15

20

5

In this embodiment, the ovarian cancer nucleic acid probes are attached to biochips as outlined herein for the detection and quantification of ovarian cancer sequences in a particular cell. Alternatively, PCR are used. Thus, a series are used with dispensed primers in desired wells. A PCR reaction can then be performed and analyzed for each well.

Expression monitoring are performed to identify compounds that modify the expression of one or more ovarian cancer-associated sequences, e.g., a polynucleotide sequence set out in Tables 1-3. In a preferred embodiment, a test modulator is added to the cells prior to analysis. Moreover, screens are also provided to identify agents that modulate ovarian cancer, modulate ovarian cancer-associated proteins, bind to a ovarian cancer-associated protein, or interfere with the binding of a ovarian cancer-associated protein and an antibody or other binding partner.

The term "test compound" or "drug candidate" or "modulator" or grammatical equivalents as used herein describes any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for the capacity to directly or indirectly alter the ovarian cancer phenotype or the expression of a ovarian cancer sequence, e.g., a nucleic acid or protein sequence. In preferred embodiments, modulators alter expression profiles, or expression profile nucleic acids or proteins provided herein. In one embodiment, the modulator suppresses a ovarian cancer phenotype, e.g. to a normal tissue fingerprint. In another embodiment, a modulator induced a ovarian cancer phenotype. Generally, a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e.,

at zero concentration or below the level of detection.

5

10

20

25

30

Drug candidates encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Preferred small molecules are less than 2000, or less than 1500 or less than 1000 or less than 500 Daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides.

In one aspect, a modulator will neutralize the effect of a ovarian cancer-associated protein. By "neutralize" is meant that activity of a protein is inhibited or blocked and the consequent effect on the cell.

In certain embodiments, combinatorial libraries of potential modulators will be screened for an ability to bind to a ovarian cancer polypeptide or to modulate activity. Conventionally, new chemical entities with useful properties are generated by identifying a chemical compound (called a "lead compound") with some desirable property or activity, e.g., inhibiting activity, creating variants of the lead compound, and evaluating the property and activity of those variant compounds. Often, high throughput screening (HTS) methods are employed for such an analysis.

In one preferred embodiment, high throughput screening methods involve providing a library containing a large number of potential therapeutic compounds (candidate compounds). Such "combinatorial chemical libraries" are then screened in one or more assays to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis by combining a number of chemical

"building blocks" such as reagents. For example, a linear combinatorial chemical library, such as a polypeptide (e.g., mutein) library, is formed by combining a set of chemical building blocks called amino acids in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds are synthesized through such combinatorial mixing of chemical building blocks (Gallop et al., 1994, J. Med. Chem. 37(9):1233-1251).

5

10

30

35

Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries, peptoids, encoded peptides, random bio-oligomers, nonpeptidal peptidomimetics, analogous organic syntheses of small compound libraries, nucleic acid libraries, peptide nucleic acid libraries, antibody libraries, carbohydrate libraries and small organic molecule libraries.

The assays to identify modulators are amenable to high throughput screening. Preferred assays thus detect enhancement or inhibition of ovarian cancer gene transcription, inhibition or enhancement of polypeptide expression, and inhibition or enhancement of polypeptide activity.

High throughput assays for the presence, absence, quantification, or other properties of particular nucleic acids or protein products are well known to those of skill in the art. Similarly, binding assays and reporter gene assays are similarly well known. Thus, e.g., U.S. Patent No. 5,559,410 discloses high throughput screening methods for proteins, U.S. Patent No. 5,585,639 discloses high throughput screening methods for nucleic acid binding (i.e., in arrays), while U.S. Patent Nos. 5,576,220 and 5,541,061 disclose high throughput methods of screening for ligand/antibody binding.

In addition, high throughput screening systems are commercially available (see, e.g., Zymark Corp., Hopkinton, MA; Air Technical Industries, Mentor, OH; Beckman Instruments, Inc. Fullerton, CA; Precision Systems, Inc., Natick, MA, etc.). These systems typically automate entire procedures, including all samlsle and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detectors) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. The manufacturers of such systems provide detailed protocols for various high throughput systems. Thus, e.g.,

97

Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like.

In one embodiment, modulators are proteins, often naturally occurring proteins or fragments of naturally occurring proteins. Thus, e.g., cellular extracts containing proteins, or random or directed digests of proteinaceous cellular extracts, are used. In this way libraries of proteins are made for screening in the methods of the invention. Particularly preferred in this embodiment are libraries of bacterial, fungal, viral, and mammalian proteins, with the latter being preferred, and human proteins being especially preferred. Particularly useful test compound will be directed to the class of proteins to which the target belongs, e.g., substrates for enzymes or ligands and receptors.

In a preferred embodiment, modulators are peptides of from about 5 to about 30 amino acids, with from about 5 to about 20 amino acids being preferred, and from about 7 to about 15 being particularly preferred. The peptides are digests of naturally occurring proteins as is outlined above, random peptides, or "biased" random peptides. By "randomized" or grammatical equivalents herein is meant that each nucleic acid and peptide consists of essentially random nucleotides and amino acids, respectively. Since generally these random peptides (or nucleic acids, discussed below) are chemically synthesized, they may incorporate any nucleotide or amino acid at any position. The synthetic process are designed to generate randomized proteins or nucleic acids, to allow the formation of all or most of the possible combinations over the length of the sequence, thus forming a library of randomized candidate bioactive proteinaceous agents.

25

30

10

15

20

In one embodiment, the library is fully randomized, with no sequence preferences or constants at any position. In a preferred embodiment, the library is biased. That is, some positions within the sequence are either held constant, or are selected from a limited number of possibilities. For example, in a preferred embodiment, the nucleotides or amino acid residues are randomized within a defined class, e.g., of hydrophobic amino acids, hydrophilic residues, sterically blased (either small or large) residues, towards the creation of nucleic acid binding domains, the creation of cysteines, for cross-linking, prolines for SH-3 domains, serines, threonines, tyrosines or histidines for phosphorylation sites, etc., or to purines, etc.

Modulators of ovarian cancer can also be nucleic acids, as defined below. As described above generally for proteins, nucleic acid modulating agents are naturally occurring nucleic acids, random nucleic acids, or "biased" random nucleic acids. For example, digests of procaryotic or eucaryotic genomes are used as is outlined above for proteins.

5

10

15

20

25

In certain embodiments, the activity of a ovarian cancer-associated protein is down-regulated, or entirely inhibited, by the use of antisense polynucleotide, *i.e.*, a nucleic acid complementary to, and which can preferably hybridize specifically to, a coding mRNA nucleic acid sequence, e.g., a ovarian cancer-associated protein mRNA, or a subsequence thereof. Binding of the antisense polynucleotide to the mRNA reduces the translation and/or stability of the mRNA.

In the context of this invention, antisense polynucleotides can comprise naturally-occurring nucleotides, or synthetic species formed from naturally-occurring subunits or their close homologs. Antisense polynucleotides may also have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur containing species which are known for use in the art. Analogs are comprehended by this invention so long as they function effectively to hybridize with the ovarian cancer-associated protein mRNA. See, e.g., Isis Pharmaceuticals, Carlsbad, CA; Sequitor, Inc., Natick, MA.

Such antisense polynucleotides can readily be synthesized using recombinant means, or are synthesized *in vitro*. Equipment for such synthesis is sold by several vendors, including Applied Biosystems. The preparation of other oligonucleotides such as phosphorothioates and alkylated derivatives is also well known to those of skill in the art.

30

35

Antisense molecules as used herein include antisense or sense oligonucleotides. Sense oligonucleotides can, e.g., be employed to block transcription by binding to the anti-sense strand. The antisense and sense oligonucleotide comprise a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target mRNA (sense) or DNA (antisense) sequences for ovarian cancer molecules. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment generally at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, e.g., Stein & Cohen (Cancer Res. 48:2659 (1988 and van der Krol et al. (BioTechniques 6:958 (1988)).

In addition to antisense polynucleotides, ribozymes are used to target and inhibit transcription of ovarian cancer-associated nucleotide sequences. A ribozyme is an RNA molecule that catalytically cleaves other RNA molecules. Different kinds of ribozymes have been described, including group I ribozymes, hammerhead ribozymes, hairpin ribozymes, RNase P, and axhead ribozymes (see, e.g., Castanotto et al., Adv. in Pharmacology 25: 289-317 (1994) for a general review of the properties of different 5 ribozymes).

Methods of preparing ribozymes are well known to those of skill in the art (see, e.g., WO 94/26877; Ojwang et al., Proc. Natl. Acad. Sci. USA 90:6340-6344 (1993); Yamada et al., Human Gene Therapy 1:39-45 (1994); Leavitt et al., Proc. Natl. Acad. Sci. USA 92:699- 703 (1995); Leavitt et al., Human Gene Therapy 5:1151-120 (1994); and Yamada et al., Virology 205: 121-126 (1994)).

15

20

25

5

Polynucleotide modulators of ovarian cancer are introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell. Alternatively, a polynucleotide modulator of ovarian cancer are introduced into a cell containing the target nucleic acid sequence, e.g., by formation of an polynucleotide-lipid complex, as described in WO 90/10448. It is understood that the use of antisense molecules or knock out and knock in models may also be used in screening assays as discussed above, in addition to methods of treatment.

As noted above, gene expression monitoring is conveniently used to test candidate modulators (e.g., protein, nucleic acid or small molecule). After the candidate agent has been added and the cells allowed to incubate for some period of time, the sample containing a target sequence to be analyzed is added to the biochip. If required, the target sequence is prepared using known techniques. For example, the sample are treated to lyse the cells, using known lysis buffers, electroporation, etc., with purification and/or amplification such as PCR performed as appropriate. For example, an *in vitro* 

transcription with labels covalently attached to the nucleotides is performed. Generally, the nucleic acids are labeled with biotin-FITC or PE, or with cy3 or cy5.

In a preferred embodiment, the target sequence is labeled with, e.g., a fluorescent, a chemiluminescent, a chemical, or a radioactive signal, to provide a means of detecting the target sequence's specific binding to a probe. The label also are an enzyme, such as, alkaline phosphatase or horseradish peroxidase, which when provided with an appropriate substrate produces a product that are detected. Alternatively, the label are a labeled compound or small molecule, such as an enzyme inhibitor, that binds but is not catalyzed or altered by the enzyme. The label also are a moiety or compound, such as, an epitope tag or biotin which specifically binds to streptavidin. For the example of biotin, the streptavidin is labeled as described above, thereby, providing a detectable signal for the bound target sequence. Unbound labeled streptavidin is typically removed prior to analysis.

15

20

35

5

10

As will be appreciated by those in the art, these assays are direct hybridization assays or can comprise "sandwich assays", which include the use of multiple probes, as is generally outlined in U.S. Patent Nos. 5,681,702, 5,597,909, 5,545,730, 5,594,117, 5,591,584, 5,571,670, 5,580,731, 5,571,670, 5,591,584, 5,624,802, 5,635,352, 5,594,118, 5,359,100, 5,124,246 and 5,681,697, all of which are hereby incorporated by reference. In this embodiment, in general, the target nucleic acid is prepared as outlined above, and then added to the biochip comprising a plurality of nucleic acid probes, under conditions that allow the formation of a hybridization complex.

A variety of hybridization conditions are used in the present invention, including high, moderate and low stringency conditions as outlined above. The assays are generally run under stringency conditions which allows formation of the label probe hybridization complex only in the presence of target. Stringency are controlled by altering a step parameter that is a thermodynamic variable, including, but not limited to, temperature, formamide concentration, salt concentration, chaotropic salt concentration pH, organic solvent concentration, etc.

These parameters may also be used to control non-specific binding, as is generally outlined in U.S. Patent No. 5,681,697. Thus it are desirable to perform certain steps at higher stringency conditions to reduce non-specific binding.

The reactions outlined herein are accomplished in a variety of ways. Components of the reaction are added simultaneously, or sequentially, in different orders, with preferred embodiments outlined below. In addition, the reaction may include a variety of other reagents. These include salts, buffers, neutral proteins, e.g. albumin, detergents, etc. which are used to facilitate optimal hybridization and detection, and/or reduce non-specific or background interactions. Reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may also be used as appropriate, depending on the sample preparation methods and purity of the target.

The assay data are analyzed to determine the expression levels, and changes in expression levels as between states, of individual genes, forming a gene expression profile.

Screens are performed to identify modulators of the ovarian cancer phenotype. In one embodiment, screening is performed to identify modulators that can induce or suppress a particular expression profile, thus preferably generating the associated phenotype. In another embodiment, e.g., for diagnostic applications, having identified differentially expressed genes important in a particular state, screens are performed to identify modulators that alter expression of individual genes. In an another embodiment, screening is performed to identify modulators that alter a biological function of the expression product of a differentially expressed gene. Again, having identified the importance of a gene in a particular state, screens are performed to identify agents that bind and/or modulate the biological activity of the gene product.

In addition screens are done for genes that are induced in response to a candidate agent. After identifying a modulator based upon its ability to suppress a ovarian cancer expression pattern leading to a normal expression pattern, or to modulate a single ovarian cancer gene expression profile so as to mimic the expression of the gene from normal tissue, a screen as described above are performed to identify genes that are specifically modulated in response to the agent. Comparing expression profiles between normal tissue and agent treated ovarian cancer tissue reveals genes that are not expressed in normal tissue or ovarian cancer tissue, but are expressed in agent treated tissue. These agent-specific sequences are identified and used by methods described herein for ovarian cancer genes or proteins. In particular these sequences and the proteins they encode find use in marking or identifying agent treated cells. In addition,

antibodies are raised against the agent induced proteins and used to target novel therapeutics to the treated ovarian cancer tissue sample.

Thus, in one embodiment, a test compound is administered to a population of ovarian cancer cells, that have an associated ovarian cancer expression profile. By "administration" or "contacting" herein is meant that the candidate agent is added to the cells in such a manner as to allow the agent to act upon the cell, whether by uptake and intracellular action, or by action at the cell surface. In some embodiments, nucleic acid encoding a proteinaceous candidate agent (i.e., a peptide) are put into a viral construct such as an adenoviral or retroviral construct, and added to the cell, such that expression of the peptide agent is accomplished. Regulatable gene administration systems can also be used.

5

10

15

20

25

30

35

Once the test compound has been administered to the cells, the cells are washed if desired and are allowed to incubate under preferably physiological conditions for some period of time. The cells are then harvested and a new gene expression profile is generated, as outlined herein.

Thus, e.g., ovarian cancer tissue are screened for agents that modulate, e.g., induce or suppress the ovarian cancer phenotype. A change in at least one gene, preferably many, of the expression profile indicates that the agent has an effect on ovarian cancer activity. By defining such a signature for the ovarian cancer phenotype, screens for new drugs that alter the phenotype are devised. With this approach, the drug target need not be known and need not be represented in the original expression screening platform, nor does the level of transcript for the target protein need to change.

In a preferred embodiment, as outlined above, screens are done on individual genes and gene products (proteins). That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of either the expression of the gene or the gene product itself are done. The gene products of differentially expressed genes are sometimes referred to herein as "ovarian cancer-associated proteins" or a "ovarian cancer modulatory protein". The ovarian cancer modulatory protein are a fragment, or alternatively, be the full length protein to the fragment encoded by the nucleic acids referred to in Tables 1-3. Preferably, the ovarian cancer modulatory protein is a fragment. In a preferred embodiment, the ovarian cancer amino acid sequence which is used to determine sequence identity or similarity is encoded by a

103

nucleic acid referred to in Tables 1-3. In another embodiment, the sequences are naturally occurring allelic variants of a protein encoded by a nucleic acid referred to in Tables 1-3. In another embodiment, the sequences are sequence variants as further described herein.

5

10

15

20

25

Preferably, the ovarian cancer modulatory protein is a fragment of approximately 14 to 24 amino acids long. More preferably the fragment is a soluble fragment. Preferably, the fragment includes a non-transmembrane region. In a preferred embodiment, the fragment has an N-terminal Cys to aid in solubility. In one embodiment, the C-terminus of the fragment is kept as a free acid and the N-terminus is a free amine to aid in coupling, i.e., to cysteine.

In one embodiment the ovarian cancer-associated proteins are conjugated to an immunogenic agent as discussed herein. In one embodiment the ovarian cancer-associated protein is conjugated to BSA.

Measurements of ovarian cancer polypeptide activity, or of ovarian cancer or the ovarian cancer phenotype are performed using a variety of assays. For example, the effects of the test compounds upon the function of the ovarian cancer polypeptides are measured by examining parameters described above. A suitable physiological change that affects activity are used to assess the influence of a test compound on the polypeptides of this invention. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as, in the case of ovarian cancer associated with tumours, tumour growth, tumour metastasis, neovascularization, hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as cGMP. In tire assays of the invention, mammalian ovarian cancer polypeptide is typically used, e.g., mouse, preferably human.

30

35

Assays to identify compounds with modulating activity are performed *in* vitro. For example, a ovarian cancer polypeptide is first contacted with a potential modulator and incubated for a suitable amount of time, e.g., from 0.5 to 48 hours. In one embodiment, the ovarian cancer polypeptide levels are determined *in vitro* by measuring the level of protein or mRNA. The level of protein is measured using immunoassays such as western blotting, ELISA and the like with an antibody that selectively binds to the ovarian cancer

polypeptide or a fragment thereof. For measurement of mRNA, amplification, e.g., using PCR, LCR, or hybridization assays, e.g., northern hybridization, RNAse protection, dot blotting, are preferred. The level of protein or mRNA is detected using directly or indirectly labeled detection agents, e.g., fluorescently or radioactively labeled nucleic acids, radioactively or enzymatically labeled antibodies, and the like, as described herein.

Alternatively, a reporter gene system are devised using the ovarian cancer-associated protein promoter operably linked to a reporter gene such as luciferase, green fluorescent protein, CAT, or (beta-gal. The reporter construct is typically transfected into a cell. After treatment with a potential modulator, the amount of reporter gene transcription, translation, or activity is measured according to standard techniques known to those of skill in the art.

In a preferred embodiment, as outlined above, screens are done on individual genes and gene products (proteins). That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of the expression of the gene or the gene product itself are done. The gene products of differentially expressed genes are sometimes referred to herein as "ovarian cancer-associated proteins." The ovarian cancer-associated protein are a fragment, or alternatively, be the full length protein to a fragment shown herein.

In one embodiment, screening for modulators of expression of specific genes is performed. Typically, the expression of only one or a few genes are evaluated. In another embodiment, screens are designed to first find compounds that bind to differentially expressed proteins. These compounds are then evaluated for the ability to modulate differentially expressed activity. Moreover, once initial candidate compounds are identified, variants are further screened to better evaluate structure activity relationships.

In a preferred embodiment, binding assays are done. In general, purified or isolated gene product is used; that is, the gene products of one or more differentially expressed nucleic acids are made. For example, antibodies are generated to the protein gene products, and standard immunoassays are run to determine the amount of protein present. Alternatively, cells comprising the ovarian cancer-associated proteins are used in the assays.

5

10

15

20

25

30

10

15

20

25

30

35

Thus, in a preferred embodiment, the methods comprise combining a ovarian cancer-associated protein and a candidate compound, and determining the binding of the compound to the ovarian cancer-associated protein. Preferred embodiments utilize the human ovarian cancer-associated protein, although other mammalian proteins may also be used, e.g. for the development of animal models of human disease. In some embodiments, as outlined herein, variant or derivative ovarian cancer-associated proteins are used.

Generally, in a preferred embodiment of the methods herein, the ovarian cancerassociated protein or the candidate agent is non-diffusably bound to an insoluble support having isolated sample receiving areas (e.g. a microtiter plate, an array, etc.). The insoluble supports are made of any composition to which the compositions are bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports are solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, teflon™, etc. microtitre plates and arrays are especially convenient because a large number of assays are carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusable. Preferred methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or agent, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

In a preferred embodiment, the ovarian cancer-associated protein is bound to the support, and a test compound is added to the assay. Alternatively, the candidate agent is bound to the support and the ovarian cancer-associated protein is added. Novel binding agents include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for agents that have a low toxicity for human cells. A wide variety of assays are used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility

PCT/AU2003/001166

5

10

15

20

shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

The determination of the binding of the test modulating compound to the ovarian cancer-associated protein are done in a number of ways. In a preferred embodiment, the compound is labeled, and binding determined directly, e.g., by attaching all or a portion of the ovarian cancer-associated protein to a solid support, adding a labeled candidate agent (e.g., a fluorescent label), washing off excess reagent, and determining whether the label is present on the solid support. Various blocking and washing steps are utilized as appropriate.

In some embodiments, only one of the components is labeled, e.g., the proteins (or proteinaceous candidate compounds) are labeled. Alternatively, more than one component are labeled with different labels, e.g., <sup>125</sup>I for the proteins and a fluorophor for the compound. Proximity reagents, e.g., quenching or energy transfer reagents are also useful.

In one embodiment, the binding of the test compound is determined by competitive binding assay. The competitor is a binding moiety known to bind to the target molecule (i.e., a ovarian cancer-associated protein), such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there are competitive binding between the compound and the binding moiety, with the binding moiety displacing the compound. In one embodiment, the test compound is labeled. Either the compound, or the competitor, or both, is added first to the protein for a time sufficient to allow binding, if present. Incubations are performed at a temperature which facilitates optimal activity, typically between 4 and 40°C. Incubation periods are typically optimized, e.g., to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

30

35

25

In a preferred embodiment, the competitor is added first, followed by the test compound. Displacement of the competitor is an indication that the test compound is binding to the ovarian cancer-associated protein and thus is capable of binding to, and potentially modulating, the activity of the ovarian cancer-associated protein. In this embodiment, either component are labeled. Thus, e.g., if the competitor is labeled, the presence of

label in the wash solution indicates displacement by the agent. Alternatively, if the test compound is labeled, the presence of the label on the support indicates displacement.

In an alternative preferred embodiment, the test compound is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate that the test compound is bound to the ovarian cancer-associated protein with a higher affinity. Thus, if the test compound is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate that the test compound is capable of binding to the ovarian cancer-associated protein.

10

15

20

25

5

In a preferred embodiment, the methods comprise differential screening to identity agents that are capable of modulating the activity of the ovarian cancer-associated proteins. In this embodiment, the methods comprise combining a ovarian cancer-associated protein and a competitor in a first sample. A second sample comprises a test compound, a ovarian cancer-associated protein, and a competitor. The binding of the competitor is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to the ovarian cancer-associated protein and potentially modulating its activity. That is, if the binding of the competitor is different in the second sample relative to the first sample, the agent is capable of binding to the ovarian cancer-associated protein.

Alternatively, differential screening is used to identify drug candidates that bind to the native ovarian cancer-associated protein, but cannot bind to modified ovarian cancer-associated proteins. The structure of the ovarian cancer-associated protein are modeled, and used in rational drug design to synthesize agents that interact with that site. Drug candidates that affect the activity of a ovarian cancer-associated protein are also identified by screening drugs for the ability to either enhance or reduce the activity of the protein.

Positive controls and negative controls are used in the assays. Preferably control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples are counted in a scintillation counter to determine

the amount of bound compound.

A variety of other reagents are included in the screening assays. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc. which are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., are used. The mixture of components are added in an order that provides for the requisite binding.

In a preferred embodiment, the invention provides methods for screening for a compound capable of modulating the activity of a ovarian cancer-associated protein. The methods comprise adding a test compound, as defined above, to a cell comprising ovarian cancer-associated proteins. Preferred cell types include almost any cell. The cells contain a recombinant nucleic acid that encodes a ovarian cancer-associated protein. In a preferred embodiment, a library of candidate agents are tested on a plurality of cells.

15

20

25

30

10

5

In one aspect, the assays are evaluated in the presence or absence or previous or subsequent exposure of physiological signals, e.g. hormones, antibodies, peptides, antigens, cytokines, growth factors, action potentials, pharmacological agents including chemotherapeutics, radiation, carcinogenics, or other cells (i.e. cell-cell contacts). In another example, the determinations are determined at different stages of the cell cycle process.

In this way, compounds that modulate ovarian cancer agents are identified. Compounds with pharmacological activity are able to enhance or interfere with the activity of the ovarian cancer-associated protein. Once identified, similar structures are evaluated to identify critical structural feature of the compound.

In one embodiment, a method of inhibiting ovarian cancer cell division is provided. The method comprises administration of a ovarian cancer inhibitor. In another embodiment, a method of inhibiting ovarian cancer is provided. The method comprises administration of a ovarian cancer inhibitor. In a further embodiment, methods of treating cells or individuals with ovarian cancer are provided. The method comprises administration of a ovarian cancer inhibitor.

In one embodiment, a ovarian cancer inhibitor is an antibody as discussed above. In another embodiment, the ovarian cancer inhibitor is an antisense molecule.

10

15

20

25

30

A variety of cell growth, proliferation, and metastasis assays are known to those of skill in the art, as described below.

5 Soft agar growth or colony formation in suspension

Normal cells require a solid substrate to attach and grow. When the cells are transformed, they lose this phenotype and grow detached from the substrate. For example, transformed cells can grow in stirred suspension culture or suspended in semi-solid media, such as semi-solid or soft agar. The transformed cells, when transfected with tumour suppressor genes, regenerate normal phenotype and require a solid substrate to attach and grow. Soft agar growth or colony formation in suspension assays are used to identify modulators of ovarian cancer sequences, which when expressed in host cells, inhibit abnormal cellular proliferation and transformation. A therapeutic compound would reduce or eliminate the host cells' ability to grow in stirred suspension culture or suspended in semisolid media, such as semi-solid or soft.

Techniques for soft agar growth or colony formation in suspension assays are described in Freshney, *Culture of Animal Cells a Manual of Basic Technique* (3rd ed., 1994), herein incorporated by reference. *See also*, the methods section of Garkavtsev *et al.* (1996), supra, herein incorporated by reference.

Contact inhibition and density limitation of growth

Normal cells typically grow in a flat and organized pattern in a petri dish until they touch other cells. When the cells touch one another, they are contact inhibited and stop growing. When cells are transformed, however, the cells are not contact inhibited and continue to grow to high densities in disorganized foci. Thus, the transformed cells grow to a higher saturation density than normal cells. This are detected morphologically by the formation of a disoriented monolayer of cells or rounded cells in foci within the regular pattern of normal surrounding cells. Alternatively, labeling index with (<sup>3</sup>H)-thymidine at saturation density are used to measure density limitation of growth. See Freshney (1994), supra. The transformed cells, when transfected with tumour suppressor genes, regenerate a normal phenotype and become contact inhibited and would grow to a lower density.

In this assay, labeling index with (<sup>3</sup>H)-thymidine at saturation density is a preferred method of measuring density limitation of growth. Transformed host cells are transfected

10

15

20

30

35

with a ovarian cancer-associated sequence and are grown for 24 hours at saturation density in non-limiting medium conditions. The percentage of cells labeling with (<sup>3</sup>H)-thymidine is determined autoradiographically. See, Freshney (1994), supra.

## 5 Growth factor or serum dependence

Transformed cells have a lower serum dependence than their normal counterparts (see, e.g., Temin, J. Natl. Cancer Insti. 37:167-175 (1966); Eagle et al., J. Exp. Med. 131:836-879 (1970)); Freshney, supra. This is in part due to release of various growth factors by the transformed cells. Growth factor or serum dependence of transformed host cells are compared with that of control. Tumor specific markers levels Tumor cells release an increased amount of certain factors (hereinafter "tumour specific markers") than their normal counterparts. For example, plasminogen activator (PA) is released from human glioma at a higher level than from normal brain cells (see, e.g., Gullino, Angiogenesis, tumour vascularization, and potential interference with tumour growth. in Biological Responses in Cancer, pp. 178-184 (Mihich (ed.) 1985)). Similarly, Tumor angiogenesis factor (TAF) is released at a higher level in tumour cells than their normal counterparts. See, e.g., Folkman, Angiogenesis and Cancer, Sem Cancer Biol. (1992)). Various techniques which measure the release of these factors are described in Freshney (1994), supra. Also, see, Unkless et al., J. Biol. Chem. 249:4295-4305 (1974); Strickland & Beers, J. Biol. Chem. 251:5694-5702 (1976); Whur et al., Br. J. Cancer 42:305 312 (1980); Gullino, Angiogenesis, tumour vascularization, and potential interference with tumour growth. in Biological Responses in Cancer, pp. 178-184 (Mihich (ed.) 1985); Freshney Anticancer Res. 5:111-130 (1985).

## 25 Invasiveness into Matrigel

The degree of invasiveness into Matrigel-or some other extracellular matrix constituent are used as an assay to identify compounds that modulate ovarian cancer-associated sequences. Tumor cells exhibit a good correlation between malignancy and invasiveness of cells into Matrigel or some other extracellular matrix constituent. In this assay, tumourigenic cells are typically used as host cells. Expression of a tumour suppressor gene in these host cells would decrease invasiveness of the host cells.

Techniques described in Freshney (1994), supra, are used. Briefly, the level of invasion of host cells are measured by using filters coated with Matrigel or some other extracellular matrix constituent. Penetration into the gel, or through to the distal side of the filter, is rated as invasiveness, and rated histologically by number of cells and

111

distance moved, or by prelabeling the cells with 125 1 and counting the radioactivity on the distal side of the filter or bottom of the dish. See, e.g., Freshney (1984), supra.

## Tumor growth in vivo

15

20

WO 2004/022778

5 Effects of ovarian cancer-associated sequences on cell growth are tested in transgenic or immune-suppressed mice. Knock-out transgenic mice are made, in which the ovarian cancer gene is disrupted or in which a ovarian cancer gene is inserted. Knock- out transgenic mice are made by insertion of a marker gene or other heterologous gene into the endogenous ovarian cancer gene site in the mouse genome via homologous recombination. Such mice can also be made by substituting the endogenous ovarian cancer gene with a mutated version of the ovarian cancer gene, or by mutating the endogenous ovarian cancer gene, e.g., by exposure to carcinogens.

A DNA construct is introduced into the nuclei of embryonic stem cells. Cells containing the newly engineered genetic lesion are injected into a host mouse embryo, which is re-implanted into a recipient female. Some of these embryos develop into chimeric mice that possess germ cells partially derived from the mutant cell line. Therefore, by breeding the chimeric mice it is possible to obtain a new line of mice containing the introduced genetic lesion (see, e.g., Capecchi et al., Science 244:1288 (1989)). Chimeric targeted mice are derived according to Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory (1988) and Teratocarcinomas and Embryonic Stem Cells: A Practical Approach, Robertson, ed., IRL Press, Washington, D.C., (1987).

Alternatively, various immune-suppressed or immune-deficient host animals are used. For example, genetically athymic "nude" mouse (see, e.g., Giovanella et al., J. Natl. Cancer Inst. 52:921 (1974)), a SCID mouse, a thymectomized mouse, or an irradiated mouse (see, e.g., Bradley et al., Br. J. Cancer 38:263 (1978); Selby et al., Br. J. Cancer 41:52 (1980)) are used as a host. Transplantable tumour cells (typically about 10<sup>6</sup> cells) injected into isogenic hosts will produce invasive tumours in a high proportions of cases, while normal cells of similar origin will not. In hosts which developed invasive tumours, cells expressing a ovarian cancer-associated sequences are injected subcutaneously. After a suitable length of time, preferably 4 to 8 weeks, tumour growth is measured (e.g. by volume or by its two largest dimensions) and compared to the control. Tumours that have a statistically significant reduction (using, e.g. Student's T test) are said to have inhibited growth.

#### Administration

therapeutic reagents of the invention are administered to patients, therapeutically. Typically, such proteins/polynucleotides and substances may preferably be combined with various components to produce compositions of the invention. Preferably the compositions are combined with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition (which are for human or animal use). Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition of the invention are administered by direct injection. The composition are formulated for parenteral, intramuscular, intravenous, subcutaneous, intraocular, oral, vaginal or transdermal administration. Typically, each protein are administered at a dose of from 0.01 to 30 mg/kg body weight, preferably from 0.1 to 10 mg/kg, more preferably from 0.1 to 1 mg/kg body weight.

Polynucleotides/vectors encoding polypeptide components for use in modulating the activity of the ovarian cancer-associated proteins/polynucleotides are administered directly as a naked nucleic acid construct. When the polynucleotides/vectors are administered as a naked nucleic acid, the amount of nucleic acid administered may typically be in the range of from 1 µg to 10 mg, preferably from 100 µg to 1 mg.

20

25

30

35

10

Uptake of naked nucleic acid constructs by mammalian cells is enhanced by several known transfection techniques for example those including the use of transfection agents. Example of these agents include cationic agents (for example calcium phosphate and DEAE-dextran) and lipofectants (for example lipofectam<sup>TM</sup> and transfectam<sup>TM</sup>). Typically, nucleic acid constructs are mixed with the transfection agent to produce a composition.

Preferably the polynucleotide or vector of the invention is combined with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition are formulated for parenteral, intramuscular, intravenous, subcutaneous, oral, intraocular or transdermal administration.

The pharmaceutical compositions are administered in a range of unit dosage forms depending on the method of administration. For example, unit dosage forms suitable for oral administration include, powder, tablets, pills, capsules and lozenges. Orally

113

administered dosage forms will typically be formulated to protect the active ingredient from digestion and may therefore be complexed with appropriate carrier molecules and/or packaged in an appropriately resistant carrier. Suitable carrier molecules and packaging materials/barrier materials are known in the art.

5

10

15

The compositions of the invention are administered for therapeutic or prophylatic treatments. In therapeutic applications, compositions are administered to a patient suffering from a disease (e.g. ovarian cancer) in an amount sufficient to cure or at least partially ameliorate the disease and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose". An amount of the composition that is capable of preventing or slowing the development of cancer in a patient is referred to as a "prophylactically effective dose".

The routes of administration and dosages described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and dosage for any particular patient and condition.

The present invention is further described with reference to the accompanying drawings and the following non-limiting examples.

20

114

#### **EXAMPLE 1**

# Gene expression profiling to identify differentially-expressed genes in ovarian cancer

#### 1. Tissue Bank and Database

Tissue was collected from patients undergoing treatment at the GCC, we have established an Ovarian Cancer Tissue Bank and Clinical Database that currently holds data on over 400 cases treated at the GCC between 1986 and 2002. Tissue (currently 149 fresh/frozen and 292 archival fixed paraffin-embedded samples) was acquired from patients undergoing cytoreductive surgery and does not interfere with the collection of tissue for the normal processing of diagnostic specimens. Patient consent, included in all our studies, was collected prior to surgery. Tissue specimens and their associated pathology reports were coded in order to maintain patient confidentiality. Uncoded data was electronically and/or physically locked with restricted access by appropriate senior investigators only. Clinical (diagnosis, treatment, residual disease) and pathological data (tumour grade, stage) were collected and updated (disease recurrence, patient survival) at regular intervals. This study has ethical approval from the South Eastern Sydney Area Health Service Research Ethics Committee, Australia. Clinical data and tissue collection are ongoing.

#### 2. Genetic profiling of ovarian cancers

In order to identify those genes differentially regulated in epithelial ovarian cancer 51 ovarian cancer tumor samples were manually dissected from biological samples derived from subjects undergoing cytoreductive surgery. These samples comprised 8 endometrioid tumors, 4 mucinous tumors and 31 serous epithelial ovarian tumors, 12 corresponding omental deposits and 8 borderline (low-malignant potential) tumors.

25

30

5

10

15

20

RNA was isolated from the tumor samples in addition to 4 normal ovary samples using Trizol reagent (Life Technologies, Rockville, MD, USA) essentially according to manufacturer's instructions. RNA was then reverse transcribed using an oligo(dT) anchored oligonucleotide that additionally comprised a T7 promoter sequence. Isolated cDNA was then transcribed *in vitro* using the T7 MEGAscript kit (Ambion, Austin, TX, USA) according to manufacturer's instructions. Transcription was performed with biotinylated nucleotides (Bio-11-CTP and Bio-16-UTP) to enable detection of the transcribed cRNA.

Levels of gene expression in the cancer samples was then determined by analysing the transcribed cDNA samples using customized Affymetrix GeneChip® microarrays that

115

comprise 59,618 oligonucleotide probe sets. These probe sets facilitate analysis of 46,000 gene clusters, representing over 90% of the predicted expressed human genome.

Data were normalized, and changes in gene expression detected using a ranked penalized t-statistic with p-values adjusted for multiple testing using the Holm procedure. Analysis was performed using the LIMMA package (available from Bioconductor, Biostatistics Unit of the Dana Farber Cancer Institute at the Harvard Medical School/Harvard School of Public Health).

Gene expression in 186 samples representing 52 different tissues of the body was also determined using the previously described methods to facilitate the identification of changes in gene expression that are specific for ovarian cancer.

Using this method 284 up-regulated transcripts and 186 down-regulated transcripts were identified.

In order to determine the efficacy of such a method of analysis for determining gene expression changes associated with ovarian cancer, those genes identified were compared to results of published expression profile studies. Using this method, 71 genes were identified in the present study that had been previously identified, including, for example, genes known to be over-expressed in ovarian cancer, such as, for example MUC1 and E-cadherin.

The ovarian cancer-associated genes and proteins set forth in Table 1 include sequences that are up-regulated or down-regulated in ovarian cancer subjects, including subjects suffering specifically from serous, encodmetrioid, mucinous or clear cell ovarian cancer, or non-invasive (borderline) ovarian cancers of any phenotype, and subjects that suffered from recurrences of ovarian cancer in the medium term, or died within the medium term.

30

5

15

20

25

Data presented in Table 2 indicate those genes that are expressed at significantly higher levels or significantly reduced levels in patients suffering from serous cancer relative tot he level of expression of the same genes in a normal or healthy subject.

116

## **EXAMPLE 2**

Validation of gene expression profiling results using tissue microarrays

Each of the transcripts identified as being differentially-expressed specifically in ovarian cancer was then further analysed using *in situ* hybridization or immunohistochemical staining of tissue microarrays constructed from a large cohort of primary ovarian tumor tissue. Such analysis confirms upregulation, down-regulation or total loss of expression of the transcripts identified in the microarray analysis of tumor samples.

Furthermore, as each of the samples in the tissue microarray have been clinicopathologically characterized (for example to identify cancer grade and/or disease stage) and the subjects from whom the tumors were isolated continuously monitored (to detect for example, death or relapse of cancer), changes with gene expression were also analysed for correlation with such parameters in order to determine predictive changes in gene expression.

15

20

25

10

5

The relative intensity and percentage of cells staining was determined and evaluated for associations with clinical stage and grade of disease and disease relapse using the Kaplan Meier method and log-rank test, and by univariate and bivariate analyses in a Cox proportional hazards model for gene expression and other clinical and pathologic predictors of outcome to determine the potential independent prognostic value of the markers being assessed.

Immunohistochemical analysis has been performed on several genes identified in gene profiling analysis of ovarian cancer samples. For example, SOX17, Ep-CAM and claudin 3 were shown by gene profiling analysis to be specifically up-regulated in ovarian cancer compared to normal ovaries (Figure 1 and Figure 2). Using immunohistochemical analysis, it was determined that SOX17, Ep-CAM and claudin 3 are upregulated in serous cancer, mucinous cancer, endometroid cancer and clear cell ovarian cancer.

Furthermore, immunohistochemical analysis has been used to analyse the expression of several other genes that are specifically upregulated in mucinous ovarian cancer. In particular the expression of LI-cadherin (cadherin 17), meprin alpha and Galectin 4 as detected using immunohistochemistry is shown in Figure 3. There was a significant increase in protein detected in the mucinous ovarian cancer samples compared to the normal ovary sample and serous ovarian cancer sample.

Immunohistochemical analysis was also performed to analyse the expression of three genes that are known to be upregulated in ovarian cancer (CA125, MUC-1 and E-cadherin) (Figures 1 and 2).

5

10

15

20

25

30

35

## **EXAMPLE 3**

### Identification of prognostic markers of ovarian cancer

Using a classical survival analysis to mine expression profiling data several genes that are associated with poor patient outcome (ie death or cancer relapse) have been identified (Tables 2 and 3). Such genes have clinical utility as prognostic indicators of disease.

Using detailed clinicopathological and postoperative data on all of the 51 patients included in our transcriptional profiling studies, including details of biochemical (eg. rising serum CA-125) and/or clinical recurrence of disease and overall survival, expression profiles were correlates with clinical parameters.

A preliminary survival analysis was performed on the 33 serous cancers within this cohort. The median follow-up time for these patients was 25.5 months from the date of primary laparotomy to the date of last follow-up or the date of death, and 21 of these patients (66%) were deceased from causes related to their malignancy.

Preliminary analysis of the expression profiles of these tumors identified several potential gene clusters that were associated with an increased risk of biochemical and clinical recurrence and overall survival, including the *EDD* gene (SEQ ID NO: 63). Exemplary prognostic markers for detecting ovarian cancer are shown in Tables 1 and 3. Preferred markers are indicated in Table 3.

Using immunohistochemical analysis two genes have been confirmed to be upregulated in serous ovarian cancer. In particular, sFRP4, a negative signalling protein of the Wnt pathway, and SOCS3, a negative signaller of IL-6 induced signalling are specifically upregulated in serous ovarian cancer when compared to normal ovarian tissue (Figure 4A).

Furthermore, using clinical patient data and correlating this information with gene expression levels using a Cox proportional hazards model, it has been shown that high

118

expression of sFRP4 correlates with a poor outcome in patients (n=127) with serous ovarian cancer (p=0.0056) (Figure 4B).

## **EXAMPLE 4**

Validation of gene expression profiling results using quantitative RT-PCR Candidate diagnostic genes are screened by quantitative RT-PCR against ovarian cancer cell lines to both validate the transcript profiling data (ie check their up- or down-regulation). Candidate diagnostic genes are screened using mRNA isolated from a panel of 9 ovarian tumour cell lines, (A2780, SKOV3, OVCAR-3, IGROV-1, CAOV3, OV-90, SW626, TOV-21G and TOV-112D), in addition to several other tumour cell lines including lines derived from breast, prostate and colorectal tumours, and immortalised (non-transformed) human ovarian surface epithelial cells and a primary normal breast epithelial cell line (184).

Total RNA is isolated from the normal and tumour cell lines, reverse transcribed into cDNA and used as template in a quantitative PCR using a LightCycler system (Roche Diagnostics). The relative amount of each gene product is determined by comparison to a standard housekeeping gene (GAPDH).

EXAMPLE 5

5

10

20

25

30

35

Identification of Novel Genes for Diagnosis of Ovarian Cancer

We identified candidate genes with diagnostic potential from our list of aberrantly regulated genes by applying the following selection procedure: genes with a good transcript profile and low p-value (ie highly significantly up- or down-regulated in ovarian cancer, as determined in Example 1); and mapping to areas of the genome that have been shown to be amplified or lost in ovarian cancer. Accordingly, it is likely that these genes are involved in the development and progression of ovarian cancer (ie putative oncogenes and tumour suppressor genes). Additional parameters for analysis included known or putative function in oncogenesis (eg signal transduction, regulation of cellular proliferation, apoptosis etc); and association with other forms of other tumours. Genes identified in this analysis are shown in Table 3.

One method for the diagnosis of cancer comprises detecting modified DNA shed by the developing tumour into the blood stream. This can include the detection of mutations in both oncogenes and tumour suppressor genes involved in the development and progression of ovarian cancer. Furthermore, it has been recently shown that aberrant

119

methylation of tumour suppressor genes, specifically hypermethylation of their gene promoters, frequently accompanies gene silencing in cancers, and indeed in some cases appears to be the predominant mechanism of gene silencing.

Combined with the knowledge of tumour nucleic acids circulating in the blood that reflect the biological characteristics of a tumour, the detection of methylation-specific tumour suppressor gene signatures for any given tumour type has promise as a specific and sensitive molecular test for detecting and monitoring cancer. Aberrant methylation is a frequent epigenetic event in epithelial ovarian cancer and many candidate tumour suppressor genes of epithelial ovarian cancer have been shown to be hypermethylated in epithelial ovarian cancer, such as, for example BRCA1.

In particular, expression of the candidate tumor suppressor gene MCC, has been shown to be down-regulated in epithelial ovarian cancer compared to normal ovarian tissue.

MCC appears to be involved in critical cell growth regulatory processes and maps to a chromosomal region hypothesised as containing a tumor suppressor gene in ovarian cancer. Furthermore, we have identified a CpG island within the predicted promoter sequence of the MCC gene, a critical feature of genes that are subject to gene silencing by hypermethylation and a known characteristic of tumor suppressor genes. Taken together these data strongly implicate MCC as a candidate tumor suppressor gene involved in epithelial ovarian cancer.

	P value	<b>.</b>	0	0	0	0	0	<b>o</b>	0	0
	bjects suffering from ovarian cancer Putative Function	Lymphocyte antigen, plasma membrane, tumor antigen. Member of the CA733 family, C archoma-associated antigen expressed on most normal epithelial cells and gastrointestinal cardinomas and functions as a homotypic calclum-independent cell adhesion molecule. The partition is holytected or homotypic and the collection of homotypic and the collection.	amygan is being used as a talyet for immunicuted apy used into the human carcinomas. Home saplens cDNA FL/10920 fis, done OVARC1000384-resourcerer.	contains 3 RNA recognition motifs	weakly similar to a drosophila transcription factor	Mai2 T-cell differentiation protein; found thru interaction with TPD52 which is overexpressed in breast cancer, 4 TM are involved in vesicle transport	Tumor suppressor. Ca2+-dependent glycoprotein, mediates cell-cell interactions in epithelial cells. Mutations correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function thought to contribute to progression in cancer by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the protein in consider the contribute of the contribute	Sychastics contains required for internatization.  Early endosome, membrane fraction, non-selective vesicle docking, non-selective vesicle transport, protein complex assembly, synaptic vesicle. Member of a family involved in docking or fusion of synaptic vesicles. Associated with the perhudear vesicular structures of the party and contactines.	Cell shape and cell size control, cell-cell adherens junction, epidermal differentiation, intermediate filament, structural constituent of cytoskeleton. Acts as a site of attachment for intermediate filaments in desmosomes (intercellular junction in vertebrate epitheliala cells). Compound heterozgosity for non-sense and miscone on definer independent of the programment of the compound heterozgosity for non-sense and miscone on definer independent.	ocias indexacts of controlled statements of the statement
Table 1	Genes having modified expression in subjects suffering from ovarian cancer Gene symbol and title	Ep-CAM; TACSTD1, tumor-associated calcium signal transducer 1; epithelial gylcoprotein	HSPC195, hypothetical protein HSPC195	FLJ20171, hypothetical protein FLJ20171	FLJ13782, Hypothetical protein FLJ13782	MALZ	CDH1, cadherin 1, type 1, E-cadherin (epithelial)	VAMP8, vesicle-associated membrane protein 8 (endobrevin)	DSP, desmoplakin (DPI, DPII)	CD24: CD24 antigen (small cell lung carcinoma cluster 4 antigen)
	UniGene Mapping	Hs.692:235	Hs.15093:210; Hs.290304:1	Hs.24743:94	Hs.257924:13	Hs.76550:164	Hs.194657:233	Hs.172684:89	Hs.349499	Hs.286124:357; Hs.375108
	Accession number	a. upregulated genes NM_002354	BC006428	NM_017697	AW419198	AW630088	NM_004360	NM_003761	NM_004415	NM_013230

P value	inhibitor. Member 0 Hepatocyte growth Tic for HGF activator solytic activation of	0.0001		te niaments; type ii 0.0002 Disruption of ssion in vivo could attc disorders	epidermal growth 0.0002 1, protein amino family of recentor	nannmary tumor Ilating HGL- irmal growth factor for interaction with.	0.0002	osphate to uridine 0.0003	s. An allele of this munity to	is. An allele of this munity to incrotubule 0.0004 mization and pressed in cells of and cell oules, and may ated mouse protein equired for
Putative Function	Extracellular, membrane fraction, serine protease inhibitor. Member of the Kunitz family of serine protease inhibitors. Hepatocyte growth factor activator inhibitor is a potent inhibitor specific for HGF activator and is thought to be involved in regulation of proteolytic activation of	HGF in injured tissues. Function unknown	Function unknown	Cell su double, Cytoskeletal. May form intermediate tilaments; type in keratin, member of a family of structural proteins. Disruption of mechanisms that normally regulate keratin expression in vivo could be related to inflammatory and neoplastic pancreatic disorders (Casanova 1999).	Transmembrane receptor protein tyrosine kinase, epidermal growth factor receptor, integral plasma membrane protein, protein amino acid phosphorylation. Member of the ERBB gene family of recentor	tyrosine kinases, elevated levels in certain human mammary tumor cell lines. A receptor for heregulin, capable of mediating HGL-stimulated tyrosine phosphorylation of itself. Epidermal growth factor contains both positive and negative determinants for interaction with enth-2/1Erlbl-3 helenclines (Sturders 2002)	Function unknown	Catalyzes the phosphorylation of uridine monophosphate to uridine diphosphate. First step in production of pyrimidine nucleoside	triphosphates required for RNA and DNA synthesis. An allele of this gene may play a role in mediating nonhumoral immunity to	triphosphates required for RNA and DNA synthesis. An allele of this gene may play a role in mediating nonhumoral immunity to Hemophilus influerizae type B. Establishment and/or maintenance of cell polarity, microtubule associated protein, microtubule cytoskeleton organization and plogenesis, structural molecule. Predominantly expressed in cells of epithelial origin. Involved in microtubule dynamics and cell polarization and differentiation. Stabilizes microtubules, and may modulate microtubule functions. Studies of the related mouse protein suggest an essential role in microtubule function required for
Gene symbol and title	SPINT1, serine protease inhibitor, Kunitz type 1. Hepatocyte growth factor activator inhibitor.	FLJ90586, hypothetical protein	KRT8 keralin 8		ERBB3, v-erb-b2 enythroblastic leukemia viral oncogene homolog 3 (avian)		ESTs, Weakly similar to CYL1_HUMAN CYLICIN I [H.sapiens]	UMPK, uridine monophosphate kinase		MAP7, microtubule-associated protein 7
UniGene Mapping	Hs.233950:84,Hs.182 265:2,Hs.7771:1	Hs.17558:16 He 24543:36	Hs.242463:1		Hs.199067:46	•	Hs.294142:167	Hs.75939:33,Hs.1708 64:1		Hs.146388:47,Hs.113 919:3
Accession number	NM_003710	NM_153345 NM_015238	AI282759		Al393742			NM_012474; W70171		AA165082

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
NM_004433	Hs.166098:170	ELF3, E74-like factor 3 (ets domain transcription factor, epithelial-specific)	Embryogenesis and morphogenesis, transcription co-activator, transcription factor, transcription from Pol II promoter. ETS domain transcriptional activator, activates expression of epithelial cell	0.0004
AW247252	Hs.75514:181	NP, nucleoside phosphorylase	specific genes.  DNA modification, nucleobase nucleoside nucleotide and nucleic acid metabolism, purine-nucleoside phosphorylase. Enzyme purine nucleoside phosphorylase together with adenosine deaminase (ADA) serves a key role in purine catabolism, referred to as the salvage pathway. Mutations in either enzyme result in a severe	0.0004
NM_015925	Hs.361379, Hs.95697:59,Hs.9364 9:1	LISCH7, Liver-spedfic bHLH-Zip transcription factor	combined immunodefidency (SCID). LISCH protein	0.0004
NM_022454	Hs.97984:22	SOX17, SRY (sex determining region Y)-box 17	Likely ortholog of mouse SRY-box containing gene 17; allas SOX17	0.0005
A1124756	Hs.5337:191	IDH2, isocitrate dehydrogenase 2 (NADP+), mitochondrial	Carbohydrate metabolism, mitochondrion	0.0006
NM_003084	Hs.313:273,Hs.29789 5:1	SPP1, secreted phosphoprotein 1 (osteopontin, bone staloprotein I, early T-lymphocyte activation 1)	Osteopontin (bone statoprotein); bone and blood vessel extracellular matrix protein involved in calcification and atheroscierosis, increased expression is associated with breast tumor metastasis (Urquidi 2002). Role in HCC, especially in cancer-stromal interactions (Gotoh 2002). Association between levels of a biomarker, osteopontin, and protein cancer strong in the principal process.	0.0006
BE382756	Hs.169902:319,Hs.27 5406:1	SLC2A1, Solute carrier family 2 (facilitated glucose transporter), member 1	Glucose transporter, membrane fraction, SLC241/GLUT1. Glucose transporter, membrane fraction, SLC241/GLUT1. facilitated glucose transporter, Glucose transporter an integral membrane glycoprotein that is involved in transporting glucose into most cells. 12 TMs. Role in transport of glucose across the bloodbrain barrier. Consistent marker of ovarian epithelial malignancy (Kalir 2002). Marker for discriminating hepatocellular carcinoma from	0.0006
BE512730	Hs. 65114:718, Hs. 279 437:1	KRT18, keratin 18	other carcinomas (Zimmeman 2002). Cell shape and cell size control, embryogenesis and morphogenesis, intermediate filament, structural constituent of cytoskeleton. Component of intermediate filaments; type I epidemal keratin, strongly similar to murine Endo B. Expressed in single layer epithelial tissues of the body. Mutations linked to cryptogenic cirrhosis.	0.0006

P value	0.0006	0.0007	0.0009	0.0009	0.0009	0.001	0.0011	0.0011	0.0011	0.0012	0.0012
Putative Function	Plasma membrane, integral plasma membrane protein. Member of the transmembrane 4 superfamily (TM4SF); may mediate platelet activation and aggregation. Cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily	Contains four RNA recognition motifs (RRM, RBD, or RNP)	Endopeptidase inhibitor, extracellular space, proteolysis and peptidolysis, spermatogenesis. Epididymis-specific secreted protein; may have a role in sperm maturation; arelong to a family of extracellular proteinase inhibitors. Expressed in pulmonary epithelial cells, and also expressed in some ovarian cancers.	Amiloride-sensitive sodium channal, excretion, integral plasma membrane protein, membrane fraction, sodium transport. Alpha subunit of the amiloride-sensitive epithelial sodium channel; functions in nonvoltage-gated channel	Histogenesis and organogenesis, embryogenesis and morphogenesis, thyroid-stimulating hormone receptor, transcription factor. Member of the paired domain family of nuclear transcription factors; are involved in the ribosome assembly, required for normal thyroid development. PAX genes play critical roles during fetal development and cancer growth.	Function unknown	Function unknown	Antimicrobial humoral response (sensu invertebrata), cell-cell signaling, chemokine chemotaxis. Cytokine A16; lymphocyte and monocyte chemoattractant.	KiAA0175 gene product; serine/threonine protein kinase domain	amino acid transporter A1 (ATA1), likely ortholog of mouse N-system amino acid transporter protein NAT2.	cell adhesion, cell-cell signalling, G-protein linked receptor, integral plasma membrane protein, G-protein linked receptor protein signalling pathway. Member of the G protein-coupled receptor family; similar to secretin and calcitonin receptors. 7 transmembrane domains, a mucin-like domain and cysteine box in the N-terminal region. Expressed in range of tissues, highest levels in thyroid, selectively within the monolayer of cuboidal epithelial cells of the smaller, more actively secreting follicles of human thyroid. Differentially expressed in melanoma cell lines with different
Gene symbol and title	CD9: CD9 antigen (p24)	FLJ20273, RNA-binding protein	WFDC2, WAP four-disuffide core domain 2	SCNN1A, sodium channel, nonvoltage-gated 1 alpha	PAX8, paired box gene 8	ESTs	ESTs	SCYA16, small Inducible cytokine subfamily A (Cys-Cys), member 16.	MELK, likely ortholog of maternal embryonic leucine zipper kinase.	SLC38A1, solute carrier family 38, member 1	GPR58, G protein-coupled receptor 56
UniGene Mapping	Hs.1244:227,Hs.2305 59:1,Hs.242020:1	Hs.95549:147,Hs.229 556:1	Hs.2719:108,Hs.5445 1:1	Hs.438580	Hs.73149:72,Hs.2130 08:1	Hs.120912:12	Hs.7956:28	Hs.10458:10	Hs.184339:27	Hs.18272:81	Hs.6527:201
Accession number	NM_001769	AI791905; NM_019027	NM_006103	U81961	X69698; NM_013952	A1027643	AA173992	AB018249	NM_014791	NM_030674	NM_005682

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
			metastatic potential (Zendman et al 1999).	
AI669760	Hs.188881:6,Hs.1993 54:1	ESTs	dbEST Library Tissue Type restricted to prostate	0.0013
NM_001730	Hs.84728:127	KLF5, Kruppel-like factor 5 (Intestinal)	RNA polymerase II transcription factor, transcription from Pol II promoter. Zinc finger transcriptional activator, localizes to the nucleus and binds the epidermal growth factor response element, hinds GC hoves	0.0014
Al355761	Hs.242463:2	KRT8, keratin 8	Cell structure, Cytoskeletal. May form intermediate filaments; type II keratin, member of a family of structural proteins. Disruption of mechanisms that normally regulate keratin expression in vivo could be related to inflammatory and neoplastic pancreatic disorders (Casanova 1989).	0.0014
BE019020	Hs.85838:171	SLC16A3, solute carrier family 16 (monocarboxylic acid transporters), member 3 (MCT3)	Integral plasma membrane protein, membrane fraction, monocarboxylic acid transport, monocarboxylic acid transporter. Member of monocarboxylate transporter family; may function as a transporter (MCT3).	0.0015
NM_001307	Hs.278562:101	CLDN7, claudin 7	Integral membrane protein, tight junction. Similar to murine Cldn7;	0.0016
NM_002266	Hs.168557:394	KPNA2, karyopherin alpha 2 (RAG cohort 1, Importin alpha 1)	DNA metabolism, G2 phase of mitotic cell cycle. NLS-bearing substrate-nucleus import, cytoplasm, importin alpha-subunit, nuclear focalization sequence binding, nucleoplasm, regulation of DNA recombination, spindle pole body and microtubule cycle (sensu Saccharomyces). Karyopherin alpha 2 (importin alpha 1); subunit of the NLS (nuclear localization signal) receptor. KPNA2 protein interacts with the NLSs of DNA helicase Q1 and SV40 T antigen and are involved in the nuclear transport of proteins. KPNA2 also may	0.0016
AW176120	Hs.9061:77	MGC2477, hypothetical protein MGC2477	function unknown	0.0016
BE265489	Hs.3123:49	LLGL2, lethal giant larvae (Drosophila) homolog 2	Cytoskeleton, structural molecule. May associate with nonmuscle myosin if heavy chain. cDNA source cancer cell lines. 57% ID to m.musculus 1920382A tumor suppressor gene mgl1	0.0016
BE279383	Hs.26557:77	PKP3, plakophilin 3	Cell adhesion, intercellular junction. Desmosomal plaque proteins are members of the 'armadillo-repeat' multigene family and have important functions in cytoskeletonoell membrane interactions.	0.0016

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
J05581; NM_002456	Hs.89603:128,Hs.296 789:1	MUC1, mucin 1, transmembrane	Integral plasma membrane protein. Cell surface much glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoletic cell lineages. Afterations in glycosylation in epithelial cancer cells. Marker for hepatocellular carcinoma. MUC1 metabolic complex conserved in tumor-derived and normal epithelial cells. Expression predictor of surgical outcome in mass-forming intrahepatic cholangiocarcinoma. Tyrosine kinase c-Src constitutes a bridge between cystic fibrosis transmembrane regulator channel	0.0016
AA531276 AW167128	Hs.59509:9 Hs.231934:3	ESTs (unnamed protein product) ESTs; weakly similar to A57717 transcription factor EC2	railure and MUC1 overexpression in cystic fibrosis. Function unknown Function unknown	0.0017
AW368226	Hs.67928:25,Hs.2298 40:1	Ets-related transcription factor, ESX, epithelium- restricted Ets protein ESX-not in Unigene, but found using resourcerer.	Embryogenesis and morphogenesis, transcription co-activator, transcription factor, transcription from Poi II promoter.	0.0021
AK000733	Hs.23900:82	RACGAP1, Rac GTPase activating protein 1	Strongly similar to murine Racgap1 GTPase-activating protein for rac. The plexin-B1/Rac interaction inhibits PAK activation and partitions Common Strong Minding	0.0024
NM_014736 NM_014586	Hs.81892:95 Hs.109437:17	KIAA0101 gene product HUNK, hormonally upregulated neu tumor- associated kinase	ennatices Serina+D rigatio princing function unknown; no significant hits with Superfamily Developmental processes, protein serine/threonine kinase, signal transduction, protein kinase containing SNF1 (fam of	0.0025
Al885516	Hs.95612:31,Hs.2516 88:1	desmocollin type 2a, desmocollin 2, isoform Dsc2b preproprotein; desmosomal glycoprotein II/III; desmocollin-3-not in Unigene, but found using resourcerer.	semeurredning knases) doman; progesterone and estradion regulated. Similar to murine Hunk. Cell adhesion, intercellular junction	0.0027
AW194426 NM_001982	Hs.20726:17 Hs.199067:83,Hs.167 386:1	ESTs ERBB3, HER3 (c-erb-B3), v-erb-b2 erythroblastic leukemia viral oncogene homolog 3 (avian)	Function unknown  Epidermal growth factor receptor, integral plasma membrane protein, protein amino acid phosphoryfation. Member of the ERBB gene family of receptor tyrosine klnases, elevated levels in certain human mammany tumor cell lines. A recentor for herectulin, canable of	0.0027
NM_007019	Hs.93002:85	UBE2C, ubiquitin carrier protein E2-C	mediating HGL-stimulated tyrosine phosphorylation of itself. Ubiquitin-dependent protein degradation, degradation of cyclin, protein modification, positive control of cell proliferation. Subunit of a complex with ubiquitin ligase activity, complex that exhibits cyclinselective ubiquitin ligase activity.	0.0031

	Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
	BE184455	Hs.251754:128,Hs.24 5742:1	SLPI, secretory leukocyte protease inhibitor (antileukoprotelnase)	Plasma protein, proteinase inhibitor. Secreted inhibitor which protects epithelial tissues from serine proteases. Found in various secretions including seminal plasma, cervical mucus, and bronchial secretions, has affinity for trypsin, leukocyte elastase, and cathepsin G. Its inhibitory effect contributes to the immune response by protecting epithelial surfaces from attack by endogenous proteolytic enzymes; the protein is also thought to have broad-spectrum anti-	0.0034
	Y00815; NM_002840	Hs.75216:262,Hs.228 792:1,Hs.245083:1	PTPRF, protein tyrosine phosphatase, receptor type, F	biolic activity.  Cell adhesion, integral plasma membrane protein, transmembrane receptor protein, tyrosine phosphatase signaling pathway. Receptor-type protein tyrosine phosphatase F; interacts with the insulin	0.0035
-	AA706017	Hs.119944:14	ESTs	receptor; nas ig-like and FIN-III repeats in the extracellular domain Function unknown	0.0038
-	AA256641	Hs.236894:24	ESTs, Highly similar to S02392 alpha-2- macroglobulin receptor precursor	Function unknown	0.0041
	AW055308	Hs.31803:15	ESTs, Weakly similar to TRHY_HUMAN TRICHOHYALI [H.sapiens]	Function unknown	0.0043
•	Al301558	Hs.290801:35, Hs.356228	EST	Function unknown	0.0044
•	<b>T18997</b>	Hs.180372:119; Hs.394609	BCL2-like 1, Homo saplens cDNA FLJ20750 fis, clone HEP05174 (hypothetical protein	Function unknown	0.0044
•	AI798863	Hs.87191:8	ESTs	Finction inknown	0000
•	J03258	Hs.2062:146	VDR, vitamin D (1,25- dihydroxyvitamin D3) receptor	DNA binding, signal transduction, vitamin D3 receptor. Zino-finger DNA-binding transcription factor. Genetic polymorphism determines bone mineral density. Start-vitamin D receptor interactions antagonize 1,25-dihydroxyvitamin D transcriptional activity and	0.0049
•	AA151647	Hs.68877:141,Hs.228 686:1	CYBA, cytochrome b-245, alpha polypeptide	ennance start-mediated transcription. Cytochrome b, membrane, mitochondrion, superoxide metabolism. Alpha-subunit of cytochrome b245, primary component of the microbicidal oxidase system of phagocytes. CYBA deficiency is associated with chronic granulomatous disease (CGD).	0.005

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
Al538613	Hs.135657:8	TMPRSS3 Transmembrane protease, serine 3	integral membrane protein, proteobysis and peptidolysis. Contains a serine protease domain, a transmembrane domain, a LDL receptorlike domain, and a scavenger receptor cystelne-rich domain. Serlne proteases are known to be involved in a variety of biological processes, whose malfunction often leads to human diseases and disorders. Expressed in fetal cochlea and many other tissues, and is thought to be involved in the development and maintenance of the inner ear or the contents of the pertlymph and endolymph. Missense mutations in autosomal recessive sensorineural deafness, Identified	0.0051
NM_018000	Hs.79741:18	FLJ10116, hypothetical protein FLJ10116	as a tumor associated gene that is overexpressed in ovarian tumors.  Function unknown	0.0051
NM_144724 AJ278016	Hs.124740:18 Hs.55565:35	hypothetical protein FLJ30532 ANKRD3, ankyrin repeat domain 3	59% identity to human Zinc finger protein 91 ATP binding, protein amino acid phosphorylation, protein binding, protein serine/threonine kinase.	0.0051
NM_013894	Hs.75562:147	DDR1, discoidin domain receptor family, member 1	Cell adhesion, Integral plasma membrane protein, transmembrane receptor, protein tyrosine kinase. Epithelial-specific receptor protein tyrosine kinase; are involved in cell adhesion; has putative discoldin motifs in extracellular domain. DDR1 (CD167a) is a RTK that is widely expressed in normal and transformed epithelial cells and is activated by various tyros of collaren.	0.0055
T09997: NM_001312	Hs.70327:196,Hs.211	CRIP-2, cysteine-rich protein 2	activated by various typics of companies and protein containing 2 Zn-finger LIM domain protein;208-amino acid protein containing 2 LIM domains	0.0055
BE302796	Hs.105097:115	TK1, thymidine kinase 1, soluble	Cytoplasm, thymidine kinase. Generates thymidylate for DNA synthesis. TK1 gene expression together with TS, TP and DPD gene expression may play important roles in influencing the malignant behavior of explain contact Fillward R 2002.	0.006
NM_001067	Hs.156346:184,Hs.27 0810:2	TOP2A, topolsomerase (DNA) II alpha (170kD)	DNA binding, DNA topolsomerase (ATP-hydrolyzing), nucleus. DNA topolsomerase (ATP-hydrolyzing), nucleus. DNA topolsomerase I apha; may relax DNA torsion upon replication or transcription. Involved in processes such as chromosome condensation, chromatid separation, and the relief of torsional stress that occurs during DNA transcription and replication. Catalyzes the transient breaking and rejoining of two strands of duplex DNA. The gene encoding this enzyme functions as the target for several anticancer agents and a variety of mutations in this gene have been associated with the development of drug resistance. Reduced activity of this enzyme may also play a role in ataxia-telangiectasia.	900.0

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
U46455	Hs.252189:148,Hs.24 8217:1	SDC4, syndecan 4 (amphiglycan, ryudocan)	Integral plasma membrane, proteoglycan syndecan. Syndecans are transmembrane heparan sulfate proteoglycans that appear to act as recenting or consenting the proteographs included in integral, and appears to act as	0.0061
M79141 A1955040	Hs.13234:39 Hs.301584:5,Hs.2653 98:3	ESTs ESTs, Moderately similar to hypothetical protein FLJ20378 [Homo sapiens]	Members of the MYC gene family and 4 members of the syndecan gene family are closely situated on 4 different chromosomes. Function unknown	0.0062
NM_005560	Hs.11669:81,Hs.2310 10:1	LAMAS, laminin, alpha 5	Basement lamina, structural molecule. Widely expressed in adult tissues, with highest levels in lung, heart, and kidney. Fifth member of the alpha subfamily of vertebrate laminin chains. Possible	0.0066
BE563085	Hs.833:97	ISG15, interferon-stimulated protein, 15 kDa	basament membrane protein; contains laminin EGF-like domain, two extracellular laminin G domains. Cell-cell signaling, cytoplasm, extracellular space, protein binding. Protein that is induced by interferon.	0.0068
BE278288	Hs.155048:119	LU, Lutheran blood group (Auberger b antigen included)	Blood group antigen, cell adhesion, Integral plasma membrane protein, signal transduction, transmembrane receptor. Lutheran	0.0069
NM_020859	Hs.278628:52	ShrmL, Shroom-related protein (KIAA1481 protein)	blood group glycoprotein; may play role in cell-cell, cell-matrix adhesion, signal transduction; member of the lg superfamily, has integrin-binding motifs, SH3 domains. Amilonde-sensitive sodium dhannel (weakly similar to Mus musculus PD7 domain activity blooms provided.	0.0074
Al262789	Hs.93659:52	ERP70, protein disulfide isomerase related protein (calcium-binding protein, intestinal-related)	Endoplasmic reticulum lumen, protein secretion. Strongly similar to rat Rn.4070 (CABP2); may bind calcium.	0.008
NM_006147	Hs.11801:77	IRF6, interferon regulatory factor 6	Member 6 of the Interferon regulatory factor transcription factor family; has low similarity to IRF4, which is a lymphocytic transcription	0.0082
R61463	Hs.16165:50	LAK-4P, expressed in activated T/LAK iymphocytes	ractor that stimulates B cell proliferation. expressed in activated T/LAK tymphocytes	0.0082
AI878857; NM_016185	Hs.109706;285	HN1, hematological and neurological expressed 1 protein	Strongly similar to murine Hn1	0.0087
AK001783	Hs.73239:37	FLJ10901, hypothetical protein FLJ10901	B link shows some homology to KIAA1294 but no known function	0.009

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AC004770	Hs.4756:99	FEN1, flap structure-specific endonuclease 1	DNA repair enzyme, DNA replication, UV protection, double-strand break repair, double-stranded DNA binding, double-stranded DNA specific exodeoxyribonuclease, endonuclease, fatty acid desaturation, membrane fraction. Removes 5' overhanging flaps in DNA repair and processes the 5' ends of Okazaki fragments in location strand but symbols.	0.0093
AI567421	Hs.273330:137	AGRN: agrin	Agrin is a neuronal aggregating factor that induces the aggregation of acetylcholine receptors and other postsynaptic proteins on muscle fibers and is crucial for the formation of the neuromuscular junction. Acts at the nerve-muscle synapse in the glomerular basal membrane	0.0093
AW161386	Hs.13561:49	MGC4692: hypothetical protein MGC4692	ata on Tyriprovyco. Function unknown	0.0103
M85430	Hs.155191:546	VIL2, villin 2 (ezrin)	Cytoskeletal anchoring, microvilius. Regulates cell adhesion and cortical morphogenesis. The cytoplasmic peripheral membrane protein encoded by this gene functions as a protein-tyrosine kinase substrate in microvilii. As a member of the ERM protein family, this protein serves as an intermediate between the plasma membrane and the actin cytoskeleton. It plays a key role in cell surface structure adhesion prication and order protein.	0.0106
AW250380	Hs.109059:124,Hs.24 756:11	MRPL12, mitochondrial ribosomal protein L12	Protein synthesis, General cellular role, Ribosomal subunit, Mitochondral, RNA-binding protein, Ribosome-associated.	0.0114
AI733848; NM_021220	Hs.71935:13	ZNF339, zinc finger protein 339	Zinc finger protein	0.0115
AF111856; NM_006424	Hs.105039:48	SLC34A2, solute carrier family 34 (sodlum phosphate), member 2	SLC34A2: solute carrier family 34 (sodium phosphate), member 2; contains 8 predicted TMs and a cysteine-rich N-terminal region. Type 2 sodium-dependent phosphate transporter, member of the renal type II ro-transporter family.	0.0121
BE386983; NM_138410	Hs.343214	CKLFSF7; chemokine-like factor super family 7	chemokine-like factor gene superfamily; transmb 4 superfamily	0.0131
AA433988	Hs.98502:8	MUC16, mucin 16, CA125	Mucin 16. Alias CA125 ovarian cancer antigen	0.0137
AW248314	Hs.9622:83	MRPS18A, mitochondrial ribosomal protein S18A	Mitochondrial small ribosomal subunit, protein biosynthesis, structural constituent of ribosomeribosomal mitochondrial protein	0.0149

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AA454501	Hs.43668:65	PTP4A3, protein tyrosine phosphatase type IVA, member 3	Prenylated protein tyrosine phosphatase. PTPs are cell signaling molecules that play regulatory roles in a variety of cellular processes. Strong similarity to murine Ptp4a3 (Mm.4124). Overexpression of this gene in mammalian cells was reported to Inhibit angiotensin-II induced cell calcium mobilization and promote cell growth. PRL3 (PTP4A3) expressed at high levels cancer metastases (Saha et al. 2001), PRL3 gene is important for colorectal cancer metastasis.	0.016
U33446	Hs.75799:116	PRSS8, protease, serine, 8 (prostasin)	Extracellular space, plasma membrane, serine type peptidase. A trypslnogen, member of the trypsin family of serine proteases. Highly expressed in prostate epithelia, one of several proteolytic enzymes found in seminal fluid. Protease-mediated regulation of sodium absorption is a function of human airway epithelia, and prostasin is a likely candidate for this arrivals.	0.0166
X98654	Hs.93837:43	PITPNM, phosphatidylinositol transfer protein, membrane-associated	Brain development, lipid metabolism, membrane fraction, phosphatidylinositol transporter, phototransduction. Catalyzes the transport of phosphatidylinositol between membranes; similar to Drocophila men	0.0167
AI660149	Hs.44865:39,Hs.3008 19:19,Hs.293904:14	LEF1, Lymphold enhancer-binding factor-1.	Very strongly similar to murine Leff; may act as a transcription factor. Expressed in pre-B and T cells. Binds to T-cell receptor-alpha enhancer and confers maximal enhancer activity. A target gene ectopically activated in colon cancer, from selective activation of a promoter for a full-length LEF1 isoform that binds beta-catenin	0.0172
AF098158; NM_012112	Hs.9329:152	C20orf1, chromosome 20 open reading frame 1	ATP binding, GTP binding, cell proliferation, mitosis, nucleus spindle. Proliferation-associated nuclear protein; associates with the spindle note and mitoric squalle nuclear protein.	0.0183
AB014551	Hs.337774 Hs.337774	ARHGEF2, rho/rac guanine nucleotide exchange factor (GEF) 2	Cell shape and cell size control, cell surface receptor linked signal transduction, guanyi-nucleotide exchange factor, microtubule cytoskeleton. Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form complex with G proteins and stimulate Rho-dependent signals. Rho/Rac guannine nucleotide exchange factor (GEP) 2; associates	0.0206
Al278023	Hs.89986:24,Hs.2907 80:1	ESTs	with micromothes, summates GFF officially on hac and Nito Function unknown	0.0208

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
Z95152	Hs.178695:25,Hs.791 07:1	MAPK13, mitogen-activated protein kinase 13	MAP kinase, antimicrobial humoral response (sensu Invertebrata), cell surface receptor, signal transduction, chemotaxis, stress response. MAP kinases act as an integration point for multiple blochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. Are activated by proinfiammatory cytokines and cellular stress. Transcription factor ATF2, and microtubule dynamics regulator stathmin are substrates of this	0.0217
AW840171	Hs.265398:7	ESTs, Moderately similar to hypothetical protein FLJ20378 [Homo sapiens]	Kinase. Function unknown	0.0222
D49441	Hs.155981:53	MSLN, mesothelin	Cell adhesion, cell surface antigen, membrane. Pre-promegakaryocyte potentiating factor. An antibody that reacts with ovarian cancers and mesotheliomas was used to isolate a cell surface antigon named mesothelin. Although the function of mesothelin is unknown, it may play a role in cellular adhesion and is	0.0225
AW797437	Hs.69771:282,Hs.444: 1,Hs.294163:1	EST, CM1-UM0039-030400-173-a09	present on mesothellum, mesothellomas, and ovarian cancers. Function unknown	0.0229
BE396290	Hs.5097;261	SYNGR2, synaptogydn 2	Integral plasma membrane protein, member of a family of transmembrane synaptic vesicle proteins, specialized secretory organelles that store neurotransmitters in nerve terminals, and release them by fusing with the presynaptic plasma membrane	0.0229
Al656168; NM_025080	Hs.7331	ASRGL1; asparaginase like 1	during exocytosis. glycoprotein catabolism	0.02
NM_002145	Hs.2733;25	HOXB2, homeo box B2, Hox2H protein	Circulation, developmental processes, transcription factor. Member of homeodomain family of DNA binding proteins; may regulate gene expression, morphogenesis, and differentiation. Genes of the HOXB (or HOX2) complex are expressed specifically in erythromegakaryocytic cell lines, some are expressed only in	0.024
AW959311	Hs.87019:8; Hs.172012	Hypothetical protein DKFZp434J037	hematopoietic progenitors. probabie serine/threonine protein kinase; KIAA0537	0.0251

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
NM_000269	Hs.118638:166,Hs.27 6104:1,Hs.276127:1,H s.276246:1	NME1, non-metastatic cells 1, protein (NM23A)	Transcription factor and nucleoside diphosphate kinase; has a role in the transcriptional regulation of c-myc expression. Mutations in NME1 have been identified in aggressive neuroblastomas.	0.0257
AA379597	Hs.5198:87,Hs.27719 2:1	HSPC150, HSPC150 protein similar to ublquitin- conjugating enzyme	Similar to ubiquitin conjugating enzyme	0.0259
BE148235	Hs.193063:100	Homo saplens cDNA FLJ14201 fis, clone NT2RP3002955	high homology to ARP-3 actin-like protein	0.0259
Al683243; Al587638	Hs.97258	ESTs	Mod similarity to S29539 ribosomal protein L13a	0.03
AF111713	Hs.286218:64	JAM1, junctional adhesìon molecule	Cell motility, Inflammatory response, intercellular junction. Role in the regulation of tight junction assembly in epithelia. Ligation of JAM is required for reovin-induced activation of NF-kappa-B and	0.0261
BE381635	Hs.75725:450,Hs.274 751:1,Hs.277482:1,Hs .277468:1	TAGLN2, transgelin 2	applyings, room in supplyings from the protein transgetin, which complex assembly protein. Homolog of the protein transgetin, which is one of the earliest markers of differentiated smooth muscle. Function not yet determined. Are an actin-binding protein.	0.0275
D14697	Hs.77393:201,Hs.247 769:1	FDPS, farnesyl diphosphate synthase (farnesyl pyrophosphate synthetase, dimethylallyltranstransferase, geranyltranstransferase)	Farnesyl pyrophosphate synthetase (farnesyl diphosphate synthase); part of the cholesterol synthesis pathway.	0.0278
AW194364	Hs.94814	MGC2865, Hypothetical protein MGC2885	Function unknown.	0.0295
<b>T47364</b>	Hs.278613:145	IFI27, interteron, alpha-inducible protein 27	Integral membrane protein, Isolated from estradiol-treated human breast carcinoma cells, Induced by Interferon-alpha in human cell lines of different origin, expression is independent of the presence of estradial receives the cells	0.03
U17760	Hs.301103:71,Hs.755 17:24,Hs.199068:1	LAMB3, Laminin, beta 3 (nicein (125kD), kalinin (140kD), BM600 (125kD)) (Accn NM_000228)	estation receptor in the cano.  Epidemad differentiation, laminin-6, structural molecule. Member of a family of basement membrane proteins. LAMB3 serves as the beta chain in lamitin-6, Mutations in LAMB3 have been identified as the cause of various trace of parlemanticis buildes.	0.0304
AU076517	Hs.184276:142	SLC9A3R1, solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulatory factor 1	Actin cytoskelebn, protein complex assembly. Regulatory cofactor of the NHE3 (SLC9A3) sodium/hydrogen antiporter; interacts with merlin (NF2) and ERM family members; has two PDZ domains. Structural determinants in interaction of beta 2 adrenergic and platelet-derived growth factor receptors	0.0312

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AW880841	Hs.96908 , Hs.74427:112	PIG11 , p53-inducad protein	Negative control of cell proliferation, stress response. May generate or respond to oxidative stress, may have a role in p53-dependent apoptosis Polyak K, Xia Y, Zweier JL, Kinzler KW, Vogelstein B. A model for p53-induced apoptosis. Nature. 1997 Sep 18:389(6548):300-5.	0.0314
H24185	Hs.92918:91	BM-009, hypothetical protein BM-009	Function unknown	0.0314
BE614410	Hs.23044:51	MGC16386, hypothetical protein, similar to RIKEN cDNA	Function unknown.	0.0326
H16423	Hs.82685:37	CD47: CD47 antigen (Rh-related antigen, integrinassociated signal transducer)	Oncogenesis, plasma membrane, plasma glycoprotein, cell-cell matrix adhesion, integral plasma membrane proteoglycan, Integrin receptor signal signalling pathway. Similar to Rh-antigen; may interact with integrins and have a role in intracellular calclum increase during cell adhesion.	0.0336
AU076611; NM_006636	Hs. 154672:123	MTHFD2, methylene tetrahydrofolate dehydrogenase (NAD+ dependent); methenyltetrahydrofolate cyclohydrolase	Electron fransporter, methenyltetrahydrofolate cyclohydrolase, mitochondrion. encodes a ruclear-encoded mitochondrial bifunctional enzyme with methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate cyclohydrolase activities, may provide formyltetrahydrofolate for formylmethionyl tRNA synthesis; Involved in inflation of mitochondrial protein synthesis.	0.0342
Al859390	Hs.288940:49	TMEM8, five-span transmembrane protein M83; type i transmembrane protein	Integral plasma membrane protein. Type I transmembrane protein; contains five membrane-spanning domains	0.0345
AA159216	Hs.55505:57	FLJ20442, hypothetical protein FLJ20442	Contains a dual specificity protein phosphatase catalytic domain; 34% similar to protein-tyrosine phosphatase	0.0354
AF119665; NM_021129	Hs.184011:156	PP, pyrophosphatase (inorganic)	Inorganic diphosphatase, phosphate metabolism. Catalyzes the hydrolysis of pyrophosphate to inorganic phosphate	0.0358
BE513613; NM_005720	Hs.11538:275	ARPC1B, actin related protein 2/3 complex, subunit 1A (41,kD)	Cell motility, structural constituent of cytoskeleton. Arp2/3 complex, subunit 1A; involved in assembly of the actin cytoskeleton, may have a role in profit sion of lamellinodia.	0.0367
NM_012153	Hs.182339	EHF: ets homologous factor	DNA binding, tumor suppressor, cell proliferation, developmental processes, transcription activating factor. Member of the ESE subfamily of fits transcription factors.	0.0404
AW772298	Hs.21103:40,Hs.2667 84:2,Hs.102950:1	Homo saplens mRNA; cDNA DKFZp564B076 (from clone DKFZp564B076)	Alias coat protein gamma-cop	0.0423
H16646	Hs.118666:66	PP591, hypothetical protein PP591	Hypothetical protein PP591 (Novel Human cDNA clones with function of inhibiting cancer cell growth; unpublished)	0.043

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AA279661	Hs.83753:244,Hs.301 236:3	SNRPB, small nuclear ribonucleoprotein polypeptides B and B1	Spliceosome, mRNA splicing, small nuclear ribonucleoprotein. Ut and U2 snRNP protein; component of snRNP complexes, required units of the spliceosome	0.0446
BE001596	Hs.85266:102	iTGB4, integrin, beta 4	Cell adhesion receptor, integrin, invasive growth, oncogenesis. Beta 4 subunit of integrin; involved in cell-cell and cell-matrix interactions; member of a family of cell-surface proteins. Binding of beta 4 to plectin is essential for the proper formation and function of hemidesmosomes.	0.0453
BE246444	Hs.283685:148,Hs.23	FLJ20398, hypothetical protein FLJ20398	100%/175aa unnamed protein g7020468	0.0453
X54942	Hs.83758.34	CKS2, CDC28 protein kinase 2	Cell proliferation, regulation of CDK activity. Similar to S. pombe p13suc1; binds and regulates CDK-cyclin complexes, expressed in different patterns through the cell cycle in HeLa cells, which reflects	0.0478
AA30559B	Hs.238205:36	PRO2013, hypothetical protein PRO2013	Specialized for it is an action provide. Function unknown	0.0483
AF019228	Hs.8036:84	RAB3D, member RAS oncogene family	RAB small monoment GTPase, hemocyte development. GTP- binding protein; are involved in vesicle transport; member of the RAB family of small GTPases. Allas GOV, that is overexpressed in glicoblastoms multiforme tissue as compared to normal brain tissue. GOV is also highly expressed in recurrent glioma, colon tumor	0.0485
NM_001949	Hs.1189:65,Hs.29693 9:2	E2F3, E2F transcription factor 3	metastatic to brain, breast tumors, prostate tumors, and several tumor cell lines Protein binding, transcription factor, transcription initiation from Pol II promoter. Involved in cell cycle regulation, binds retinoblastoma protein (Rb), E2P family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of cycle and action of tumor suppressor proteins and is also a target of	0.049
AF217513	Hs.279905:73,Hs.283 649:4	ANKT, nucleolar protein ANKT	the transforming proteins of sinear browning wissons. clone HQ0310 PRO0310p1 nucleolar protein ANKT ~ no functional data	0.0504
AW513143 AJ245671	Hs.98367:8 Hs.12844:73	ESTs EGFL0, EGF-like-domaln; multiple 6	Expressed in uterus Cell cycle, oncogenesis, integrin ligand, extracellular space. Member of the epidernal growth factor (EGF) repeat superfamily, contains an EGF-like-domain. Expressed early during development, and its	0.0568
AA084248	Hs.85339:64	GPR39, G protein-coupled receptor 39	expression has been detected in lung and meningioma tumors. G-protein linked receptor, G-protein coupled receptor protein signaling pattway, integral plasma membrane protein.	0.19

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
U62801	Hs.79361:65	KLK6, kallikrein 6 (neurosin, zyme)	Serine type peptidase, pathogenesis. Neurosin (protease M, zyme); a serine protease that cleaves amyloid precursor protein (APP). Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease humanders.	0.0159
D49441	Hs. 155981:53	MSLN, mesothelin	usease Julianels.  Cell adhesion, cell surface antigen, membrane. Pre-pro- megakaryocyte potentiating factor. An antibody that reacts with  ovarian cancers and mesotheliomas was used to isolate a cell  surface antigen named mesothelin. Although the function of  mesothelin is unknown, it may play a role in cellular adhesion and is  present on mesothelium mesothelium and ovarian cancers.	0.147
X51630	Hs.1145:22,Hs.29685 1:1	WT1, Wilms tumor 1	Nucleus, transcription factor, transcription regulation, 4 Zn finger domains. Functions in kidney and gonad proliferation and differentiation. Mutations in this gene are associated with the development of Volims tumors in the kidney or with abnormalities of the negligible of the propriet of the properties of the propriet of the properties of the propriet of the properties of the properti	0.2938
AB018305	Hs.5378:149	SPON1, spondin 1, (f-spondin) extracellular matrix protein	by Sommon and the state of the strongly similar to rat F-spondin (Rn. 7546); may have a role in the growth and guidance of axons.	0.3394
AA433988	Hs.98502:8	MUC16, mucin 16, CA125	Mucin 16. Alias CA125 ovarian cancer antigen	0.6568
NM_006149	Hs.5302:132	LGALS4, lectin, galactoside-binding, soluble, 4 (galectin 4)	Lectin, cytosol, cell adhesion, plasma membrane. Binds to beta galactoside, involved in cell adhesion, cell growth regulation, inflammation, immunomodulation, apoptosis and metastasis; member of a family of lectins. LGALS4 is an S-type lectin that is strongly underexpressed in colorectal cancer.	0.0001
AA315933	Hs.120879:17	Homo sapiens, clone MGC:32871 IMAGE:4733535, mRNA, complete cds	Function unknown	0.0001
U47732	Hs.84072:110	TM4SF3, transmembrane 4 superfamily member 3	Integral plasma membrane protein, lysosome, pathogenesis, protein amino acid glycosylation, signal transducer, tumor antigen. Cell surface glycoprotein defined by the monoclonal antibody CO-029 is a 27- to 34-kD membrane protein expressed in gastific, colon, rectal, and pancreatic carcinomas but not in most normal tissues	0.0028

Accession number	UnlGene Mapping	Gene symbol and title	Putative Function	P value
NM_005588	Hs.179704	MEP1A, meprin A alpha, PABA peptide hydrolase	metalloprotease located apically and secreted by epithelial cells in normal colon; degrades broad range of ECM components in vitro; proposed role in tumour progression by facilitating migration, intravasation and metastasis	0.01
AW503395	Hs.5541:112	ATP2A3, ATPase, Ca++ transporting, ubiquitous	Endoplasmic retrculum, adenosinetriphosphatase, small molecule transport, calcium-transporting ATPase, Integral plasma membrane protein. Sarco/endoplasmic retrculum Ca2+-ATPase; pumps calcium.	0.0154
NM_004063	Hs.89436.50	CDH17, cadherin 17, Ll cadherin (liver-intestine)	Cell adhesion, integral plasma membrane protein, membrane fraction, small molecule transport, transporter. Member of the cadherin family of calclum-dependent glycoproteins; facilitates uptake of peptide-based drugs, may mediate cell-cell interactions. Component of the gastrointestinal tract and pancreatic ducts, intestinal proton-dependent peptide transporter in the first step in oral absorption of many medically important peptide-based drugs.	0.0172
AI073913	Hs.100686:20	LOC155465, antentor gradient protein 3	Oncogenesis	0.0266
A1928445	Hs.92254:80	SYTL2: synaptotagmin-like 2	Synaptotagmin-like protein of the C2 domain-containing family of proteins. Although the specific function of the synaptotagmin-like proteins is unknown, a role in regulation of synaptic vesicle trafficking via their C2 domains has been suggested. Region of weak	0.08
W40460	Hs.144442:5	PLA2G10: phospholipase A2, group X	similarity to murine Uph Extracellular, secreted phospholipase A2. Group X secretory phospholipase_a2; hydrolyzes the phospholipid sn-2 ester bond; member of the phospholipase family	0.1888
AA132961	Hs.212533:4	Homo saplens cDNA: FL/22572 fis, done HSi02313	. Function unknown	0.1965
AF111856	Hs.105039:48	SLC34A2, solute carrier family 34 (sodium phosphate), member 2	SLC34A2: solute carrier family 34 (sodium phosphate), member 2; contains 8 predicted TMs and a cysteine-rich N-terminal region. Type 2 sodium-dependent phosphate transporter, member of the renal type II co-transporter family.	0.5078
AA143654		zo65a02.r1 Stratagene pancreas (#937208) Homo saplens cDNA clone IMAGE:591722 5', mRNA sequence	Function unknown	0.036

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
b. prognostic indicators AA046217 NM_015902	Hs. 105370;2	ESTs EDD:Homo saplens progestin induced protein (DD5), mRNA. VERSION NM_020987.1 GI	Function unknown Soluble fraction, cell proliferation, ublquitin—protein ligase, ubiquitin conjugating enzyme, ublquitin-dependent protein degradation. Member of the HECT family of proteins; may function as an E3 ublquitin-protein ligase. This gene is localized to chromosome 8q22, a locus disrupted in a variety of cancers. This gene potentially has a role in regulation of cell proliferation or differentiation.	0.00
T8382 #(NOCAT)	Hs.97927:20	ESTs NM_001615*:Homo saplens actin, gamma 2, smooth muscle, enteric (ACTG2), mRNA. variant 1, mRNA.	Function unknown Structural protein of muscle. Gamma 2 actin; enteric-type, smooth muscle cell actin.	0.01
AB040888		Homo sapiens mRNA for KIAA1455 protein, partial cds	Function unknown	0.01
AA628980	Hs.192371:3	DSCR8 down syndrome critical region protein DSCR8	Function unknown	0.01
AI623351	Hs.172148:51	ESTs	Function unknown	0.01
AW614420	Hs.204354:383	ARHB ras homolog gene family, member B	RHO small monomeric GTPase, RHO protein signal transduction, peripheral plasma membrane protein. Ras-related GTP binding protein of the rho subfamily, member B; may regulate assembly of actin stress fibers and focal adhesions; very strongly similar to murine Arth.	0.01
AA243499	Hs.104800:23	hypothetical protein FLJ10134	Highly similar to murine p19.5; are a membrane protein	0.01
AF251237	Hs.112208:16	GAGED2 XAGE-1 protein	GAGE genes are expressed in a variety of tumors and in some fetal and reproductive tissues. This gene is strongly expressed in Ewing's sarcoma, alveolar rhabdomyosarcoma and normal testis. The protein encoded by this gene contains a nuclear localization signal and shares a sequence similarity with other GAGE/PAGE proteins. Because of the expression pattern and the sequence similarity, this protein also belongs to a family of CT (cancer-testis) antigens.	
AI970797	Hs.64859:16	ESTs	Function unknown	0.01
AF145713	Hs.61490:51	SCHIP1 schwannomin-Interacting protein 1	Cytoplasm. Associates with the neurofibromatosis type 2 protein schwannomin (NF2); contains a colled-coll domain/Proteome	0.01

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
X78565	. Hs.289114:173,Hs.74 637:1	TNC hexabrachion (tenascin C, cytotactin)	Cell adhesion, extracellular matrix, cell adhesion receptor, ligand bluding or carrier. Hexabrachion (tenascin c), an extracellular matrix glycoprotein; has epidermal growth factor-like repeats	0.01
T97307		gb:ye53h05.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA done IMAGE:121497 3', mRNA sequence.	Function unknown	0.01
BE243845	Hs.75511:418	CTGF connective tissue growth factor	Cell motility, plasma membrane, soluble fraction, response to wounding, extracellular matrix, extracellular space, epidermal differentiation, cell growth and maintenance, insulin-like growth factor binding, insulin-like growth factor receptor binding protein. Connective tissue growth factor, blinds IGF, may have a role in requisition ormal and neonjastic cell emowship.	0.01
AW068302	Hs.182183:214,Hs.32 5474:172,Hs.283080: 7	CALD1 caldesmon 1	Cytoskeleton, actin blinding, calmodulin blinding, tropomyosin binding. Protein of unknown function. Actomyosin regulatory protein, non-muscle form	0.01
AL133561 BE313555	Hs.241428:5 Hs.7252:158	DKFZP434B081 protein RAI17 refinite and Indured 17	Function unknown Function unknown	0.01
X07820	Hs.2258:1	MMP10 matrix metalloproteinase 10 .(MMP10; stromelysin 2)	Zinc binding, extracellular space, extracellular matrix, metalloendopeptidase, proteolysis and peptidolysis. Stromelysin 2; matrix metalloprotease that degrades connective tissue	0.02
Al973016	Hs.15725:77	IERS Immediate early response 5	Function unknown. A related mouse gene may play an important role in mediating the cellular response to mitogenic signals.	0.02
AF084545		Homo saplens versican Vint isoform, mRNA, partial cds	Function unknown	0.02
U41518	Hs.74602:146,Hs.767 :1	AQP1 aquaporin 1 (channel-forming integral protein, 28kD)	Excretion, water transport, water transporter, integral plasma membrane protein. Aquaporin 1 (channel-forming integral protein); member of a family of water-transporters	0.02
Z11894		H.sapiens rearranged mRNA for immunoglobulin kappa chain (VNJ)		0.02
AW138190	Hs.180248:8	ZNF124	DNA binding, C2H2 zino-finger protein 124	0.02
BE086548	Hs.42346:83,Hs.6975 :42	MYOZZ myozenin 2 myozenin 2	calcineurin-binding protein calsardin-1	0.02

Accession number UniGene Mapping	UniGene Mapping	Gene symbol and title	Putative Function	P value
W47196	Hs.166172:50	ARNT aryl hydrocarbon receptor nuclear translocator	Nucleus, transcription factor, transcription co-activator, transcription, DNA-dependent, protein-nucleus import, translocation, aryl hydrocarbon receptor nuclear translocator. Aryl hydrocarbon receptor nuclear translocator, used in receptor furchear translocator, used in receptor furchear translocator, used in receptor furchear translocator.	0.02
Al796870	Hs.54277:76	DXS9928E DNA segment on chromosome X (unique) 9928	Nucleus. Has many charged residues and a possible nuclear localization signal	0.02
X02761	Hs.287820:73,Hs.321 592:1	expressed sequence FN1 fibronectin 1	Cell adhesion, cell motility, cell adhesion, soluble fraction, signal transduction, extracellular matrix, extracellular space. Fibronectin 1; member of family of proteins found in plasma and extracellular matrix.	0.02
AW968613	Hs.79428:166	BNIP3 BCL2/adenovirus E1B 19kD-interacting protein 3	Anti-apoptosis, apoptosis Inhibitor, Bci2-related protein 3; binds antiapoptotic viral E18 19 kDa protein and cellular Bci2 protein	0.05
AW972565	Hs.32399:24	ESTs, Weakly similar to S51797 vasodilator- stimulated phosphoprotein [H.saplens]	Function unknown	0.02
AF045229	Hs.82280:81	RGS10 regulator of G-protein signalling 10	Regulator of G protein signaling (RGS) family members are regulatory molecules that act as GTPase activating proteins (GAPs) for G alpha subunits of heterotimeric G proteins. RGS proteins are able to deactivate G protein subunits of the Gl alpha, Go alpha and Gq alpha subtypes. They drive G proteins into their inactive GDP-bound forms.	0.02
AW953853	Hs.292833:19	PAEP progestagen-associated endometrial protein (placental protein 14, pregnancy-associated endometrial alpha-2-globulin, alpha uterine	Developmental processes. Placental protein 14 (Glycodelin); member of lipocalin superfamily, highly similar to beta-lactoglobulins	0.02
U52426	Hs.74597:75,Hs.1576 15:3	STIM1 STIM1 stromal interaction molecule 1	Integral plasma membrane protein, positive control of cell proliferation. Very strongly similar to murine Stim1; are a transmembrane stromal cell protein	0.02
F06700	Hs.7879:115	IFRD1 Interferon-related developmental regulator 1	Myoblast determination. Strongly similar to rat interferon-related developmental regulator 1; may play a role in muscle differentiation	0.02
A1798863 NA	Hs.87191:8	ESTs C4001170:gi 6863176 gb AAF30402.1 AF109924_ 1 (AF109924) sulfatase 1 precursor [Helix poma	Function unknown	0.03
H52761	Hs.141475:24	Homo sapiens cDNA done IMAGE:178663	Function unknown	0.03

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
BE546947	Hs. 44276:43	HOXC10 homeo box C10	Embryogenesis and morphogenesis, positive control of cell proliferation, RNA polymerase II transcription factor. Homeobox C10, member of the homeobox developmental regulator family; binds with HOXA13 and HOXC13 to the Lamin B2 origin; ortholog of Drosophila Andminal R	0.03
AU076643	Hs.313:257,Hs.32991 0:1	SPP1. secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1)	Ossimirario extracellular matrix, skeletal development. Osteopontin (bone staloprotein); bone and blood vessel extracellular matrix protein involved in calcification and atherosclerosis	0.03
#(NOCAT)		NM_015902*:Homo saplens progestin Induced protein (DD5), mRNA. VERSION NM_020967.1 GI		0.03
U20538	Hs.3280;20	CASP6 caspase 6, apoptosis-related cysteine protease	Induction of apoptosis, cysteine-type peptidase, proteolysis and peptidolysis. Caspase 6; a cysteine (thiol) protease; related to the ICE-subfamily of caspases	0.03
AA581602	Hs.41840:7	ESTs	Function unknown	0.03
AJ245210		gb:Homo sapiens mRNA for immunoglobulin gamma heavy chain variable region, partial, clone 1A-4G21.	Function unknown	0.03
X65965		H.sapiens SOD-2 gene for manganese superoxide dismutase		0.03
AI808770	Hs.30258:9	ESTs	Function unknown	0.03
BE386490	Hs.279663:51	PJR Plrin	Nucleus, transcription co-factor, transcription from Pol II promoter. Putative cofactor of the NFI/CTF1 transcriptional activator	0.03
AW581992	Hs.301434:104,Hs.32 8017:1	KIAA1387 KIAA1387 protein	Function unknown	0.03
U77534		Human clone 1A11 immunoglobulin variable region (VH5-D-JH4) gene, partial cds	Function unknown	. 0.03
AL034417	Hs.11169:194,Hs.109 58:1,Hs.74137:1	Gene 33/Mig-6	Function unknown	0.03
L10343	Hs.112341:96,Hs.196 8:1	Homo saplens elafin precursor, gene, complete cds	Function unknown	0.03
AW518944	Hs.76325:80,Hs.2312 99:1	IGJ immunoglobulin J polypeptide, linker protein for immunoglobulin alpha and mu polypeptides	Linker protein for immunoglobulin alpha and mu polypeptides	0.03

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
W28729	Hs.236510:6	Human retina cDNA randomly primed sublibrary Homo sapiens cDNA, mRNA sequence	Function unknown	0.03
AI640160	Hs.74131:4	ARSE arylsulfatase E (chondrodysplasia punctata 1)	Aryisulfatase, skeletal development. Aryisulfatase E; likely involved in warfarin embryopathy.	0.03
U11862	Hs.75741:62	ABP1 amiloride binding protein 1 (amine oxidase (copper-containing))	Metabolism, peroxisome, amine oxidase, drug binding. Diamine oxidase (D-amino-acid oxidase histaminase, amiloride-binding profeln); deaminates putrescine and histamine	0.03
AW295980	Hs.252741:3	ESTs	Function unknown	0.03
X59135	Hs.158110:4	H.sapiens mRNA for immunoglobulin 0-81VL		0.03
BE466173	Hs.379794	Homo sapiens mRNA; cDNA DKFZp686N0118 (from clone DKFZp686N0118)	Function unknown	0.03
#(NOCAT)		Target Exon		0.03
Al354722	Hs.127216:24	hypothetical protein FLJ13465	Function unknown	0.04
M90464	Hs.169825:45,Hs.408 :1	Human collagen type IV afpha 5 chain (COL4A5) gene, 5' end	Function unknown	0.04
AA829286	Hs.332053:48,Hs.336 462:10	SAA1 serum amyloid A1	Inflammatory response, high-density lipoprotein. Member of the serum amyloid A protein family, member of high density applicantities.	0.04
Al333771	Hs.82204:8,Hs.22836 3:1	ESTs	Function unknown	0.04
BE465867; NM_014992	Hs.197751:66	DAAM1 dishevelled associated activator of morphogenesis 1	The protein encoded by this gene contains FH domains and belongs to a novel FH protein subfamily implicated in cell polarity, thought to function as a scaffolding protein.	0.04
BE616902	Hs.285313:145,Hs.40 55:43	COPEB core promoter element binding protein	A trancriptional activator, capable of activating transcription approximately 4-fold either on homologous or heterologous promoters. The DNA binding and transcriptional activity of this protein, in conjunction with its expression pattern, suggests that this protein may participate in the regulation and/or maintenance of the basal expression of pregnancy-specific glycoprotein gene and possibly other TATA box-less genes.	40.0

P value	0.04	.004	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
Putative Function	Function unknown	Function unknown	Function unknown	Function unknown	Function unknown	Function unknown	Function unknown	Function unknown	DNA repair, DNA recombination, cell cycle control, 1-phosphatidylinositol 3-kinase, inositol/phosphatidylinositol kinase. FKBP-rapamycin associated protein; phosphatidylinositol kinase that may mediate rapamycin inhibition of the cell cycle progression throub G1.	Extracellular, serine-type peptidase. Putative serine protease	Energy pathways, secretory vesicle, cytochrome b5 reductase, secretory vesicle membrane, integral plasma membrane protein. Cytochrome b561; serves as a biological marker for adrenergic secretory vesicles.	Drug resistance, Immune response, cell growth and maintenance. Semaphorin E; member of a protein family involved in neuronal growth cone guidance	Nucleus, mRNA splicing, mRNA processing, pre-mRNA splicing factor, May have a role in pre-mRNA splicing; contains arginine/serine-rich domain and an RRM domain		Nucleus, spiloeosome, mRNA splicing, mRNA processing, pre- mRNA splicing factor. Spliceosome-associated protein 3a, subunit 3; component of the essential heterotrimeric splicing factor SF3a; contains a zinc finger
Gene symbol and title	gb:zw20f11.s1 Soares ovary fumor NbHOT Homo sapiens cDNA done IMAGE:769869 3' similar to gb:M63438 iG KAPPA CHAIN PRECURSOR V-III REGION (HUMAN);, mRNA sequence.	ESTs	CTHRC1 collagen triple helix repeat containing 1	ESTs	Target Exon	KIAA0591 protein	ESTs	MGC5469 hypothetical protein MGC5469	FRAP1 FK506 binding protein 12-rapamydn associated protein 1	KLK10 kallikrein 10 (KLK10) (PRSSL1) (nes1)	CYB561 cytochrome b-561	SEMA3C sema domain, immunoglobulin domain (Ig), short basic domain, secreted, (semaphorin) 3C	SFRS11 splicing factor, arginine/serine-rich 11	ENSP00000231844*:Ecotropic virus integration 1 site protein.	SF3A3 splicing factor 3a, subunit 3, 60kD
UniGene Mapping		Hs.271565:3	Hs.283713:68	Hs.39168:16				Hs.19574:123	Hs.338207:139,Hs.14 6559:1	Hs.69423:46,Hs.2754 64:1	Hs.118625:62	Hs.171921:50	Hs.11482:19		Hs.77897:149
Accession number	AA430373	R27430	BE387335	AW264102	NA	AW952323	AA088177	BE614567	AL079658	NM_002776	BE261944	NM_006379	A1002238	#(NOCAT)	X81789

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
NM_002122	Hs.198253:21	HIA-DQA1 major histocompatibility complex, class II, DQ alpha 1	Pathogenesis, class II major histocompatibility complex antigen. Alpha 1 chain of HLA-DQ1 class II molecule (la antigen); complex binds peptides and presents them to CD4+ T lymphocytes[Proteome	0.00
AB001914		Homo sapiens PACE4 gene, exon 23-25, complete cds	Function unknown	0.04
AA311919	Hs.69851:24	NOLA1 nucleoiar protein family A, member 1 (H/ACA small nucleoiar RNPs)	Involved in various aspects of rRNA processing and modification. Localize to the dense fibrillar components of nucleoil and to coiled (Cajal) bodies in the nucleus.	0.04
Al381750	Hs.283437:122,Hs.10 065:58	HTGN29 protein	Function unknown	0.04
#(NOCAT)		NM_000636*:Homo sapiens superoxide dismutase 2, mitochondrial (SOD2), mRNA. expression) (RFX2), mRNA.	Mitochondrion, oxidative stress response, manganese superoxide dismutase. Manganese superoxide dismutase; intramitochondrial free radical scavenging enzyme; has strong similarity to murine Sod2.	0.04
AA292998	Hs.163900:25	ESTs	Function unknown	0.04
BE439580	Hs.75498:40	SCYA20 small inducible cytokine subfamily A (Cys-Cys), member 20	Chemokine, chemotaxis, immune response, signal transduction, extracellular space, cell-cell signalling, inflammatory response, antimicrobial humoral response. Cytokine A20 (exodus): chemotactic factor for iymphocytes, but not a chemotactic factor for monocytes	0.04
AI677897	Hs.76640:124	RGC32 RGC32 protein	Cytoplasm, cell cycle regulator, regulation of CDK activity. Strongly similar to RGC-32.	0.04
#(NOCAT)		Target Exon	Function unknown	0.04
N72403		Homo saplens cDNA done IMAGE:245132	Function unknown	0.05
BE003054	Hs.1695:46	MMP12 matrix metalloproteinase 12 (macrophage	Zinc binding, cell motility, macrophage elastase, extracellular matrix, proteolysis and peptidolysis. Matrix metalloprotease; degrades elastin	0.05
AL035588	Hs.153203:26,Hs.233 91:1	Human DNA sequence from clone 696P19 on chromosome 6p12.3-21.2. Contains the gene for TFEB, an NPM1 (Nucleophosmin, Numatrn) pseudogene and the MDFI gene for MyoD family inhibitor (myogenic repressor I-MF). Contains ESTs, STSs, GSSs and two putative CpG Islands, complete sequence	Function unknown	0.05
Ai080491	Hs.93270:3	ESTs, Moderately similar to S65657 alpha-1C-adrenergic receptor splice form 2 [H.saplens]	Function unknown	0.05

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AW770994	Hs.30340:125	hypothetical protein KIAA1165	Function unknown	0.05
H24177	Hs.75262:69,Hs.2389 12:1	CTSO cathepsin O	Cysteine-type endopeptidase, proteolysis and peptidolysis. Cathepsin O; cysteine (thiol) protease	0.05
AF146781	Hs.20450:29	BCM-like membrane protein precursor	Function unknown	0.05
NM_001955	Hs.2271:45, Hs.306:1	EDN1 endothelin 1	Circulation, peptide hormone, soluble fraction, signal transduction, extracellular space, cell-cell signalling, blood pressure regulation, positive control of cell proliferation. Preproendothelin 1; precursor of the hormone endothelin 1	0.05
A1680737	Hs.289068:204,Hs.32 6198:1	TCF4 transcription factor 4	Nucleus, RNA polymerase II transcription factor, transcription regulation from Pol II promoter. Transcriptional activator, interacts with (TCF3); contains basic helix-loop-helix domain Proteome	0.05
AI752666	Hs.76669:183	NNMT nicotinamide N-methyltransferase	Nicotinamide N-methyltransferase; catalyzes the N-methylation of nicotinamide and other pyridines, structurally-related drugs and xenobiotics Proteome	0.05
AA505445	Hs.300697:21	IGHG3 immunoglobulin heavy constant gamma 3 (G3m	Constant region of heavy chain of igG3	0.05
BE246849; NM_003955	Hs.345728	Sichard Sichard STAT-Inhibitor 3; suppressor of STAT induced STAT-Inhibitor 3; suppressor of cytokine signalling 3	suppression of IL-6 mediated signalling	0.02
M86849	Hs.323733:62,Hs.300 816:5	GJB2 gap junction protein, beta 2, 26kD (connexin 26)	Hearing, connexon, plasma membrane, connexon channel, cell-cell signalling, small molecule transport. Connexin 26; gap junction promises of in various fissues including cochlea.	0.00
AW863419	Hs.155223:20	STC2 stannlocalcin 2	Person of the cell cell signalling, glycopeptide hormone, nutritional response pathway, cell surface receptor linked signal transduction. Stannlocalcin 2, may regulate metal ion homeostasis and inhibits phosphate innake.	0.00
BE298665	Hs.14846:132	Homo sapiens mRNA; cDNA DKFZp564D016 (from clone	Function unknown	0.00
AK000837	Hs.46624:11	· HSPC043 HSPC043 protein	Function unknown	0.00
BE077546	Hs.31447:27	ESTs, Moderately similar to A46010 X-linked retinopathy protein [H.sapiens]	Function unknown	0.00
T97307		gb:ye53h05.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA done IMAGE:121497 3',	Function unknown	0.00

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
		mRNA sequence.		
R24601	Hs.108300:46	Homo sapiens adenylosuccinate synthetase isozyme (ADSS) mRNA, complete cds	Function unknown	0.00
BE090176	Hs.179902:95	Interim-CDw92 antigen	choline transporter-like protein	0.00
AA393907	Hs.97179:22	ESTs	Function unknown	0.00
W28729	Hs.236510:6	Homo sapiens mRNA; cDNA DKFZp666D074 (from clone DKFZp666D074)	Function unknown	0.00
BE313754	Hs.13350:52	Homo sapiens mRNA; cDNA DKFZp586D0918	Function unknown	0.01
AW673081	Hs.54828:9	ESTs	Function unknown	0.01
AA356694	Hs.94011:42,Hs.7744 :2,Hs.231043:1	HCA4 Hepatocellular cardnoma-associated protein HCA4	Function unknown	0.01
L08239	Hs.5326:11	MG61 Porcupine	amino acid system N transporter 2;	0.01
BE397649	Hs.94109:40	Homo sapiens cDNA FLJ34399 fis, clone HCHON2001359	Function unknown	0.01
NM_012317	Hs.45231:36	LDOC1 Leucine zipper, down-regulated in cancer 1	Nucleus, negative control of cell proliferation. Nuclear protein; contains a leucine zipper-like motif	0.01
NM_000947	Hs.74519:20	PRIM2A primase, polypeptide 2A (58kD)	DNA primase, DNA replication, priming, alpha DNA polymerase; primase complex. Subunit of DNA primase polypeptide 24: part of the DNA polymerase alpha-primase complex.	0.01
AJ250562	Hs.82749:133	Homo saplens partial TM4SF2 gene for tetraspanin protein, exon 1 and joined CDS	Function unknown	0.01
AL040183	Hs.123484:24,Hs.326 906:1	Homo sapiens mRNA; cDNA DKFZp686E1934 (from done DKFZp686E1934)	Function unknown	0.01
BE207573	Hs.83321:32	NMB neuromedin B	Peptide hormone, soluble fraction, signal transduction, cell-cell signalling. Precursor of neuromedin B, a C-terminally amidated pendide hormone: similar to hombesin	0.01
BE564162	Hs.250820:45	FLJ14827 hypothetical protein FLJ14827	Function unknown	0.01

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
BE439580	Hs.75498:40	SCYA20 Small Inducible cytokine subfamily A (Cys-Cys), member 20	Chemokine, chemotaxis, immune response, signal bransduction, extracellular space, cell-cell signalling, inflammatory response, antimicrobial humoral response. Cytokine A20 (exodus); chemotactic factor for lymphoxies, but not a chemotactic factor for monocytes.	0.01
AW067800	Hs.155223:52	STC2 stannlocaldin 2	Peptide hormone, cell-cell signalling, glycopeptide hormone, nutritional response pathway, cell surface receptor linked signal transduction. Stanniccalcin 2; may regulate metal ion homeostasis and inhibits observate untake.	0.01
AA569756	Hs.87803:10	Homo sapiens cDNA FLJ30156 fis, clone BRACE2000487	Function unknown	0.01
AW138190	Hs.180248:8	ZNF124 zlnc finger protein 124 (HZF-16)	DNA binding. C2H2 zino-finger protein 124	0.01
AF126245	Hs.14791:48	ACAD8 acyl-Coenzyme A dehydrogenase family, member 8	Lipid metabolism, acyl-CoA dehydrogenase. Member of the acyl- Coenzyme A dehydrogenase family, alpha,beta-dehydrogenates aryl-CoA esters	0.01
L10343	Hs.112341:96,Hs.196 8:1	Homo sapiens elafin precursor, gene, complete cds	elastase-specific inhibitor in bronchial secretions	0.01
NM_002514	Hs.235935:38	NOV nephroblastoma overexpressed gene	insulin-like growth factor receptor binding protein. Insulin-like growth factor binding protein; may play a role in nephrogenesis	0.01
 AI863735	Hs.186755:3	ESTs	Function unknown	0.01
NM_005397	Hs.16426:160,Hs.248 780:1	PODXL podocalyxin-like	Integral plasma membrane protein. Transmembrane protein similar to rodent podocalyxins	0.01
W26391	Hs.301206:100	KIF3B kinesin family member 38	Plus-end kinesin, microtubule motor, anterograde axon cargo transport, plus-end-directed kinasin ATPase, determination of left-right asymmetry. Similar to murine Kif3b; may have a role in intracellular organelle transport, may act in refer-fight determination in the contraction of minary in the contraction.	0.01
H15474	Hs.132898:156	FADS1 fatty acid desaturase 1	eninayoganesis, are a midousobare associated protein. C-5 sterol desaturase, fatty and desaturation, integral membrane protein. Delta-6 desaturase; catalyzes production of polyenoic fatty printein, controlled and con	0.01
U51188	Hs.173824:106	TDG Thymine-DNA giycosylase	actus such as a racinoting actual DNA binding, base-excision DNA repair, nucleoplasm, damaged DNA binding, base-excision repair, G/T-mismatch-specific trymine-DNA glycosylase. Trymine-DNA glycosylase, excises uracil and trymine from mispairs with	0.01
AA243499	Hs.104800:23	FLJ10134 hypothetical protein FLJ10134	guantonie Highly similar to murine p18.5; are a membrane protein	0.01
AW408807	Hs.34497:46	FLJ22116 hypothetical protein FLJ22116	Function unknown	0.01

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AI738719	Hs.198427:98	HK2 Hexokinase 2	Hexokinase, cell cycle control, glucose catabolism, glucose metabolism, mitochondrial outer membrane. Hexokinase II; converts	0.01
AB040888	Hs.41793:110	Homo saplens mRNA for KIAA1455 protein, partial cds	aldo- and keto-nexose sugars to the nexose-o-phosphate Function unknown	0.01
BE313077	Hs.93135:40,Hs.2283 57:1	Homo sapiens cDNA FLJ39971 fis, clone SPLENZ028066	Function unknown	0.01
AI677897	Hs.76640:124	RGC32 RGC32 protein	Cytoplasm, cell cycle regulator, regulation of CDK activity. Strongly similar to RGC-32	0.01
C14898	Hs.192986:5	ESTs	Function unknown	0.01
AI821730	Hs.116524:7	Homo sapiens cDNA FLJ35800 fis, done TESTI2005933	Function unknown	0.01
AF007393	Hs.177574:111	PRKRIR protein-kinase, interferon-Inducible double stranded RNA dependent inhibitor, repressor of (P58 repressor)	Stress response, protein binding, signal transduction, translational regulation, negative control of cell proliferation. Regulates interferoninduced protein kinase PKR (PRKR) activity by binding and inhibiting	0.01
H65423	Hs.17631:42	DKFZP434E2135 hypothetical protein DKFZp434E2135	ura raka-eguadoi Pooira (raka) Function unknown	0.01
N46243	Hs.110373:26	ESTs, Highly similar to T42626 secreted leucinerich repeat-containing protein SLIT2 - mouse (fragment) IM misserties	Function unknown	0.01
AA095971	Hs.198793:56,Hs.309 674:7	Homo sapiens cDNA: FLJ22463 fls, clone HRC10126	Function unknown	0.01
U20350	Hs.78913:33	CX3CR1 chemokine (C-X3-C) receptor 1	Virulence, chemotaxis, coreceptor, cell adhesion, plasma membrane, chemokine receptor, response to wounding, cellular defense response, integral plasma membrane protein, G-protein linked receptor protein signalling pathway. CX3C chemokine receptor; G protein-coupled receptor, mediates leukocyte migration and adhesion, binds the CX3C chemokine fractalkine and signals	0.01
NM_005756	Hs.184942:18	GPR64 G protein-coupled receptor 64	undugir a perussis oxin sensitive c-protein Spermatogenesis, G-protein linked receptor, integral plasma membrane protein, G-protein linked receptor protein signalling	0.01
D19589	Hs.13453:87	FLJ14753 hypothetical protein FLJ14753	patrway. Member of the G protein-coupled receptor family Function unknown	0.02
AW957446	Hs.301711:74	ESTs	Function unknown	0.02

 $\mathcal{B}_{i}$  , the probability of the state o

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AW294647	Hs.233634:40	C20orf39 chromosome 20 open reading frame 39	Function unknown	0.02
BE159718	Hs.85335:46	Homo sapiens, done IMAGE:4513159, mRNA	Function unknown	0.02
A1888490	Hs.56902:22	EDG3 endothelial differentiation, sphingolipid G- protein-coupled receptor, 3	Lipid binding, plasma membrane, Inflammatory response, G-protein linked receptor, embryogenesis and morphogenesis, integral plasma membrane protein, positive control of cell proliferation, cytostolic calcium ion concentration elevation, G-protein linked receptor protein signalling pattway. Lysosphingolipid receptor, a G protein-coupled receptor, activates calcium flux and serum response element driven transcorted.	0.02
AA022569	Hs.29802:35,Hs.2717 85:1	ESTs	ranschour Function unknown	0.02
BE147740	Hs.104558:21	ESTs, Moderately similar to hypothetical protein FLJ20378 [Homo sapiens]	Function unknown	0.02
AI798863	Hs.87191:8	ESTs	Function unknown	0.02
BE464341	Hs.21201:18	Interim-DKFZP566B0846: nectin 3	Low similarity to PVRL1; are a membrane glycoprotein; contains an immunoglobulin (Ig) domain	0.02
AL080235	Hs.35861:34,Hs.2890 68:1	RIS1 Ras-induced senescence 1	Rat brain specific binding protein	0.02
AI557212	Hs.17132:102,Hs.330 782:1	ESTs	Function unknown	0.02
X75208	Hs.2913:41	ЕРНВЗ ЕрһВЗ	Signal transduction, Integral plasma membrane protein, transmembrane receptor protein tyrosine kinase. Eph-related receptor protein kinase R3	0.02
AA628980	Hs.192371:3	DSCR8 Down syndrome critical region protein DSCR8	Melanoma-testis-associated protein 2	0.02
BE242587	Hs.118651:39	HHEX hematopoietically expressed homeobox	Nucleus, DNA binding, transcription factor, developmental processes, antimicrobial humoral response. Member of the homeodomain family of DNA binding proteins; may regulate gene excression, muchocenesis, and differentiation	0.02
NM_005512	Hs.151641:65	GARP glycoprotein A repetitions predominant	Integral plans membrane protein. Putative transmembrane cell surface protein, has an extracellular domain comprised largely of laurine-rich menats.	0.02
AW953853	Hs.292833:19	PAEP progestagen-associated endometrial protein (placental protein 14, pregnancy-associated endometrial alpha-2-globulin, alpha uterine	Developmental processes. Placental protein 14 (Glycodelin); member of lipocalin superfamily, highly similar to beta-lactoglobulins	0.02

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
•		protein)		
AU076611	Hs.154672:122	MTHFD2 methylene tetrahydrofolate dehydrogenase (NAD dependent), methenyltetrahydrofolate cyclohydrolase	Mitochondrion, electron transporter, methenyltetrahydrofolate cyclohydrolase, methylenetetrahydrofolate dehydrogenase. NADdependent methylene tetrahydrofolate dehydrogenase-cyclohydrolase; may provide formyltetrahydrofolate for formylmethlonyl RRNA synthesis; involved in initiation of mitochondrial modeln synthesis.	0.02
AW968613	Hs.79428:166	BNIP3 BCL2/adenovirus E1B 19kD-Interacting protein 3	Anti-apoptosis, apoptosis inhibitor. Bct2-related protein 3; binds antiapoptotic viral E1B 19 kDa protein and cellular Bct2 protein	0.02
AL353944	Hs.50115:14	Homo sapiens mRNA; cDNA DKFZp761J1112 (from clone DKFZp761J1112)	Function unknown	0.02
BE614149	Hs.20814:29,Hs.3066 26:27	LOC51072: C21orf19-like protein	Function unknown	0.02
AA292998	Hs.163900:25	ESTs	Highly similar to winged helix/forkhead transcription factor	0.02
H12912	Hs.274691:138	AK3 adenylate kinase 3	Nucleobase, nucleoside, nucleotide and nucleic acid metabolism. Adenylate kinase 3; strongly similar to murine Ak4	0.02
AA188763	Hs.36793:4	SLC12A8 solute carrier family 12 (potasslum/chloride transporters), member 8	Solute carrier family 12 (potasslum/chloride transporters), member 8	0.02
AK000596	Hs.3618:56	HPCAL1 hippocalcin-like 1	Calclum-binding protein with similarity to hippocalin (human HPCA); expressed only in the brain.	0.02
Al970797	Hs.64859:16	ESTs	Function unknown	0.02
AW519204	Hs.40808:22	ESTs	Function unknown	0.02
Z42387	Hs.83883:114	TMEPAl transmembrane, prostate androgen Induced RNA	Function unknown	0.02
AF145713	Hs.61490:51	SCHIP1 schwannomin-interacting protein 1	Cytoplasm. Associates with the neurofibromatosis type 2 protein schwannomin (NF2); contains a coiled-coil domain	0.02
AA972412	Hs.13755:41	FBXW2 f-box and WD-40 domain protein 2	Protein modification, ublquitin—protein ligase, proteolysis and peptidolysis, ublquitin conjugating enzyme. F-box and WD-40 domain protein 2: putative SCF ublquitin ligase subunit involved in protein and an E-box	0.02
AK001564	Hs.104222:139,Hs.29 6267:4	Homo sapiens cDNA FLJ10702 fis, done NT2RP3000759, weakly similar to ADP-RIBOSYLATION FACTOR	Member of the ADP-ribosylation factor (ARF) family; putative GTP-binding protein involved in protein trafficking	0.02

Accession number	UniGené Mapping	Gene symbol and title	Putative Function	P value	
AW959861	Hs.290943;28	ESTs	Function unknown	0.02	
BE313555	Hs.7252:158	RAI17 retinoic acid induced 17	Function unknown	0.02	
W25005	Hs.24395:199	zb67e02.r1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:308666 5', mRNA	Function unknown	0.02	
A193356	Hs.160316:3	sequence ESTs	Function unknown	0.02	
AF111108	Hs.3382:223	PPP4R1 Protein phosphatase 4, regulatory subunit 1	Protein phosphatase	0.02	
AJ130740	Hs.6241:116	PIK3R1 phosphoinositide-3-kinase, regulatory subunit, polypeptide 1 (p85 alpha)	A family of erzymes that phosphorylate the 3'-hydroxyl of phosphatidylinositol (Ptdins).	0.02	
AA985190	Hs.246875:42	FLJ20059 hypothetical protein FLJ20059	Contains four Kelch motif domains	0.02	
BE221880	Hs.268555:144	XRN2 5'-3' exortbonuclease 2	Nucleus, nuclease, recombination, RNA catabolism, RNA processing, 5'-3' Exoribonuclease; similar to Schizosaccharomyces pombe Dhafa	0.03	
AF084545		Homo saplens versican Vint Isoform, mRNA, partial cds	Function unknown	0.03	
R26584	Hs.267993:43	TAPBP-R: TAP binding protein related	Has low similarity to TAPBP (Tapasin); contains two immunoglobulin (Ig) domains/Proteome	0.03	
AW247380	Hs.12124:116	ELAC2 elaC homolog 2 (€. ∞II)	putative prostate cancer susceptibility protein	0.03	
AA364261	Hs.131365:7	ESTs	Weakly similar to T31613 hypothetical protein Y50E8A.i - Caenorhabditis elegans [C.elegans]	0.03	
U25849	Hs.75393:141	ACP1 Human red cell-type low molecular weight acid phosphatase (ACP1) gene, exon 6 and 7, complete cds	Acid phosphatase	0.03	
AF262992	Hs.123159:14	SPAG4 Sperm associated antigen 4	Spermatogenesis, structural protein. Sperm associated antigen 4; predicted ortholog of rat SPAG4, which Interacts with rat ODF27, the 27kDa outer dense ther protein of elongaling spermatifis.	0.03	
AW342140	Hs.182545:1	ESTs, Weakly similar to POL2_MOUSE Retrovirus-related POL polyprotein	Function unknown	0.03	
AL133572	Hs.199009:58	PCCX2 protein containing CXXC domain 2	DNA-binding protein with PHD finger and CXXC domain, is regulated by proteolysis.	0.03	
		•			

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AI497778	Hs.20509:4	HBXAP Hepatitis B virus x associated protein	Weakly similar to Drosophila CG8677	0.03
AI745379	Hs.42911:31	TAF13 TAF13 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 18 kD	TFIID complex, protein binding, transcription factor, general RNA polymerase if transcription factor. TBP-associated factor K;	0.03
U51712	Hs.13775:135	LAGY: lung cancer-associated Y protein	component of IrilD complexes containing IAFIESU (IAFZH) The protein encoded by this gene is a lung cancer associated protein. The function of the protein is not known. Multiple alternatively spliced transcript variants have been described for this gene but some of their full length sequence has not been	0.03
AW375974	Hs.156704:4	ESTs	aetermineu. Function unknown	0.03
AF251237	Hs.112208:16	GAGED2 G antigen, family D, 2	GAGE genes are expressed in a variety of tumors and in some fetal and reproductive tissues. This gene is strongly expressed in Ewing's sarcoma, alvolar thabdomyosarcoma and normal testis. The protein encoded by this gene contains a nuclear localization signal and shares a sequence similarity with other GAGE/PAGE proteins. Because of the expression pattern and the sequence similarity, this protein also belongs to a family of CT (cancer-testis) antigens.	0.03
. NM_000636		Homo saplens superoxide dismutase 2, mitochondria (SOD2), mRNA. expression) (RFX2), mRNA.	Mitochondrion, oxidative stress response, manganese superoxide dismutase. Manganese superoxide dismutase; intramitochondrial free radical scavenging enzyme; has strong similarity to murine	0.02
AA130986	Hs.271627:1	ESTs	5002. Function unknown	0.01
AA216363	Hs.262958:48,Hs.327 737:2	DKFZP434B044 hypothetical protein DKFZp434B044	Function unknown	0.01
AA628980	Hs.192371:3	DSCR8 down syndrome critical region protein DSCR8	Function unknown	0.00
AA811657	Hs.220913:9	Homo sapiens cDNA FLJ40827 fis, clone TRACH2011500	Function unknown	0.02
AA897108		gb:am08a06.s1 Soares_NFL_T_GBC_S1 Homo sapiens cDNA clone 3', mRNA sequence	Function unknown	0.01
AB040888	Hs.41793:110	Homo sapiens mRNA for KIAA1455 protein, partial cds	Function unknown	0.02

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
AF212225	Hs.283693:104	Homo sapiens BM022 mRNA, complete cds	Function unknown	0.02
AI089575	Hs.9071:52	ESTs	Function unknown	0.02
AI282028	Hs.25205:10	ESTs	Function unknown	0.02
Al368826	Hs.30654:15	FLJ10849: hypothetical protein FLJ10849	Moderately similar to members of the septin family	0.02
AI718702	Hs.308026:11,Hs.194 490:6	HLA-DRB3 major histocompatibility complex, class ii, DR beta 5	Signal transduction, integral plasma membrane protein, class II major histocompatibility complex antigen. Beta 3 chain of HLA-DR; subunit of MHC class II molecule, complex binds peptides and	0.02
AI827248	Hs.224398:3	Homo saplens cDNA FLJ11469 fis, clone HEMBA1001658	presents them to CD4+ T lymphocytes Function unknown	0.01
AK002039	Hs.26243:38	MRVI1 murine retrovirus integration site 1 homolog	Oncogenesis, tumor suppressor, endoplasmic reticulum membrane. Similar to human MLRP; may act as a tumor suppressor	0.02
AL109791	Hs.241559:3	Homo saplens mRNA full length insert cDNA clone EUROIMAGE 151432	Function unknown	0.00
AW090198	Hs.4779:29	LOC127829: hypothetical protein BC015408	Function unknown	0.01
AW296454	Hs.24743:92	FLJ20171: hypothetical protein FLJ2017	Contains three RNA recognition motifs (RRM, RBD, or RNP)	0.02
AW445034	Hs.256578:4	ESTs	Function unknown	0.00
AW452948	Hs.257631:3	ESTs	Function unknown	0.01
AW470411	Hs.288433:27	HNT: neurotrimin	Cell adhesion, neuronal cell recognition, integral plasma membrane protein. Neurotrimin; may function as a GPI-anchored neural cell	0.02
AW885727	Hs.301570:22	FST follistatin	adhesion molecule; member of the immunoglobulin superfamily Developmental processes. Follistatin; inhibits the release of follicleselimitating homogone IESH)	0.01
AW970859	Hs.313503:4	ESTs	Function unknown	0.02
AW979189	Hs.283367:3	ESTs	Function unknown	0.01

•	Accession number	UniGene Mapping	Gene symbol and title	Putative Function	P value
Ι"	BE165866	Hs.83623:66	Human XIST, coding sequence "a" mRNA (locus DXS399E)	XIST mRNA	0.01
	BE175582		gb:RC5-HT0580-100500-022-C01 HT0580 Homo sapiens cDNA, mRNA sequence	Function unknown	0.01
_	BE242587	Hs.118651:39	HHEX hematopoietically expressed homeobox	Nucleus, DNA binding, transcription factor, developmental processes, antimicrobial furmoral response. Member of the homeodomain family of DNA binding proteins; may regulate gene expression, morphonenesis, and differentiation	0.01
	BE271927	Hs.87385:31,Hs.3079 40:4	LOC115416: hypothetical protein BC012331	Function unknown	0.01
	BE439580	Hs.75498:40	SCYA20 small inducible cytokine subfamily A (Cys-Cys), member 20	Chemokine, chemotaxis, immune response, signal transduction, extracellular space, cell-cell signalling, inflammatory response, antimicrobial humoral response. Cytokine A20 (exodus); chemotactic factor for lymphocytes, but not a chemotactic factor for monocytes	0.02
	BE464016	Hs.238956:35	Homo sapiens cDNA FLJ37793 fis, clone BRHIP3000473	Function unknown	0.02
	D63216	Hs.153684:137	FRZB frizzled-related protein	Membrane, extracellular, skeletal development. Frizzled-related protein; similar to frizzled family of receptors	0.02
:	F34856	Hs.292457:120	Homo saplens, done MGC:16362 IMAGE:3927795, mRNA, complete cds	Function unknown	0.02
	M83822	Hs.62354:112	LRBA LPS-responsive vesicle trafficking, beach and anchor containing	May mediate protein-protein interactions; contains two WD domains (WD-40 repeats) and a belge/BEACH domain Proteome	0.02
	N33937	Hs.10336:6	ESTs	Function unknown	0.01
	N49068	Hs.93966:4	ESTs	Function unknown	0.01
	N51357	Hs.260855:62	NSE1: NSE1	Function unknown	0.02
	N80486	Hs.39911:17	Homo saplens mRNA for FLJ00089 protein, partial cds	Function unknown	0.02
	NM_000954 .	Hs.8272:265,Hs.3323 55:1	PTGDS prostaglandin D2 synthase (21kD, brain)	Membrane, prostaglandin-D synthase. Glutathione-independent prostaglandin D2 synthase; membrane associated, catalyzes synthesis of prostaglandin D; member of the lipocalin family of synthase.	0.02
	NM_005756	Hs.184942:18	GPR64 G protein-coupled receptor 64	ransporters Spermatogenesis, G-protein linked receptor, integral plasma membrane protein, G-protein linked receptor protein signalling pathway. Member of the G protein-coupled receptor family	0.02

Accession number UniGene Mapping	UniGene Mapping	Gene symbol and title	Putative Function	P vaíue
NM_016652	Hs.268281:61	CRNKL1 Cm, crooked neck-like 1 (Drosophila)	Function unknown	0.02
R26584	Hs.267993:43	TAPBP-R: TAP binding protein related	Has low similarity to TAPBP (Tapasin); contains two immunoglobulin (ig) domains	0.01
R31178	Hs.287820:6	FN1 fibronectin 1	Cell adhesion, cell motility, cell adhesion, soluble fraction, signal transduction, extracellular matrix, extracellular space. Fibronectin 1; member of family of proteins found in plasma and extracellular matrix.	0.02
W05391	Hs.83623:8	Homo saplens cDNA FLJ30298 fis, clone BRACE2003172	Function unknown	0.02
W25005	Hs.24395:199	zb67e02.r1 Soares_fetal_lung_NbHL19W Homo sapiens cDNA clone IMAGE:308666 5', mRNA sequence	Function unknown	0.01
W45393	Hs.55888:15	ATF7 activating transcription factor 7	Transcription factor. Leucine zipper DNA-binding protein; recognizes a cAMP response element (CRE), involved in the regulation of adenovirus Ela-responsive and cellular cAMP-inducible promoters	0.02
W68815	Hs.301885:20	Homo sapiens cDNA FLJ33794 fis, clone CTONG1000009	Function unknown	0.01
X65965		H.saplens SOD-2 gene for manganese superoxide dismutase	Mitochondrion, oxidative stress response, manganese superoxide dismutase. Manganese superoxide dismutase; intramitochondrial free radical scavenging enzyme; has strong similarity to murine Sod2.	0.01
X76732	Hs.3164:58	NUCB2 nucleobindin 2	Cytosof, DNA binding, plasma membrane, calcium binding, extracellular space. Nucleobindin 2; may bind DNA and calcium; has DNA-binding and Ef-band domains, and a leucine-zloper	0.02
Z45051	Hs.22920:25	C20orf103 chromosome 20 open reading frame 103	Low similarity to a region of murine Lamp1 Proteome	0.02

Accession number UniGene Mapping	UniGene Mapping	Gene symbol and title	Putative Function	P value
c. downregulated genes	Ø			c
NM_022117	Hs.136164:23	SE20-4, cutaneous T-cell lymphoma-associated tumor antigen se20-4se20-4	Cutaneous T-cell lymphoma-associated tumor antigen se20-4se20-4; differentially expressed nucleolar TGF-beta1 target protein	<b>.</b>
NM_005460	Hs.200445:4 Hs.300445:4	SNCAIP, synuciein, alpha interacting protein (synphilin)	County 1, and whom to controlled by Apphilin-1; promotes formation of cytosolic inclusions in neurons (SNCAIP). Syruclein alpha interacting protein contains several protein-protein interaction domains and interacts with alpha synuclein in neurons. Mutations of SNCAIP have been linked to Parkinson disease.	0
NM_002387	Hs.1345:5	MCC, mutated in colorectal cancers	Receptor, signal transduction, tumor suppressor. Similar to the G protein-coupled m3 muscarint acetylcholine receptor. MCC is a candidate for the putative colorectal tumor suppressor gene. The MCC gene product are involved in early stages of colorectal neoplasta in both sporadic and familial tumors.	0
A1745249 A1694200	Hs.23660:30 Hs.356620, Hs.227913:11	Homo sapiens, done MGC:9889 IMAGE:3868330 ESTs	Function unknown Function unknown	0.0009

Table 2 Genes having modified expression in serous ovarian cancer relative to normal ovarian tissue

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
M25809	Hs.64173	ATP6V1B1, ATPase, H+ transporting, lysosomal 56/58kD, V1 subunit B, isoform 1 (Renal tubular acidosis with deafness)	Subunit B1 (beta subunit) of a vacuolar-type H+-ATPase 1; aplcal proton pump that mediates distal nephron acid secretion	1062.30
AW959311	Hs.172012	DKFZP434J037: hypothetical protein DKFZp434J037	Function unknown	227.83
H16423	Hs.82685	Homo saplens mRNA; cDNA DKFZp313F0317 (from done DKFZp313F0317)	Function unknown	74.54
AI733848	Hs.71935	ZNF339, zinc finger protein 339	Zinc finger protein	55.13
AW055308	Hs.31803	NAC1, transcriptional repressor NAC1	Function unknown	52.63
AF034102	Hs.32951	SLC29A2, solute carrier family 29 (nucleoside transporters), member 2	Nitrobenzylthloinosine-insensitive equilibrative nucleoside transporter 2; may act in the uptake of purine and pyrimidine nucleosides	44.34
AI791905	Hs.95549	FL/20273: RNA-binding protein	Contains four RNA recognition motifs (RRM, RBD, or RNP)	43.21
AW296454	Hs.24743	FLJ20171: hypothetical protein FLJ20171	Contains three RNA recognition motifs (RRM, RBD, or RNP)	38.91
Z43989	Hs.82141	Human clone 23612 mRNA sequence	Function unknown	37.89
AL043980	Hs.7886	PELI1, pellino homolog 1 (Drosophila)	Pellino protein	35.20
BE514982	Hs.38991	S100A2, S100 calcium binding protein A2	S100 calclum-binding protein A2; interacts with target proteins to link extracellular stimuli and cellular responses; member of the S100 tissue/cell specific Ca2+-binding protein family	34.53
		Target Exon	Function unknown	34.02
Al811807	Hs.108646	Homo sapiens cDNA FLJ12534 fis, clone NTZRM4000244	Function unknown	32.34
U90441	Hs.3622	P4HA2, procollagen-proilne, 2-oxogiutarate 4- dioxygenase (proline 4-hydroxylase), alpha polypeptide II	Alpha 2 subunit of prolyl 4-hydroxylase; catalyzes the formation of 4-hydroxyproline in collagers.	32.24
T98226	Hs.171952	OCLN, occludin	This gene encodes an integral membrane protein which is located at tight junctions. This protein are involved in the formation and maintenance of the tight junction.	31.56

	Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
	R35343	Hs.24988	Human DNA sequence from clone RP1-233G16 on chromosome Xq22.1-23. Contains the 5' part of a novel gene, ESTs, STSs, GSSs and a putative CpG island		31.22
	BE247295	Hs.78452	SLC20A1, solute carrier family 20 (phosphate transporter), member 1	Sodium-dependent phosphate symporter, acts as a cell-surface receptor for gibbon ape leukemia virus	30.16
	AB037734	Hs.4993	PCDH19, protocadherin C5000394*:gl 12737280 ref XP_006882.2  keratin 18 [Homo saplens]  6633	Protocadherin Function unknown	29.90 29.30
٠	AF212223	Hs.25010	Homo sapiens BM025 mRNA, complete cds	Function unknown	28.85
	AABUZ656	HS.21843	NIF3L1, NIF3 (Ngg1 interacting factor 3, S.pombe homolog)-like 1	Amyotrophic lateral sclerosis 2 (juvenile) chromosome region, candidate 1	27.73
	X14008	Hs.234734	Human lysozyme gene (EC 3.2.1.17)	Lysozyme	27.66
	AA570256		LOC116238: hypothetical protein BC014072	Function unknown	27.52
× .	AA137152	Hs.286049	PSA, phosphoserine aminotransferase	The protein encoded by this gene is likely a phosphosenine aminotransferase, based on similarity to proteins in mouse, rabbit, and Drosophila. Alternative splicing of this gene results in two transcript variants encoding different isoforms.	25.57
	BE621807		TM4SF1, transmembrane 4 superfamily member 1	L6 antigen; member of the transmembrane 4 superfamily (TM4SF)	25.40
	AB041036	Hs.57771	KLK11, kallikrein 11	Trypsin-like serine protease; has serine protease activity	25.05
	F13386	Hs.7888	Homo sapiens clone 23736 mRNA sequence	Function unknown	22.50
<b>,</b>	AA158177	Hs.118722	FUT8, fucosyltransferase 8 (alpha (1,6) fucosyltransferase)	N-linked glycosylation, oligosaccharide biosynthesis, glycoprotein 6-alpha-L-fucosyltransferase. Alpha(1,6)fucosyltransferase (GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha1-6 fucosyltransferase); transfers fucose to N-linked type complex glycopeptides from GDP-Fuc; functions in asparagine-linked glycoprotein oligosaccharide synthesis	21.90
	BE267045	Hs.75064	TBCC, tubulin-specific chaperone c	Tubulin-specific chaperone c; cofactor in the folding pathway of beta-tubulin, mediates the release of beta-tubulin polypeptides committed to the native state.	21.49
		•	NM_005936:Homo sapiens myelold/lymphoid or	Function unknown	20.46

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
AA150864	Hs.790	MGST1, microsomal glutathione S-transferase 1	Microsome, glutathione transferase. Microsomal glutathione S- transferase; catalyzes the conjugation of glutathione to electrophilic compounds; member of a family of detoxication enzymes.	20.35
AW955632	Hs.66666	EST367702 MAGE resequences, MAGD Homo saplens cDNA, mRNA sequence	Function unknown	20.26
AW837046	Hs.6527	QV1-LT0037-150200-069-e09 LT0037 Homo sapiens cDNA, mRNA sequence	Function unknown	19.60
AA286887	Hs.24724	MFHAS1, malignant fibrous histiocytoma amplified sequence 1	The primary structure of its product includes an ATP/GTP-binding site, three feucine zipper domairs, and a feucine-rich tandem repeat, which are structural or functional elements for interactions among proteins related to the cell cycle, and which suggest that overexpression might be oncorenic with respect to MFH.	
AW401864	Hs.18720	PDCD8: programmed cell death 8 (apoptosis-inducing factor)	Mitochondrial apoptosis-inducing factor, flavoprotein inducing chromatin condensation and DNA fragmentation	19.01
AA196241	Hs.73980	zp98f03.r1 Stratagene musde 937209 Homo sapiens cDNA clone IMAGE:628253 6' stmilar to gb:M19309 TROPONIN T, SLOW SKELETAL MUSCLE ISOFORMS (HUMAN);, mRNA sequence	Function unknown	18.82
NM_004998	Hs.82251	MYO1E, myosin IE	Highly similar to class I myosin; may bind proline-rich peptides; contains an Src homology 3 (SH3) and myosin head domain (motor domain)	18.62
AW873704	Hs.320831	C20orf72: chromosome 20 open reading frame 72	Function unknown	18.19
AW361666	Hs.49500	KIAA0746; KIAA0746 protein	Function unknown	18.05
BE174595	Hs.366	PTS, 6-pyruvoyltetrahydropterin synthase	6-Pyruvoyitetrahydropterin synthese; synthesizes tetrahydrobiopterin, activity requires seplapterin reductase, Mg2+, and NADPH	17.28
M31669	Hs.1735	Human inhibin beta-B-subunit gene, exon 2, and complete cds	Function unknown	16.24
AK001714	Hs.85744	FLJ10852, hypothetical protein similar to ankyrin repeat-containing priotein AKR1	Are involved in protein-protein interactions; has five ankyrin repeats and a DHHC-type zinc finger or NEW1 domain	16:09
AU076517	Hs.184276	AU076517 Sugano cDNA library Homo sapiens cDNA done ColF3365 similar to 5-end region of Homo sapiens ezrin-radiwn-moesin binding phosphoprotein- 50 mRNA, mRNA sequence	Function unknown	16.05

Ratio	15.93	15.91	15.35	14.62	14.60	14.37	14.07	14.05	13.75	13.48	12.92	12.86	12.79	12.73	12.58	12.43	12.25	12.22	12.21
Putative Function	Low similarity to beta-galactosidase a-2,3-slalytransferase SIAT4B; member of the slalytransferase family	Function unknown	Suifide dehydrogenase like	Fibroblast growth fador receptor 3; receptor tyrosine kinase that binds acidic and basic FGF	Leucine aminopeptidase	Cytokine A 2; chemotactic factor for monocytes	Ceruloplasmin; ferrous oxidase, binds copper in plasma and maintains iron homeostasis	Mitogen indudble dual specificity protein phosphatase 5; dephosphorylates extracellular signal-regulated Kinase	Function unknown	Very strongly similar to murine Whit7a; may have a role in limb development and sexual dimorphism; member of the Writ family of cell cinnelling profeins.	egreting processing from the property of the processing mediates cell-cell interactions in epithelial cells	Cell-cycle regulated HNF-3/fork head; a transcriptional regulator	Vascular endothellal growth factor, induces endothellal cell proliferation and vascular permeability	Function unknown	CXC chemokine receptor (fusin); G protein-coupled receptor binds CXC cytokines, mediates intracellular calcium flux	Strongly similar to human D6S81E; member of the DEAD/H box ATP-dependent RNA helicase family	Leucine zipper kinase	Binds to the HIV-1 TAR RNA regulatory element, may function alone or with HIV-1 Tat to disengage RNA polymerase II during transcriptional elongation; has a leuche zipper	Function unknown
Gene symbol and title	STHM, sialyltransferase	Homo sapiens cDNA FLJ14201 fis, clone NT2RP3002855	SQRDL: suffide dehydrogenase like (yeast)	FGFR3, fibroblast growth factor receptor 3 (achondroplasta, thanatophoric dwarfsm)	LAP3, leucine aminopeptidase	CCL2, chemokine (C-C motif) ligand 2	CP, ceruloplasmin (ferroxidase)	DUSP5, dual specificity phosphatase 5	EST381684 MAGE resequences, MAGK Homo sapiens cDNA, mRNA sequence	WNT7A, wingless-type MMTV Integration site family, member 7A	CDH1, cadherin 1, type 1, E-cadherin (epithelial)	FOXM1, forkhead box M1	VEGF, vascular endothelial growth factor	FLJ20160; hypothetical protein FLJ20160	CXCR4, chemokine (C-X-C motif), receptor 4 (fusin)	DDX39, DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 39	MELK, maternal embryonic feucine zipper kinase	TARBP1, TAR (HIV) RNA binding protein 1	EST365645 MAGE resequences, MAGC Homo sapiens cDNA, mRNA sequence
UniGene Mapping	Hs.288215	Hs.193063	Hs.8185	Hs.1420	Hs.182579	Hs.303649	Hs.282804	Hs.2128	Hs.86368	Hs.72290	Hs.194657	Hs.239	Hs.73793	Hs.23412	Hs.89414	Hs.311609	Hs.184339	Hs.151518	Hs.303125
Accession number	NM_006456	BE148235	AV653729	AL119671	AA393071	AL048753	Al868872	NM_004419	AW969587	AW161449	BE409838	BE540274	AF022375	AW369278	AF147204	BE242818	NM_014791	U38847	AW953575

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
AIB48095	Hs.67776	ESTs, Weakly similar to T22341 hypothetical protein F47B8.5 - Caenorhabditis elegans [C.elegans]	Homo saplens, clone IMAGE:5455669, mRNA, partial cds	12.08
BE274530	Hs.273333	FLJ10986, hypothetical protein FLJ10986	Member of the FGGY carbohydrate kinase family	11.75
AB020676	Hs.21543	KIAA0869 protein	Function unknown	11.73
		Target Exon	Function unknown	11.69
H48299	Hs.26126:33	CLDN10, claudin 10	Cell adhesion, integral plasma membrane protein, tight junction.	11.67
T34530	Hs.4210	Homo sapiens cDNA FLJ13069 fis, clone NT2RP3001762	Function unknown	11.50
NM_022454	Hs.97984	SOX17, SRY (sex determining region Y)-box 17	SRY-related HMG-box transcription factor SOX17	11.42
AA737033	Hs.7165	Homo sapiens, clone IMAGE:4428577, mRNA, partial cds	Function unknown	10.79
AA433988	Hs.98502:8	MUC16, mucin 16, CA125	Mucin 16. Alias CA125 ovarian cancer antigen	10.52
H91282	Hs.286232	Homo sapiens cDNA: FLJ23190 fis, clone LNG12190	Function unknown	10.50
AW005054	Hs.47883	LOC57118: CamKI-like protein kinase	CamKI-like protein kinase; granulocyte-specific protein kinase that activates ERK/MAP kinase activity; similar to Ca(2+)-calmodulin-dependent hinses I CamKI)	10.49
66969X	Hs.73149	PAX8, paired box gene 8	Depotation missory (Volume)  Member of the paired domain family of nuclear transcription factors; are involved in the ribosome assembly, required for normal thyroid development	10.39
AW382987	Hs.88474:42	Homo saplens cDNA, mRNA sequence	Function unknown	10.21
AW957446	Hs.301711	Homo sapkans, done MGC:23936 IMAGE:3838595, mRNA, complete cds	Function unknown	10.12
AA361562	Hs.178761	POH1: 26S proteasome-associated pad1 homolog	Ubiquitin-dependent protein degradation	10.01
AA834626		RAD54L, RAD54 (S.cerevisiae)-like	Has likely roles in mitotic and meiotic DNA recombination and repair; member of SNF2/SWI2 family of DNA-dependent ATPases	9.85
AI878927	Hs.79284	MEST, mesoderm specific transcript (mouse) homolog	Mesoderm specific protein; member of the alpha/beta hydrolase fold family	.983
AW074266	Hs.23071	LOC85439: stonin 2	Stonin 2	9.74
NM_000947	Hs.74519	PRIMZA, primase, polypeptide 2A (58kD)	Subunit of DNA primase polypeptide 2A; part of the DNA polymerase alpha-primase complex	9.72

Accession number	UniGene Napping	Gene symbol and fitte	Putative Function	Ratio
NM_006187	Hs.56009	OAS3, 2'-5'-oligoadenylate synthetase 3 (100 kD)	Member of the 2'5'-oligoadenylate synthetase family	9.68
AW276858	Hs.81256	S100A4, S100 calcium binding protein A4 (calcium protein, calvasculin, metastasin, murine piacental homolog)	Calcyclin (melastasts-associated) (\$100 calclum-binding protein A4); interacts with targets to link extracellular stimuli and cellular responses; member of the \$100 family of tissue-specific calclum-binding proteins	9.68
T18997	Hs.180372	LOC139231; hypothetical protein BC016683	Function unknown	9.49
AA262294	Hs.180383	DUSP6, dual specificity phosphatase 6	Dual specificity protein phosphatase 6; selectively dephosphorylates and inactivates MAP kinase	9.48
AA220238	Hs.94986	RPP38: ribonuclease P (38kD)	Nucleus, ribonuclease P. Subunit p38 of ribonuclease P ribonucleoprotein; processes 5' ends of precursor tRNAs	9.41
AW505308	Hs.75812	PCK2, phosphoenolpyruvate carboxykinase 2 (mitochondrial)	Phosphoenolpyruvate carboxykinase 2; forms phosphoenolpyruvate by decarboxylation of oxaloacetate at the rate-limiting step of physphononaete.	9.38
A1186431	Hs.296638	PLAB: prostate differentiation factor	Processing inhibitory cytokine; member of a subgroup of the TGF-beta superfamily	9.12
Al095718	Hs.135015	Homo sapiens cDNA FLJ40908 fis, done UTERU2004698, highly similar to Mus musculus mRNA for thrombospondin type 1 domain	Function unknown	9.04
W70171	Hs.76939	UMPK, uridine monophosphate kinase	The protein encoded by this gene catalyzes the phosphorylation of unidine monophosphale to unidine diphosphate. This is the first step in the production of the pyrmidine nucleoside triphosphates required for RNA and DNA synthesis. In addition, an allele of this gene may play a role in mediating nonhumoral immunity to Hemophilus influenzae type B.	8.97
AI580935	Hs.105898	Homo sapiens cDNA FLJ31553 fis, clone NT2Ri2001178	Function unknown	8.90
AB040914 AU076611	Hs.278628 Hs.164872	ShrmL: Shroom-related protein MTHFD2, methylene tetrahydrofolate dehydrogenase (NAD+ dependent), methenyltetrahydrofolate cyclohydrolase	Shroom-related protein NAD-dependent metrylene tetrahydrofolate defrydrogenase-cyclohydrolase; may provide formyltetrahydrofolate for formylmethionyl (RNA synthesis; involved in initiation of mitochondrial protein synthesis	8.87
A1089660	Hs,323401	LOC84661: dpy-30-like protein	dpy-30-like protein	8.71
D13668	Hs.136348:228,Hs.80988:2	OSF-2: osteobiast specific factor 2 (fasciclin I-like)	Celi achesion, skeletal development. Putative bone adhesion protein; similar to the insect protein fasciclin!	8.64
AI798863	Hs.87191	ESTs	Function unknown	8.52

	Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
	U78093	Hs.15154	SRPX, sushl-repeat-containing protein, X chromosome	Putative membrane protein with short consensus repeat (sushi) domains	8.51
	A1669760	Hs.188881	ESTs	Function unknown	8.37
	Al375726	Hs.279918	MGC2198: hypothetical protein MGC2198	Function unknown	8.37
	AW271106	Hs.133294	ESTs	Function unknown	8.30
	AK001782	Hs.15093	HSPC195: hypothetical protein HSPC195	Function unknown	8.18
	AF019226	Hs.8036	RAB3D,RAB3D, member RAS oncogene family	GTP-binding protein; are involved in vesicle transport, member of the RAB family of small GTPases	7.94
	AW968343	Hs.24255	LOC150696: prominin-related protein	Prominin-related protein	7.90
	AF111858	Hs.105039	SLC34A2, solute carrier family 34 (sodium phosphate), member 2	Sodium-dependent phosphate transporter; member of the renal type II co-transporter family	7.87
	AA863360	Hs.26040	Homo sapiens, done MGC:40051 IMAGE:5243005, mRNA.complete ods	Function unknown	7.75
	NM_005764	Hs.271473	DD98: epithelial protein up-regulated in cardnoma, membrane associated protein 17	Up-regulated in malignant epithelial cells of renal cell carcinomas, and in carcinomas of colon, breast and lung	.7.75
	AW360901	Hs.183047	MGC4399: mitochondrial carrier protein	Mitochondrial carrier protein MGC4399	7.71
	AL353944	Hs.60115	Homo sapiens mRNA; cDNA DKFZp761J1112 (from clone DKFZp761J1112)	Function unknown	7.69
	H59789	Hs.42644	TXNI.2, thioredoxin-like 2	Member of the thioredoxin family; has region of moderate similarity to glutaredoxin-like proteins	7.85
	NM_002984	Hs.75703	CCL4, chemokine (C-C motif) ligand 4	Cytokine A4	7.64
	AA642452	Hs.130881	BCL11A, B-cell CLL/lymphoma 11A (zinc finger protein)	May bind nucleic acids; contains three C2H2 type zinc finger domains	7.61
	AA789081	Hs.4029	GAS41: giloma-amplified sequence-41	Similar to the transcription factors AF-9 and ENL	7.46
	H13032	Hs.103378	MGC11034, hypothetical protein MGC11034	Function unknown	7.42
	BE384836	Hs.3454	KIAA1821: KIAA1821 protein	KIAA1821 protein	7.40
	AW067800	Hs.155223	STC2, stanniocalcin 2	Stannlocalcin 2; may regulate metal ion homeostasis and inhibits ohoshate uptake	7.36
•	T55979	Hs.115474	RFC3, replication factor C (activator 1) 3 (38kD)	Subunit of replication factor C (activator 1) 3; activator of DNA polymerases	7.35
	AJ278016	Hs.55565	ANKRD3, ankyrtn repeat domain 3	Ortholog of mouse protien kinase C-associated kinase, putative gene, anklrin like, possible dual-spedificity Ser/Thr/Tyr kinase domain	7.25

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
		NM, 025080:Homo saptens hypothetical protein FLJ22316 (FLJ22316), mRNA, VERSION NM_025078.1 Gl:13376631	Function unknown	7.22
AA084248	Hs.85339:64	GPR39, G protein-coupled receptor 39	GPR39, G protein-coupled receptor 39	7.15
BE620738	Hs.173125	PPIF, peptidytprotyl isomerase F (cydophilin F)	Cyclophilin F (peptidylprotyl Isomerase F); binds the immunosuppressant drug cyclosporin A	7.06
AF072873	Hs.114218	FZD6, frtzzled (Drosophila) homolog 6	Frtzted-6; may function in tissue polarity, development and carcinogenesis; similar to frizzled receptor family, has seven transmembrane domains	7.04
AA852773	Hs.334838	KIAA1866 protein	KIAA1866 protein	6.99
R07566	Hs.73817	CCL3, chemokine (C-C moth) ligand 3	Macrophage inflammatory protein 1 alpha; chemokine	6.98
NM_005211	Hs.174142	CSF1R, colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog	Macrophage colony stimulating factor tyrosine kinase receptor; involved in regulation of growth and differentiation of myeloid cells	6.79
AI752666	Hs.76669	NNMT, nicotinamide N-methyltransferase	Nicotinamide N-methyltransferase; catalyzes the N-methylation of nicotinamide and other pyridines, structurally-related drugs and xenoblokes.	6.52
AF182294	Hs.241578	LOC51691: U8 anRNA-associated Sm-like protein LSm8	Member of the Sm family; core constituent of snRNP complexes	6.50
AA457211	Hs.8858	BAZ1A, bromodomain adjacent to zinc finger domain, 1A	May bind DNA and act as a chromatin-mediated transcriptional regulator; contains a bromodomain and a PHD-finger	6.48
W40262	Hs.146310	zc79f0z.s1 Pancreatic islet Homo sapiens cDNA clone IMAGE:328539 3', mRNA sequence	Function unknown	6.47
AB033091	Hs.74313	KIAA1265 protein	Function unknown	6.45
AA292898	Нв.163900	ESTs, Highly similar to winged hellxforkhead transcription factor [Homo sapiens] [H.sapiens]	Function unknown	6.36
BE613269	Hs.21893	OKFZp761N0624: hypothetical protein OKFZp761N0624	Function unknown	6.35
H25836	Hs.301527	ESTs, Moderately similar to unknown [Homo saplens] [H.saplens]	Function unknown	6.27
AL037228	Hs.82043	NUDT5, nudix (nudeoside diphosphate linked molety X)-type motif 5	NDP-sugar hydrolase; converts ADP-ribose to AMP or ribose 5- phosphate; contains a Muf7 motif	6.25
AV662037	Hs.124740	FLJ30532: hypothetical protein FLJ30532	Function unknown	6.21

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
Ai674383	Hs.22891	wc38h08.x1 NCI_CGAP_Pr28 Homo sapiens cDNA clone IMAGE:2320959 3', mRNA sequence	Function unknown	6.20
AW342140	Hs.182545	ESTs, Weakly similar to POL2_MOUSE Retrovinus-related POL polyprotein (Contains: Reverse transcriptase; Endonuclease) [M.musculus]	Function unknown	6.18
BE560135	Hs.5232	HSPC125, HSPC125 protein	Function unknown	6.17
BE409857	Hs.69499	HSPC132: hypothetical protein HSPC132	Moderately similar to a region of S. cerevislae Ykl053c-ap	6.18
AW972542	Hs.289008	LOC116150: hypothetical protein, MGC:7199	Function unknown	6.16
Al523765	Hs.59236	DKFZP434L0718: hypothetical protein DKFZp434L0718	Function unknown	6.16
NM_014056	Hs.7917	DKFZP564K247: DKFZP564K247 protein	Function unknown	6.08
AI857607	Hs.181301	CTSS, cathepsin S	Cathepsin S; lysosomal cysteine (thlo!) protease that cleaves elastin	6.04
AW247529	Hs.6793	PAFAH1B3, platelet-activating factor acetyltrydrolase, isoform lb, gamma subunit (28kD)	Platelet-activating factor acetylhydrolase gamma; may play a role in brain development	5.98
AK000868	Hs.5570	Homo sapiens cDNA FLJ10006 fis, clone HEMBA1000168, weakly similar to CYLICIN I	Function unknown	6.82
AF053551	Hs.31584	MTX2, metaxin 2	Very strongly similar to murine metaxin 2 (Mm.12941); are involved in mitochondrial protein Import	5.91
Al538613	Hs.298241	TMPRSS3, Transmembrane protease, serine 3	The encoded protein contains a serine protease domain, a transmembrane domain, a LDL receptor-like domain, and a scavenger receptor cysteine-rich domain. This gene was identified as a tumor associated gene that is overexpressed in ovarian tumors.	5.86
U48508	Hs.89631	Human skeletal muscle ryanodine receptor gene (RYR1), exons 103, 104, 105, 106, and complete cds	Function unknown	5.88
T69387	Hs.76364	AIF1, allograft inflammatory factor 1	Allografi Inflammatory factor 1; cytokine Inducible protein associated with vascular Injury	5.86
AC005954	Hs.25527	Homo sapiens chromosome 19, cosmid R28784, complete sequence	Function unknown	5.86
AB037805	Hs.88442	KIAA1384 protein	Function unknown	5.84

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
AL031427	Hs.40094	Human DNA sequence from done 167A19 on chromosome 1p32.1-33. Contains three genes for novel proteins, the DIO1 gene for type I iodothyronine deloctinase (EC 3.8.1.4, TXDI1, ITDI1) and an HNRNP A3 (Heterogenous Nuclear Ribonucleoprotein A3, FBRNP) pseudogene.	Function unknown	5.83
AA340864 X89984 Al355781	Hs.278662 Hs.211563 Hs.242463	CLDN7, claudin 7 BCL7A, B-cell CLL/lymphoma 7A qt94a11.x1 NCI_CGAP_Co14 Homo seplens cDNA clone IMAGE:1962908 3' similar to gb:X74929 KERATIN, TYPE II CYTOSKELETAL 8 (HUMAN);	Similar to murine Cldn7; are an integral membrane protein Similar to the actin-binding protein caldesmon; serine-rich Function unknown	5.76 5.74 5.73
5050000	CANADA II	mRNA sequence		Ţ P
AA370409	ns. 10802	Homo Sapiens CLNA: TLJZ3513 ffs, Gone MEF 11919	rungaon unknown	6
AA310162 AW015534	Hs.169248 Hs.217493	HCS: cytochrome c ANXA2, annexin A2	Somatic cytochrome c (heart cytochrome c) Annexin II (lipocortin-2); enhances osteoclast formation and bone resorption; member of the annexin protein family	5.67 5.64
AA326108	Hs.53631:82	BHLHB3: basic hellx-loop-helix domain containing, class B, 3	Basic helix-loop-helix (bHLH) transcription factors (e.g., DEC1, also called BHLHB2; 604256) are related to Drosophila hairy/enhancer of split proteins. They are involved in the control of proliferation and development during differentiation, particularly in neurons.	5.64
AA120865	Hs.23136	ESTs, Highly similar to THYA_HUMAN Prothymosin alpha [H.saplens]	Function unknown	5.62
AK000517	Hs.6844	NALP2: NALP2 protein	Protein with low similarity to murine Op1	5.54
Z36842	Hs.57548	H.sapiens (xs85) mRNA, 209bp	Function unknown	5.53
AA831552	Hs.268016	Homo sapiens cDNA: FLJ21243 fis, clone COL01164		5.50
AL137578	Hs.27607	Homo saplens mRNA; cDNA DKFZp564N2464 (from clone DKFZp564N2464)	Function unknown	5.50
AA316181	Hs.61635	STEAP, six transmembrane epithelial antigen of the prostate	Six transmembrane epithelial antigen of the prostate; prostate-specific cell-surface antigen	5.46

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
X03835	Hs.1857	ESR1, estrogen receptor 1	Estrogen receptor, nuclear receptor transcription factor activated by liganet-binding, involved in hormone-mediated inhibition of gene	5.42
AI557280	Hs.184270	PT2.1_15_G11.r fumor2 Homo sapiens cDNA 3', mRNA sequence	akhissalar Function unknown	5,41
AW248508	Hs.279727	Homo sapiens cDNA FL/14035 fis, clone HEMBA1004638	Function unknown	5.40
N90866	Hs.276770	CDW52, CDW52 antigen (CAMPATH-1 antigen)	CAMPATH-1 antigen; GPI-anchored protein	5.39
U83115	Hs.161002	AIM1, absent in melanoma 1	Member of the beta gamma-crystallin superfamily of proteins: interactions with the cytoskeleton	5.35
AB007860	Hs.12802	DDEF2, development and differentiation enhancing factor 2	GTPase-activating protein; interacts with members of the Arf and Src family	5.35
Z46223	Hs.176663	H.saplens DNA for immunoglobulin G Fc receptor IIIB	Immunoglobulin G Fc receptor	5.31
BE264974	Hs.6566	TRIP13; thyrold hormone receptor Interactor 13	Interacts with ligand binding domain of thyroid hormone receptor and with human papiliomavirus type 16 (HPV16) E1	5.30
AA194422	Hs.22564	MYO6, myosin VI	Motor, hearing, myosin ATPase, structural protein. Class 6 myosin; motor protein; very strongly similar to murine Myo6	5.27
AF134167	Hs.169487	MAFB, v-maf musculoaponeurotic fibrosarcoma oncogene homolog B (avian)	Very strongly similar to murine Krml; may function as a basic domain- leucine zipper transcription factor	5.25
AA232119	Hs.16085	SH120: putative G-protein coupled receptor	putative G-protein coupled receptor	5.25
W58353	Hs.285123	OSBPL10, oxysterol binding protein-like 10	Member of the oxysteroLbinding protein (OSBP) family; may bind oxygenated derivetives of cholesterol	5.21
AW167128	Hs.231934	ESTs, Weakly similar to A57717 transcription factor EC2 - human [H.sapiens]	Function unknown	5.19
U70370	Hs.84136	PITX1, paired-like homeodomain transcription factor 1	Member of the homeodomain family of DNA binding proteins; may regulate gene expression and control cell differentiation	5.18
N55669	Hs.333823	MRPL13, mitochondrial ribosomal protein L13	Protein of the large 60S ribosomal subunit	5.17
BE298446	Hs.305890	BCL2L1, BCL2-like 1	BCL2-related protein; alternative form bcf-xlong inhibits apoptosis and bcf-xshort induces apoptosis	5.17
AW136551	Hs.181245	Homo saplens cDNA FLJ12532 ffs, clone NT2RM4000200	Function unknown	5.15
AW250380	Hs.109059	HGS, hepatocyte growth factor-regulated tyrosine kinase substrate	Zino-finger protein; interacts with STAM, undergoes tyrasine phosphorylation in response to IL2, CSF2, or HGF	5.13

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
AW002565	Hs.124660	Homo saplens cDNA: FLJ21763 fis, clone COLF6967	Function unknown	5,13
AI637274	Hs.105435	GMDS, GDP-mannose 4,6-dehydratase	GDP-mannose-4,6-dehydratase, epimerase converts GDP-mannose to GDP-mannose-4-keto-8-D-deoxymannose, plays a role in the synthesis of fucosylated oligosaccharides	5.11
NM_003878	Hs.78619	GGH, gamma-glutamyl hydrolase (conjugase, folylpolygammaglutamyl hydrolase)	Gamma-glutamyl hydrolase; has greater exopeptidase activity on methotrexate pentaglutamate than on diglutamate	5.11
AF052112	Hs.12540	LYPLA1, lysophospholipase I	Lysophospholipid-specific lysophospholipase 1; hydrotyzes lysophosphatidyl choline	5.09
AV654694	Hs.82316	IFI44, interferon-induced protein 44	Member of the family of Interferon-alpha/beta inducible proteins; may mediate the antiviral action of Interferon	5.09
R24601		Homo sapiens adenylosuccinate synthetase isozyme (ADSS) mRNA, complete cds	Adenylosuccinate synthetase	5.07
BE019020	Hs.85838	Homo saplens cDNA done IMAGE:2963945 5' similar to TR:O16427 O15427 MONOCARBOXYLATE TRANSPORTER: ;; mRNA sequence	Function unknown	5.04
AW163799	Hs.198365	BPGM, 2,3-bisphosphoglycerate mutase	2.3-bisphosphoglycerate mutase; has synthase, mutase, and phosphatase activities, controls 2,3-diphosphoglycerate metabolism, which is an effector for haemoglobin	5.04
AA278921	Hs.1908	PRG1, proteoglycan 1, secretory granule	Secretory granule proteoglycan 1	5.02
NM_003726	Hs.19126	SCAP1, src family associated phosphoprotein 1	Src kinase-associated phosphoprotein; acts as an adaptor protein; contains a pleckstrin homology domain and an SH3 domain	5.02
AA281167	Hs.111911	ESTs, Weakly similar to T06291 extensin homolog T9E8.80 - Arabidopsis thallana [A.thallana]	Function unknown	5.02
		C9000306*gi 12737280 ref XP_006682.2  keratin 18 [Homo saplens]  6633	Function unknown	5.01
AF098158	Hs.9329	C20orf1, chromosome 20 open reading frame 1	Proliferation-associated nuclear protein; associates with the spindle pole and mitotic spindle during mitosis	5.00

	•			
Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Kallo
AA101043	Hs,151254:19	KLK7, kalilkreln 7 (chymotryptic; stratum comeum)	Epidermal differentiation. Stratum comeum chymotryptic enzyme; senne protease. Growing evidence suggests that many kallikreins are implicated in cardinogenesis and some have potential as novel cancer and other disease biomarkers. Thought to be involved in the proteolysis of intercellular cohesive structures preceding desquamation, which is the shedding of the outsmost laver of the epidemils.	4.87
AF017986	Hs.31386:185	Homo sapiens secreted apoptosis related protein 1 (SARP1) mRNA, partial cds.	Function unknown	4.12
AW960564	Hs.3337:137	TM4SF1: transmembrane 4 superfamily member 1	Pathogenesis, plasma membrane, cell proliferation, N-linked glycosylation, integral membrane protein, integral plasma membrane protein. L6 antigen; member of the transmembrane 4 superfamily (TM4SF). The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface antigen and is highly expressed in different carchiomas.	3.62
W29092	Hs.7678:40	CRABP1 Cellular retinoic acid binding protein 1	Cytoplasm, retinoid binding, signal transduction, developmental processes. Cellular retinoic acid-binding protein 1; are involved in delivering retinoic acid to the nucleus, assumed to play an important role in retinoic acid-mediated differentiation and proliferation processes.	3.34
. Н83366	Hs.7587:84	Homo sapiens cDNA: FLJ21962 fis, clone HEP05584	Function unknown	3.29
D49441	Hs.155981:53	MSLN, mesothelin	Cell adhesion, cell surfaces antigen, membrane. Pre-pro-megakaryocyte potentiating factor. An antibody that reacts with ovarian cancers and mesotheliomas was used to isolate a cell surface antigen named mesothelin. Although the function of mesothelin is unknown, it may play a role in cellular adhesion and is present on mesothelium, may blay mesothelioms, and ovarian cancers.	3.14
AA214228	Hs.127751:21,Hs.78006:5	C20orf180: chromosome 20 open reading frame 180	Region of high similarity to tyrosine-phosphorylated protein DOK1	2.99
M31126	Hs.272620:1	PSG9: pregnancy specific beta-1-glycoprotein 9	Pregnancy, extraodilular, plasma glycoprotein. Member of the pregnancy-specific glycoprotein (PSG) and QEA families.	2.82
<b>U62801</b>	Hs.79361:65	KLK6, kalikrein 6 (neurosin, zyme)	Serine type peptidase, pathogenesis. Neurosin (protease M, zymė); a serine protease trat cleaves amyloid precursor protein (APP). Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease blomarkers.	2.77
AK001536	Hs.285803:6	Homo sapiens cDNA FLJ12852 ffs, clone NT2RP2003445	Function unknown	2.73

Accession number	UnlGene Mapping	Gene symbol and title	Putative Function	Ratio
NM_014767	Hs.74583:140 Hs.75733:129,Hs.278399:100,H s.274378:1	KIAA0275: KIAA0275 gene product AMY2A: amylase, alpha 2A; pancreatic	Function unknown Alpha-amylase, extracellular space, carbohydrate metabolism. Parcreatic alpha-amylase 2A (1.4-alpha-D-glucan glucanotrydrolase); deaves internal a-1,4 bonds between glucose monomers to digest	2.72
AA430348	Hs.288837:40	Homo saplens cDNA FLJ12927 fis, clone NT2RP2004743	starch. Function unknown	2.69
X51630	Hs.1145:22,Hs.296851:1	WT1, Wilms tumor 1	Nucleus, transcription factor, transcription regulation. 4 Zn finger domains. Functions in kidney and gonad proliteration and differentiation. Mutations in this gene are associated with the development of Wilms tumors in the kidney or with abnormalities of the genitourinary tract.	2.58
BE393948	Hs.50915:17	KLK5, kalilkreln 5	Serine type peptidase, epidermal differentiation, extracellular space. Stratum comeum tryptic errzyme (kallikrain-like protein), may function in epidermal stratum comeum desquamation and turnover. Expression in protate cancer negatively correlated with cancer aggressiveness	2.34
NM_002776	Hs.69423:46	KLK10, kaliikrein 10	If coses 2002) Putative serine protease. Expressed in normal breast tissue and benign lesions, with loss of expression during tumor progression (Dhar 2001). SINPs associated with prostate, breast, testicular, and ovarian cancers	2.24
NM_000954	Hs.8272:294	PTGDS: prostaglandin D2 synthase (21kD, brain)	(prinaraj zouzz). Membrane, prostaglandin-D synthase. Glutathione-independent prostaglandin D2 synthase; membrane associated, catalyzes synthesis of prostaglandin D2, member of the lipocalin family of transporters.	2.15
AB029000	Hs.70823:109,Hs.297970:48	KIAA1077: sulfatase FP	Function unknown	2.04
AL044315 AA334592	Hs.173094:70 Hs.79914:337	KIAA1750; KIAA1750 protein LUM: lumican	Function unknown Vision, proteoglycan, extracellular matrix, cartilage condensation, extracellular matrix glycoprotein. Member of the specialized collagens	0.95
S79895	Hs.83942:248	CTSK: cathepsin K (pycnodysostosis)	and SLRP_family Lysosome, cathepsin K, cysteine-type peptidase, proteolysis and peptidolysis. Cathepsin K (cathepsin O), a cysteine (thiol) protease; involved in bone remodeling and reabsorption	0.91
Al091195	Hs.65029:120	Homo saplens cDNA clone IMAGE:1566910 3', mRNA sequence	Function unknown	0.91

Accession number	UniGene Mapping	Gene symbol and title	Putative Function	Ratio
AF026692; NM_003014	Hs.105700:83,Hs.278611:3	SFRP4: secreted frizzled-related protein 4	Member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wht-binding site of Frtzzled proteins. SFRPs act as soluble modulators of Whit signaling. The expression of SFRP4 in venticular myocardium correlates with apoptosis related gene	0.73
AI683243	Hs.97258:15	ESTs, Moderately similar to S29539 ribosomal protein L13a, cytosolic	Function unknown	-2.96
AI287700	Hs.111128:7	Homo sapiens, clone IMAGE:4106329, mRNA	Function unknown	-5.71
AA291377	Hs.50831:23	Homo saplens Ly-6 antigen/uPA receptor-like domain- containing protein mRNA, complete cds	Function unknown	-6.78
AI420213	Hs.149722:3	cDNA clone IMAGE:2094208 3', mRNA sequence	Function unknown	-8.52
AJ245671	Hs.12844:73	EGFL6, EGF-lke-domain; multiple 6	Cell cycle, oncogenesis, integrin ligand, extracellular space. Member of the epidermal growth factor (EGF) repeat superfamily; contains an EGF-like-domain. Expressed early during development, and its expression has been detected in lung and meningioma fumors.	-9.44
AB018305	Hs.5378:149	SPON1, spondin 1, (f-spondin) extracellular matrix protein	Extracellular matrix protein. Very strongly similar to rat F-spondin (Rn.7546); may have a role in the growth and guidance of axons.	-12.55
AW872527	Hs.59761:19	ESTS; Weakly similar to DAP1_HUMAN DEATH-ASSOCIATED PROTEIN 1	Function unknown	-14.17
AF129755	Hs.117772:9,Hs.88474:1	Homo sapiens prostaglandin endoperoxide H synthase-1 mRNA, partial 3' untranslated region.	Function unknown	-21.34
A1023789	.Hs.163242;5	· Homo sapiers cDNA done IMAGE:1655725 3' similar to contains MER20.12 MER20 repetitive element;	Function unknown	41.34

Table 3

		Ð		
	۵	value o	<b>o</b> ,	o ·
ince thereof	Chromosome	<b>Location</b> 5q23.2	5922.2	unmapped
cancer or a recurre in cancer	SEQ ID NO:	SEQ ID NO: 1 (DNA) SEQ ID NO: 2 (PRT)	SEQ ID NO: 3 (DNA) SEQ ID NO: 4 (PRT)	SEQ ID NO: 5 (DNA) SEQ ID NO: 6 (PRT)
Preferred diagnostic and prognostic markers for detecting ovarian cancer or a recurrence thereof or survival of a subject suffering from ovarian cancer		Function Cytoplasm, pathogenesis, protein binding. Synphilin-1; promotes formation of cytosolic inclusions in neurons (SNCAIP). Synuclein alpha interacting protein contains several protein-protein interacting domains and interacts with alpha synuclein in neurons. Mutations of SNCAIP have been linked to Parkinson disease.	Receptor, signal transduction, tumor suppressor. Similar to the G protein-coupled m3 muscarinic acetylcholine receptor. MCC is a candidate for the putative colorectal tumor suppressor gene. The MCC gene product are involved in early stages of colorectal neoplasia in both sporadic and familial tumors.	Cutaneous T-cell lymphoma-associated tumor antigen se20-4se20-4; differentially expressed nucleolar TGF- beta1 target protein (DENTT); also known as CDA1
liagnostic and p or st <b>nes</b>		Gene Name SNCAIP, synuclein, alpha interacting protein (synphilin)	MCC, mutated in colorectal cancers	SE20-4, cutaneous T-cell lymphoma- associated tumor antigen se20- 4se20-4
Preferred diagn  A. DOWN-REGULATED GENES	Unigene	<b>Mapping</b> Hs.24948:32; Hs.300445:4	Hs.1345;5	Hs.136164:23
A. DOWN-F	Accession	NW_005460	NM_002387	A1420582; NM_022117

			Table 3 continued	:	į	1
Accession Number B. UP-REGUI	Accession Unigene Number Mapping B. UP-REGULATED GENES	Gene Name	Function	SEQ ID NO:	Chromosome Location	P vafue
BC006428; NM_016463	Hs.15093:210,Hs. 290304:1	HSPC195, hypothetical	Homo sapiens cDNA FLJ10920 fis, clone OVARC1000384-resourcerer.	SEQ ID NO: 7 (DNA) SEQ ID NO: 8 (PRT)	5q31.2	0
NM_017697	Hs.24743:94	protein HSPC185 FLJ20171, hypothetical	contains 3 RNA recognition motifs	SEQ ID NO: 9 (DNA) SEQ ID NO: 10 (PRT)	8q22.1	0
AW630088; NM_001306	Hs.76550:164	protein FLJ20171	Mat2 T-cell differentiation protein; found thru interaction with TPD52 which is overexpressed in breast cancer, 4 TM are involved in vesicle transport	SEQ ID NO: 11 (DNA) SEQ ID NO: 12 (PRT)	8q24.12	<b>o</b> ·
NM_015238	Hs.21543:36	KIAA0869, KIAA0869 protein;	Function unknown	SEQ ID NO: 13 (DNA) SEQ ID NO: 14 (PRT)	5q34	0.0002
AA284679	Hs.25640:264,Hs. 5372:2	KIBKA CLDN3, claudin 3	Integral plasma membrane protein, pathogenesis, tight junction, transmembrane receptor. Member of the claudin family of integral membrane proteins; receptor for Clostridium perfringens enterotoxin;	SEQ ID NO: 16 (DNA) SEQ ID NO: 16 (PRT)	7q11.23	0.0004
NM_022454	Hs.97984:22	SOX17, SRY (sex determining region Y)-box 17	Likely ortholog of mouse SRY-box containing gene 17; alias SOX17	SEQ ID NO: 17 (DNA) SEQ ID NO: 18 (PRT)	8q11.23	0.0005
NM_005682	Hs.6527:201	GPR56, G protein-coupled receptor 56	cell adhesion, cell-cell signalling, G-protein linked receptor, integral plasma membrane protein, G-protein linked receptor protein signalling pathway. Member of the G protein-coupled receptor family; similar to secretin and calcitonin receptors. 7 transmembrane domains, a much-like domain and cysteine box in the N-terminal region. Expressed in range of tissues, highest levels in tryroid, selectively within the monolayer of cuboidal epithelial cells of the smaller, more actively secreting follicles of human thyroid. Differentially expressed in melanoma cell lines with different mefastatic potential	SEQ ID NO: 20 (PRT)	16q13	0.0012
NM_001307	Hs.278562:101	CLDN7, claudin 7	Integral membrane protein, tight junction. Similar to murine Cldn7;	SEQ ID NO: 21 (DNA) SEQ ID NO: 22 (PRT)	17p13.1	d.0016

0.0025	0.0055	9000
15q31 20q13.12	6p21.33	17q21.2
SEQ ID NO: 23 (DNA) SEQ ID NO: 24 (PRT) SEQ ID NO: 25 (DNA) SEQ ID NO: 26 (PRT)	SEQ ID NO: 27 (DNA) SEQ ID NO: 28 (PRT)	SEQ ID NO: 29 (DNA) SEQ ID NO: 30 (PRT)
function unknown; no significant hits with Superfamily Plasma protein, proteinase inhibitor. Secreted inhibitor which protects epithelial tissues from serine proteases. Found in various secretions including seminal plasma, cervical mucus, and bronchial secretions, has affinity for trypsin, leukocyte elastase, and cathepsin G. its inhibitory effect contributes to the immune response by protecting epithelial surfaces from attack by endogenous proteolytic enzymes; the protein is also thought to have broad-spectrum anti-biotic activity.	Cell adhesion, integral plasma membrane protein, transmembrane receptor, protein tyrosine kinase. Epithelial-specific receptor protein tyrosine kinase; are involved in cell adhesion; has putative discoidin motifs in extracellular domain. DDR1 (CD167a) is a RTK that is widely expressed in normal and transformed epithelial cells and is activated by various types of collagen.	DNA binding, DNA topoisomerase (ATP-hydrolyzing), nucleus, DNA topoisomerase il alpha; may relax DNA torsion upon replication or transcription. Involved in processes such as chromosome condensation, chromatid separation, and the relief of torsional stress that occurs during DNA transcription and replication. Catalyzes the transient breaking and rejoining of two strands of duplex DNA. The gene encoding this enzyme functions as the target for several anticancer agents and a variety of mutations in this gene have been associated with the development of drug resistance. Reduced activity of this enzyme may also play a role in ataxia-telangiactasia.
KIAA0101 gene product SLPI, secretory leukocyte protease inhibitor (antifeukoproteina se)	DDR1, discoldin domain receptor family, member 1	TOP2A, topoisomerase (DNA) if alpha (170kD)
Hs.251754:128,H s.245742:1	Hs.75582:147	Hs.156346:184,H s.270810:2
NM_014736 BE184455; NM_003064	NM_013994	NM_001067

0.0131	0.0183	0.0006	0.0074	0.0004	0.0007	0.0009	0.0044	0.0051	0.0051
3p23		12p13.31		1932.1		2q13		2q35	5013.12
SEQ ID NO: 31 (DNA) SEQ ID NO: 32 (PRT)	SEQ ID NO: 33 (DNA) SEQ ID NO: 34 (PRT)	SEQ ID NO: 35 (DNA) SEQ ID NO: 36 (PRT)	SEQ ID NO: 37 (DNA) SEQ ID NO: 38 (PRT)	SEQ ID NO: 39 (DNA) SEQ ID NO: 40 (PRT)	SEQ ID NO: 41 (DNA) SEQ ID NO: 42 (PRT)	SEQ ID NO: 43 (DNA) SEQ ID NO: 44 (PRT)	SEQ ID NO: 45 (DNA)	SEQ ID NO: 46 (DNA) SEQ ID NO: 47 (PRT)	SEO ID NO: 48 (DNA)
chemokine-like factor gene superfamlly; transmb 4 superfamlly	ATP binding, GTP binding, cell proliferation, mitosis, nucleus spindle. Proliferation-associated nuclear protein; associates with the spindle pole and mitotic spindle during mitosis	Plasma membrane, integral plasma membrane protein. Member of the transmembrane 4 superfamily (TM4SF); may mediate platelet activation and aggregation. Cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily proteins.	Amiloride-sensitive sodium channel (weakly similar to Mus musculus PDZ domain actin binding protein)	Embryagenesis and morphogenesis, transcription co- activator, transcription factor, transcription from Pol II promoter. ETS domain transcriptional activator; activates expression of epithelial cell specific genes.	Contains four RNA recognition motifs (RRM, RBD, or RNP)	Histogenesis and organogenesis, embryogenesis and morphogenesis, thyrold-stimulating hormone receptor, transcription factor. Member of the paired domain family of nuclear transcription factors; are involved in the ribosome assembly, required for normal thyroid development. PAX genes play critical roles during fetal development and cancer growth.	Function unknown	Function unknown	59% identity to himan Zinc finger protein 91
CKLFSF7; chemokine-like factor super family	r C20orf1, chromosome 20 open reading frame 1	CD9: CD9 antigen (p24)	ShrmL, Shroom- related protein (KIAA1481	ELF3, E74-like factor 3 (ets domain transcription factor, epithelial- snecfin)	FLJ20273, RNA- binding protein	PAX8, paired box gene 8	EST	FLJ10116, hypothetical	hvnothefical
Hs.343214	Hs.9329:152	Hs.1244:27,Hs.2 30558:1,Hs.2420 20:1	Hs.278628:52	Hs.166096:170	Hs.95549:147,Hs. 229556:1	Hs.73149:72,Hs.2 13008:1	Hs.290801:35, Hs.356228	Hs.79741:18	He 124740-18
BE386983; NM_138410	AF098158; NM_012112	NM_001769	NM_020859	NM_004433	AI791905; NM_019027	X69699; NM_013952	Al301558	NM_018000	NM 144724

AF111856; NM_006424	Hs.105039:48	SLC34A2, solute carrier family 34 (sodlum phosphate), member 2	SLC34A2: solute carrier family 34 (sodium phosphate), member 2; contains 8 predicted TMs and a cysteinerich N-terminal region. Type 2 sodium-dependent phosphate transporter, member of the renal type !! cotransporter family.	SEQ ID NO: 50 (DNA) SEQ ID NO: 51 (PRT)	4p15.2	0.0121
AW959311	Hs.87019:8; Hs.172012	EST DKFZp434J037	probable serine/threonine protein kinase; KIAA0537	SEQ ID NO: 52 (DNA)	1q32.1	0.0251
AF111713	Hs.286218:64	JAM1, junctional adhesion molecule	Cell motility, inflammatory response, intercellular junction. Role in the regulation of tight junction assembly in epithelia. Ligation of JAM is required for reovirus-induced activation of NF-kappa-B and apoptosis. Role in lymphocyte homing.	SEQ ID NO: 53 (DNA) SEQ ID NO: 54 (PRT)		0.0261
AU076611; NM_006636	Hs.154672:123	MTHFD2, methylene tetrahydrofolate dehydrogenase (NAD+ dependent); methenyltetrahydr ofolate	Electron transporter, methenyltetrahydrofolate cydohydrolase, mitochondrion, encodes a nuclearencoded mitochondrial bifunctional enzyme with methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate cyclohydrolase activities, may provide formyltetrahydrofolate for formylmethioryl tRNA synthesis; involved in initiation of mitochondrial protein synthesis.	SEQ ID NO: 56 (DNA) SEQ ID NO: 56 (PRT)	2p13.1	0.0342

. 6p12

0.0001

Table 3 continued

0.01

Chromosome Location		19q13.2	7.77
SEQ ID NO:	SEQ ID NO: 58 (PRT)	SEQ ID NO: 59 (DNA) SEQ ID NO: 60 (PRT)	SEQ ID NO: 61 (DNA) SEQ ID NO: 62 (PRT)
Gene Name Function JES IN MUCINOUS OVARIAN CANCER ONLY	Lectin, cytosol, cell adhesion, plasma membrane. Binds to beta galactoside, involved in cell adhesion, cell growth regulation, inflammation, immunomodulation, apoptosis and metastasis; member of a family of lectins. LGALS4 is an S-type lectin that is strongly underexpressed in colorectal cancer.	Cell adhesion, integral plasma membrane protein, membrane fraction, small molecule transport, transporter. Member of the cadherin family of calciumdependent glycoproteins; facilitates uptake of peptidebased drugs, may mediate cell-cell interactions. Component of the gastrointestinal tract and pancreatic ducts, intestinal proton-dependent peptide transporter in the first step in oral absorption of many medically important peptide-based drugs.	metalloprotease located apically and secreted by epithelial cells in normal colon; degrades broad range of ECM components in vitro; proposed role in tumour progression by facilitating migration, infravasation and metastasis
Gene Name	LGALS4, lectin, galactoside- binding, soluble, 4 (galectin 4)	CDH17, cadherin 17, Ll cadherin (liver-intestine)	MEP1A, meprln A alpha, PABA peptide hydrolase
Unigene Mapping LATED GENES	Hs.5302:132	Hs.89436:50	Hs.179704
Accession Unigene Number Mapping C. UP-REGULATED GEN	AAS84890; NM_008149	·	NM_005588

			Table 3 continued			
Accession	Unigene			SEQ ID NO:	Chromosome	۵
Number	Mapping	<b>Gene Name</b>	Function	·	Location	value
D. PROGNOS	D. PROGNOSTIC MARKERS	₽ P	R SURVIVAL OR RECURRENCE			;
NM_015092	Hs.278428	005; EDD	Homo sapiens progestin induced protein (DDS), mRNA. EDD; Soluble fraction, cell proliferation, ubiquitin-protein ligase, ubiquitin conjugating enzyme, ubiquitin-pependent protein degradation. Member of the HECT family of proteins; may function as an E3 ubiquitin-protein ligase. This gene is localized to chromosome 8q22, a locus disrupted in a variety of cancers. This gene proteintially has a role in regulation of cell multiferation or differentiation.	SEQ ID NO: 63 (DNA) SEQ ID NO: 64 (PRT)	.*	0.00
					8q22.3	
BE465867; NM_014992	Hs.197751:66	DAAM1	dishevelled associated activator of morphogenesis 1 The protein encoded by this gene contains FH domains and belongs to a novel FH protein subtarnity implicated in cell polarity, thought to function as a scaffolding profein.	SEQ ID NO: 65 (DNA) SEQ ID NO: 66 (PRT)	14023.1	0.04
AA381553; NM_002122	Hs.198253:21	. HLA10A	procompatibility complex, class II, DQ alpha 1 Pathogenesis, class II major histocompatibility complex antigen. Alpha 1 chain of HLA-DQ1 class II molecule (la antigen); complex binds peptides and presents them to CD4+ T lymphocytes Proteome	SEQ ID NO: 67 (DNA) SEQ ID NO: 68 (PRT)		0.00
					6p21.3	
AF026692; NM_003014	Hs. 105700:83, Hs. 278611:3	SFRP4: secreted frizzled-related protein 4	Member of the SFRP family that contains a cysteinerich domain homologous to the putative Wnt-binding site of Frizzled proteins. SFRPs act as soluble modulators of Wnt signaling. The expression of SFRP4 in verificular myocardium correlates with apoptosis related gene expression.	SEQ ID NO: 69 (DNA) SEQ ID NO: 70 (PRT)	7p14	0.73
AW015534;	Hs.217493	ANXA2, annextn	Annexin II (lipocortin-2); enhances osteoclast formation	SEQ ID NO: 71 (DNA)	15921-22	0.00
NM_004039		A2	and bone resorption; member of the annexin protein family	SEQ ID NO: 72 (PRT)	ļ	
BE24669;			STAT induced STAT-inhibitor 3; suppressor of cytokine	SEQ ID NO: 73 (DNA) SEQ ID NO: 74 (PRT)	!	6
NM_003955 A1677897	Hs.345728	SOCS3	signalling 3; suppression of IL-6 mediated signalling	SEQ ID NO: 75 (DNA)	17q25.3	0.07
NM_014059	Hs.76640	RGC32	RGC32, hypothetical protein, unknown function	SEQ ID NO: 76 (PRT)	13q13.3	0.04

SAA1 senim Senim amviold A1: high density (innanonmytein: mile in SEQ ID NO: 78 (PRT)	amylold A1	hypothetical fireman	GJB2, gap SEQ ID NO: 81 (DNA)		26 NOV1;	Nephroblastoma SEQ ID NO: 84 (PRT) overexpressed Role in cell adhesion and migration in endothelial cells:	eueß
,	Hs.332053	He 404800	900000		Hs.323733		Hs.235935
AA829286	NM_000331	AA243489;	topolo laik	M86849;	NM_004004		NM_002514

Table 4

Correlation of expression between normal ovarian surface epithelium (OSE), non-invasive tumors (borderline, BL) and ovarian cancer (CA) as determined by ANOVA

	CA125	MUC-1	E-	CLDN3	Ep-CAM	SOX17
			cadherin			
OSE vs IC	<0.0001	<0.0001	0.7251	0.6132	0.1573	0.0854
OSE vs. BL	0.1765	<0.0001	0.0307	0.3633	0.0005	0.2287
OSE vs. CA	0.5443	<0.0001	0.1687	0.0008	<0.0001	0.6900
IC vs. BL	<0.0001	<0.0001	0.1116	0.7849	0.0913	0.2530
IC vs. CA	<0.0001	0.2707	0.4147	0.0071	0.0002	0.0544
BL vs. CA	0.0001	<0.0001	0.0615	<0.0001	0.0011	0.0152

5

## Table 5 Correlation of gene expression with patient outcome (univariate analysis ie., expression alone without the influence of covariates)

Univariate analysis for clinicopathological variables and CLDN3, Ep-CAM, SOX17, CA125, MUC1 and E-cadherin immunoreactivity with survival and relapse in 156 patients with epithelial ovarian cancer

	Disease Specific Survival Univariate		Relapse Free Survival	· .
			Univariate	
Variable .	Hazards ratio (95% CI)	p-value	Hazards ratio (95% CI)	p-value
Pathological tumor stage				
Stage 1 - 3b vs. 3c - 4b	5.89 (3.214-10.79)	< 0.0001	7.37 (3.26-16.63)	<0.0001
Tumor grade			, ,	
BL and G1vs. G2 and G3	5.508 (2.745-11.052)	< 0.0001	7.02 (2.76-17.82)	<0.0001
Age				
<50 vs.>=50	0.533 (0.288-0.988)	0.0458	0.62 (0.29-1.33)	0.2221
Residual Disease				
RD<1cmvs. >=1cm	4.192 (2.671-6.580)	< 0.0001	4.17 (2.30-7.55)	< 0.0001
CA125 level at diagnosis	, ,		•	
CA125 <500vs. >500 U/ml	1.843 (1.102-3.080)	0.0197	2.292 (1.19-4.40)	0.0128
Performance Status			•	
PS<1 vs. >1	0.270 (0.133-0.549)	0.0003	0.53 (0.16-1.74)	0.2965
CLDN3 expression				
Membranous Score 0vs. >0	2.794 (1.012-7.718)	0.0474	2.521 (0.908-6.998)	0.0758
Membranous Score <1 vs. >1	1.309 (0.763-2.246)	0.3285	1.952 (1.103-3.457)	0.0217
Ep-CAM expression				
Membranous Score <1vs. >1	1.460 (0.809-2.634)	0.2093	2.041 (0.997-4.177)	0.0509
Membranous Score <2vs. >2	1.041 (0.634-1.711)	0.873	1.449 (0.845-2.487)	0.1779
SOX17 expression				
Nuclear membranous Score 0vs. >0	0.839 (0.514-1.368)	0.481	1.311 (0.728-2.358)	0.3667
Nuclear membranous Score <1vs. >1	1.407 (0.615-3.218)	0.4183	1.037 (0.380-2.829)	0.9437
CA125 expression				
Membranous apical Score 0vs. >0	. 2.581 (1.393-4.781)	0.0026	2.725 (1.218-6.093)	0:0146
Membranous apical Score <1vs. >1	1.637 (1.045-2.564)	0.0313	1.298 (0.731-2.307)	0.3737
MUC1 expression				
Membranous apical Score 0vs. >0	2.479 (0.343-17.898)	0.368	NA	
Membranous apical Score <1vs. >1	3.745 (1.176-11.926)	0.0254	6.432 (1.562-26.483)	0.0099
Membranous apical Score <2vs. >2	1.814 (0.898-3.664)	0.0969	3.893 (1.552-9.766)	0.0038
E-cadherin expression			•	
Membranous Score 0vs. >0	0.806 (0.493-1.318)	0.3892	0.837 (0.477-1.467)	0.5341
Membranous Score <1vs. >1	1.331 (0.532-3.333)	0.5411	0.847 (0.263-2.731)	0.7814
Membranous Score <2vs. >2	0.593 (0.082-4.284)	0.6041	0.913 (0.125-6.646)	0.9284

## Table 6 Correlation of gene expression with patient outcome (multivariate analysis ie looking at expression incorporating the influence of covariates)

5

Multivariate analysis for univariate significant clinicopathological variables and CLDN3, Ep-CAM, SOX17, CA125, MUC1 and E-cadherin immunoreactivity with survival and relapse in 156 patients with epithelial ovarian cancer

	Disease Specific Survival		Relapse Free Survival		
	Multivariate		Univariate		
Variable	Hazards ratio (95% CI)	p-value	Hazards ratio (95% CI)	p-value	
Pathological tumor stage					
Stage 1 - 3b vs. 3c - 4b	5.66 (2.467-13.012)	<0.0001	5.192 (1.860-14.496)	0.0017	
fumor grade					
BL and G1 vs. G2 and G3	4.919 (2.080-11.633)	0.0003	7.989 (2.385-26.760)	0.0008	
∖ge	• •				
<50 vs.>=50	0.951 (0.482-1.877)	0.8853		•	
Residual Disease					
RD<1cm vs. >=1cm	2.974 (1.783-4.959)	<0.0001	2.779 (1.433-5.393)	0.0025	
CA125 level at diagnosis	•				
CA125 <500 vs. >500 U/ml	1.148 (0.625-2.109)	0.6563	1.289 (0.659-2.520)	0.4587	
Performance Status					
PS<1 vs. >1	0.286 (0.136-0.601)	0.0009			
CLDN3 expression					
Membranous Score 0 vs. >0	1,165 (0.325-4.183)	0.8145	•		
Membranous Score <1 vs. >1	•	•	0.953 (0.473-1.919)	0.8918	
A125 expression	•				
Membranous apical Score 0 vs. >0	0.917 (0.415-2.025)	0.8302	0.693 (0.271-1.768)	0.4427	
Membranous apical Score <1 vs. >1	1.664 (0.976-2.837)	0.0612			
/IUC1 expression					
Membranous apical Score 0 vs. >0					
Membranous apical Score <1 vs. >1	0.678 (0.255-1.804)	0.4361			
Membranous apical Score <2 vs. >2					

WO 2004/022778 PCT/AU2003/001166

181

## WE CLAIM:

5

10

- 1. A method of detecting an ovarian cancer-associated transcript in a biological sample, the method comprising contacting the biological sample with a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Table 1 or 2 or 3.
- 2. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- 20 (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- 25 (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).

15

20

25

- 3. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 5, 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 5, 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 5, 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 6, 8, 10, 12, 14, 16, 18, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 47, 49, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
  - (v) a sequence that is complementary to (i) or (ii) or (iii) or (iv).
  - 4. The method of claim 2 or 3 wherein the hybridization is enhanced in the sample from the subject being tested compared to the hybridization obtained for a sample from a control subject not having ovarian cancer.
  - 5. The method of claim 2 or 3 wherein the hybridization is reduced in the sample from the subject being tested compared to the hybridization obtained for a sample from a control subject not having ovarian cancer.

25

30

- 6. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an enhanced level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM 002387, Al745249 and Al694200;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iv).
  - 7. The method of claim 6 wherein the nucleic acid probe comprises a sequence selected from the group consisting of:
  - (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;

25

- (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 7, 9, 11, 13, 15, 17, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 45, 46, 48, 52, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- 5 (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 10, 12, 14, 16, 18, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 47, 49, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
- (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iii) or (iv).
  - 8. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a reduced level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- 20 (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, AI745249 and AI694200;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
  - (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iii) or (iv).
- 9. The method of claim 8 wherein the nucleic acid probe comprises a sequence selected from the group consisting of:

15

20

25

30

- a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, and 5;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 1, 3, and 5;
- (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 1, 3, and 5;
- (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ.ID NOs: 2, 4, and 6; and
- 10 (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iii) or (iv).
  - 10. The method according to any one of claims 1 to 9 wherein the ovarian cancer that is diagnosed is an epithelial ovarian cancer.
  - 11. The method according to any one of claims 1 to 9 wherein the ovarian cancer that is diagnosed is selected from the group consisting of serous ovarian cancer, non-invasive ovarian cancer, mixed phenotype ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer, clear cell ovarian cancer, papillary serous ovarian cancer, Brenner cell and undifferentiated adenocarcinoma.
  - 12. The method according to claim 11 wherein the ovarian cancer that is diagnosed is selected from the group consisting of serous ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer and clear cell ovarian cancer.
  - 13. A method of diagnosing a serous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein a modified level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a serous ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
  - (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession

10

25

- Number selected from the group consisting of: U62801, D49441, X51630, And AB018305;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441; X51630, And AB018305;
- (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441, X51630, And AB018305; and
- (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- 14. A method of diagnosing a mucinous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a mucinous ovarian cancer, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
  - a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, AI073913, AI928445, NM\_022454, W40460, AA132961 and AF111856;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, AI073913, AI928445, NM\_022454, W40460, AA132961 and AF111856;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
- (iv) a sequence that encodes a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of:
   NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, Al073913, Al928445, NM\_022454, W40460, AA132961 and AF111856; and

15

25

. 30

- (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- 15. The method of claim 14 wherein the nucleic acid probe comprises a sequence selected from the group consisting of:
  - (i) a sequence comprising at least about 20 contiguous nucleotides from SEQ ID NO:57 or 59 or 61;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from SEQ ID NO: 57 or 59 or 61;
- 10 (iii) a sequence that is at least about 80% identical to SEQ ID NO: 57 or 59 or 61;
  - (iv) a sequence that encodes the amino acid sequence set forth in SEQ ID NO: 58 or 60 or 62; and
  - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
  - 16. The method according to any one of claims 1 to 15 comprising performing a PCR reaction.
- 17. The method according to any one of claims 1 to 16 comprising performing a nucleic acid hybridization.
  - 18. A method of detecting an ovarian cancer-associated polypeptide in a biological sample the method comprising contacting the biological sample with an antibody that binds specifically to an ovarian cancer-associated polypeptide in the biological sample, the polypeptide being encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3.
  - 19. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a modified level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence

selected from the group consisting of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.

20. The method of claim 19 wherein the antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 6, 8, 10, 12, 14, 16, 18, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 47, 49, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.

10

15

20

25

- 21. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 1 or 2 other than a nucleic acid having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, Al745249 and Al694200.
- 22. The method of claim 21 wherein the antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 47, 49, 51, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.
- 23. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a reduced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has an ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous

WO 2004/022778 PCT/AU2003/001166

amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of NM\_022117, NM\_005460, NM\_002387, AI745249 and AI694200.

- The method of claim 23 wherein the antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, and 6.
- The method according to any one of claims 19 to 24 wherein the ovarian cancer that is diagnosed is an epithelial ovarian cancer.
  - 26. The method according to any one of claims 19 to 24 wherein the ovarian cancer that is diagnosed is selected from the group consisting of serous ovarian cancer, non-invasive ovarian cancer, mixed phenotype ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer, clear cell ovarian cancer, papillary serous ovarian cancer, Brenner cell and undifferentiated adenocarcinoma.

15

20

25

- 27. The method according to claim 26 wherein the ovarian cancer that is diagnosed is selected from the group consisting of serous ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer and clear cell ovarian cancer.
- 28. A method of diagnosing a serous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigenantibody complex to form and then detecting the complex wherein a modified level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a serous ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 2 or as set forth in Table 1 and having an Accession Number selected from the group consisting of: U62801, D49441, X51630, And AB018305.
- 35 29. A method of diagnosing a mucinous ovarian cancer in a human or animal subject being tested said method comprising contacting a biological sample from said subject

10

15

20

30

35

being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein a reduced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a mucinous ovarian cancer, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a polypeptide encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_006149, AA315933, U47732, NM\_005588, AW503395, NM\_004063, AI073913, AI928445, NM\_022454, W40460, AA132961 and AF111856.

- 30. The method according to claim 29 wherein the antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to SEQ ID NO: 58 or 60 or 62.
- 31. A method of detecting an ovarian cancer-associated antibody in a biological sample the method comprising contacting the biological sample with a polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3, wherein the polypeptide specifically binds to the ovarian cancer-associated antibody.
- 32. The method according to any one of claims 1 to 31 wherein the biological sample is contacted with a plurality of nucleic acid probes and/or antibodies and/or polypeptides.
- 25 33. The method according to any one of claims 1 to 32 wherein the subject being tested is a patient undergoing a therapeutic regimen to treat ovarian cancer.
  - 34. The method according to any one of claims 1 to 32 wherein the subject being tested is a subject suspected of having ovarian cancer.
  - 35. A method of monitoring the efficacy of a therapeutic treatment of ovarian cancer, the method comprising:
    - (i) providing a biological sample from a patient undergoing the therapeutic treatment; and
  - (ii) determining the level of a ovarian cancer-associated transcript in the biological sample by contacting the biological sample with a polynucleotide

30

that selectively hybridizes to a sequence having at least about 80% identity to a sequence as shown in any one of Tables 1-3, thereby monitoring the efficacy of the therapy.

- 5 36. The method according to claim 35 further comprising comparing the level of the ovarian cancer-associated transcript to a level of the ovarian cancer-associated transcript in a biological sample from the patient prior to, or earlier in, the therapeutic treatment.
- 37. A method of monitoring the efficacy of a therapeutic treatment of ovarian cancer, the method comprising :
  - (i) providing a biological sample from a patient undergoing the therapeutic treatment; and
  - (ii) determining the level of a ovarian cancer-associated antibody in the biological sample by contacting the biological sample with a polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3, wherein the polypeptide specifically binds to the ovarian cancer-associated antibody, thereby monitoring the efficacy of the therapy.
- 38. The method of claim 37 further comprising comparing the level of the ovarian cancer-associated antibody to a level of the ovarian cancer-associated antibody in a biological sample from the patient prior to, or earlier in, the therapeutic treatment.
- 39. A method of monitoring the efficacy of a therapeutic treatment of ovarian cancer, the method comprising :
  - (i) providing a biological sample from a patient undergoing the therapeutic treatment; and
  - (ii) determining the level of a ovarian cancer-associated polypeptide in the biological sample by contacting the biological sample with an antibody, wherein the antibody specifically binds to a polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3, thereby monitoring the efficacy of the therapy.

20

25

30

- 40. The method of claim 39 further comprising comparing the level of the ovarian cancer-associated polypeptide to a level of the ovarian cancer-associated polypeptide in a biological sample from the patient prior to, or earlier in, the therapeutic treatment.
- 5 41. The method according to any one of claims 35 to 40 wherein the ovarian cancer that is diagnosed is an epithelial ovarian cancer.
  - 42. The method according to any one of claims 35 to 41 wherein the ovarian cancer that is diagnosed is selected from the group consisting of serous ovarian cancer, non-invasive ovarian cancer, mixed phenotype ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer, clear cell ovarian cancer, papillary serous ovarian cancer, Brenner cell and undifferentiated adenocarcinoma.
- 43. The method according to claim 42 wherein the ovarian cancer that is diagnosed is selected from the group consisting of serous ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer and clear cell ovarian cancer.
  - 44. A method of determining the likelihood of survival of a subject suffering from an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
  - (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, Al970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, Al796870, X02761, AW968613, AW972565, AF045229, AW953853, U52426, F06700, Al798863, H52761, BE546947, AU076643, U20536, AA581602, AJ245210, X65965, Al806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135, BE466173, Al354722, M90464, AA829286, Al333771, BE465867, NM\_014992,

. 5

10

15

20

BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM\_002776, BE261944, NM\_006379, Al002238, X81789, NM\_002122, AB001914, AA311919, Al381750, AA292998, BE439580, Al677897, N72403, BE003054, AL035588, Al080491, AW770994, H24177, AF146761, NM\_001955, Al680737, Al752666, AA505445, BE246649, and NM\_003955;

- a sequence that hybridizes under at least low stringency hybridization conditions (ii) to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, AI970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, AI796870, X02761, AW968613, AW972565, AF045229, AW953853, U52426, F06700, Al798863, H52761, BE546947, AU076643, U20536, AA581602, AJ245210, X65965, Al806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135, BE466173, Al354722, M90464, AA829286, AI333771, BE465867, NM\_014992, BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM 002776, BE261944, NM\_006379, Al002238, X81789, NM\_002122, AB001914, AA311919, Al381750, AA292998, BE439580, Al677897, N72403, BE003054, AL035588, Al080491, AW770994, H24177, AF146761, NM\_001955, Al680737, Al752666, AA505445, BE246649, and NM\_003955;
- (iii) a sequence that is at least about 80% identical to (i) or (ii);
- a sequence that encodes a polypeptide encoded by a nucleic acid set forth in (iv) Table 1 and having an Accession Number selected from the group consisting of: 25 NM\_003014, AA046217, NM\_015902, T83882, AB040888, AA628980, Al623351, AW614420, AA243499, AF251237, AI970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, AI796870, X02761, AW968613, AW972565, AF045229, AW953853, U52426, F06700, AI798863, 30 H52761, BE546947, AU076643, U20536, AA581602, AJ245210, X65965, Al806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135, BE466173, Al354722, M90464, AA829286, AI333771, BE465867, NM\_014992, BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, 35 NM\_002776, BE261944, NM\_006379, Al002238, X81789, NM\_002122,

10

15

AB001914, AA311919, Al381750, AA292998, BE439580, Al677897, N72403, BE003054, AL035588, Al080491, AW770994, H24177, AF146761, NM\_001955, Al680737, Al752666, AA505445, BE246649, and NM\_003955; and

- (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iii) or (iv).
- 45. The method of claim 44 wherein the nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 63, 65, 67, 69, 71, and 73;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a sequence selected from the group consisting of SEQ ID NOs: 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iii) a sequence that is at least about 80% identical to a sequence selected from the group consisting of SEQ ID NOs: 63, 65, 67, 69, 71, 73, 75, 77, 79, 81 and 83;
- (iv) a sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84; and
- (v) a sequence that is complementary to (i) or (ii) or (iii) or (iv).
- 20 46. A method of determining the likelihood of survival of a subject suffering from an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigenantibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the 25 antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has has a poor probability of survival, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: NM 003014, AA046217, NM 015902, T83882, AB040888, AA628980, Al623351, 30 AW614420, AA243499, AF251237, AI970797, AF145713, X78565, T97307, BE243845, AW068302, AL133561, BE313555, X07820, Al973016, AF084545, U41518, Z11894, AW138190, BE086548, W47196, Al796870, X02761, AW968613, AW972565, AF045229, AW953853, U52426, F06700, AI798863, H52761, BE546947, AU076643, 35 U20536, AA581602, AJ245210, X65965, Al806770, BE386490, AW581992, U77534, AL034417, L10343, AW518944, W28729, Al640160, U11862, AW295980, X59135,

WO 2004/022778 PCT/AU2003/001166

195

BE466173, Al354722, M90464, AA829286, Al333771, BE465867, NM\_014992, BE616902, AA430373, R27430, BE387335, AW264102, AW952323, AA088177, BE614567, AL079658, NM\_002776, BE261944, NM\_006379, Al002238, X81789, NM\_002122, AB001914, AA311919, Al381750, AA292998, BE439580, Al677897, N72403, BE003054, AL035588, Al080491, AW770994, H24177, AF146761, NM 001955, Al680737, Al752666, AA505445, BE246649, and NM 003955.

5

10

15

20

25

- 47. The method of claim 46 wherein the antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence having at least about 80% identity to a sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, 78, 80, 82 and 84.
- 48. A method of determining the likelihood of survival of a subject suffering from a serous ovarian cancer, said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization obtained for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:
- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 71 or 73;
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 71 or 73;
- (iii) a sequence that is at least about 80% identical to (i) or (ii) and encoding an sFRP protein or a SOCS3 protein;
- (iv) a sequence that encodes a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 72 or 74; and
- 30 (v) a sequence that is complementary to any one of the sequences set forth in (i) or (ii) or (iv).
  - 49. A method of determining the likelihood of survival of a subject suffering from a serous ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced

WO 2004/022778

5

10

· 15

25

30

35

level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein said antibody binds to an sFRP polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 72 or a SOCS3 polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 74.

- 50. A method of determining the likelihood of survival of a subject suffering from a serous ovarian cancer, said method comprising contacting a biological sample from said subject being tested with at least two antibodies for a time and under conditions sufficient for antigen-antibody complexes to form and then detecting the complexes wherein an enhanced level of the antigen-antibody complexes for the subject being tested compared to the amount of the antigen-antibody complexes formed for a control subject not having ovarian cancer indicates that the subject being tested has a poor probability of survival, and wherein one antibody binds to an sFRP polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 72 and wherein one antibody binds to a SOCS3 polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 74.
- 51. The method according to any one of claims 44 to 47 wherein the ovarian cancer is an epithelial ovarian cancer.
  - 52. The method according to any one of claims 44 to 47 wherein the ovarian cancer is selected from the group consisting of serous ovarian cancer, non-invasive ovarian cancer, mixed phenotype ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer, clear cell ovarian cancer, papillary serous ovarian cancer, Brenner cell and undifferentiated adenocarcinoma.
  - 53. The method according to claim 52 wherein the ovarian cancer is selected from the group consisting of serous ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer and clear cell ovarian cancer.
  - 54. A method of determining the likelihood that a subject will suffer from a recurrence of an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with a nucleic acid probe for a time and under conditions sufficient for hybridization to occur and then detecting the hybridization wherein an elevated level of hybridization of the probe for the subject being tested compared to the hybridization

10

15

20

25

obtained for a control subject not having ovarian cancer indicates that the subject being tested has a high probability of recurrence, and wherein said nucleic acid probe comprises a sequence selected from the group consisting of:

- (i) a sequence comprising at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694. L08239, BE397649, NM\_012317, NM\_000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM\_002514, Al863735, NM 005397, W26391, H15474, U51166, AA243499, AW408807, AI738719, AB040888, BE313077, AI677897, C14898, AI821730, AF007393, H65423, N46243, AA095971, U20350, NM 005756, D19589. AW957446, AW294647, BE159718, Al888490, AA022569, BE147740, Al798863. BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM\_005512, AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763, AK000596, AI970797, AW519204, Z42387, AF145713, AA972412, AK001564, AW959861, BE313555, W25005, AI193356, AF111106, AI130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, Al497778, Al745379, U51712, AW375974, AF251237, NM\_000636, AA130986, AA216363, AA628980, AA811657, AA897108, AB040888, AF212225, Al089575, Al282028, Al368826, Al718702, AI827248, AK002039. AL109791, AW090198, AW296454. AW445034, AW452948, AW470411, AW885727, AW970859, AW979189, BE165866. BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965. X76732 and Z45051,
- (ii) a sequence that hybridizes under at least low stringency hybridization conditions to at least about 20 contiguous nucleotides from a nucleic acid set forth in Table 1
  30 and having an Accession Number selected from the group consisting of: M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694, L08239, BE397649, NM\_012317, NM\_000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM\_002514, Al863735, NM\_005397, W26391, H15474, U51166, AA243499, AW408807, Al738719, AB040888, BE313077, Al677897, C14898, Al821730, AF007393,

10

15

H65423, N46243, AA095971, U20350, NM\_005756, D19589, AW957446, AW294647, BE159718, Al888490, AA022569, BE147740, Al798863, BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM\_005512, AW953853. AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763. AK000596, AI970797, AW519204, Z42387, AF145713, AA972412, AK001564, AW959861, BE313555, W25005, Al193356, AF111106, Al130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, AI497778, AI745379, U51712, AW375974, AF251237. NM\_000636, AA130986, AA216363, AA628980, AA811657, AA897108, AB040888, AF212225, Al089575, Al282028, Al368826, Al718702, Al827248, AK002039, AL109791, AW090198, AW296454, AW445034, AW452948, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM 000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051;

- (iii) a sequence that is at least about 80% identical to (i) or (ii);
- a sequence that encodes a polypeptide encoded by a nucleic acid set forth in (iv) Table 1 and having an Accession Number selected from the group consisting 20 of:M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694, L08239, BE397649, NM 012317, NM 000947, AJ250562, AL040183, BE207573, BE564162, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM 002514, Al863735, NM 005397, W26391, H15474, U51166, AA243499, AW408807, AI738719, AB040888, BE313077, AI677897, C14898, AI821730, 25 AF007393, H65423, N46243, AA095971, U20350, NM\_005756, D19589, AW957446, AW294647, BE159718, AI888490, AA022569, BE147740, AI798863, BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM\_005512, AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763, AK000596, AI970797, AW519204, Z42387, AF145713, AA972412, 30 AK001564, AW959861, BE313555, W25005, Al193356, AF111106, Al130740, AA985190, BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, Al497778, Al745379, U51712, AW375974, NM\_000636, AA130986, AA216363, AA628980, AA811657, AF251237. AA897108, AB040888, AF212225, Al089575, Al282028, Al368826, Al718702, 35 AK002039, AL109791, AW090198, AW296454, AW445034, Al827248.

10

15

20

25

30

35

AW452948, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580, BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051; and

- (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- 55. The method of claim 54 determining the likelihood that a subject will suffer from a recurrence of an ovarian cancer, said method comprising contacting a biological sample from said subject being tested with an antibody for a time and under conditions sufficient for an antigen-antibody complex to form and then detecting the complex wherein an enhanced level of the antigen-antibody complex for the subject being tested compared to the amount of the antigen-antibody complex formed for a control subject not having ovarian cancer indicates that the subject being tested has a high probability of recurrence, and wherein said antibody binds to a polypeptide comprising an amino acid sequence comprising at least about 10 contiguous amino acid residues of a sequence encoded by a nucleic acid set forth in Table 1 and having an Accession Number selected from the group consisting of: M86849, AW963419, BE298665, AK000637, BE077546, T97307, R24601, BE090176, AA393907, W28729, BE313754, AW673081, AA356694. L08239, BE397649, NM 012317, NM 000947, AJ250562, AL040183, BE207573, BE439580, AW067800, AA569756, AW138190, AF126245, L10343, NM\_002514, Al863735, NM\_005397, W26391, H15474, U51166. AA243499, Al738719, AB040888, BE313077, Al677897, C14898, Al821730, AF007393, H65423, N46243, AA095971, U20350, NM 005756, D19589, AW957446, AW294647, BE159718, Al888490, AA022569, BE147740, Al798863, BE464341, AL080235, Al557212, X75208, AA628980, BE242587, NM\_005512, AW953853, AU076611, AW968613, AL353944, BE614149, AA292998, H12912, AA188763, AK000596, Al970797, AW519204, Z42387, AF145713, AA972412, AK001564, AW959861, BE313555. W25005, Al193356, AF111106, Al130740, AA985190. BE221880, AF084545, R26584, AW247380, AA364261, U25849, AF262992, AW342140, AL133572, Al497778, Al745379, U51712, AW375974, AF251237, NM\_000636, AA130986, AA216363, AA628980, AA811657, AA897108, AB040888, AF212225, Al089575, Al282028, Al368826, Al718702, Al827248, AK002039, AL109791, AW090198, AW296454, AW445034, AW452948, AW470411, AW885727, AW970859, AW979189, BE165866, BE175582, BE242587, BE271927, BE439580,

BE464016, D63216, F34856, M83822, N33937, N49068, N51357, N80486, NM\_000954, NM\_005756, NM\_016652, R26584, R31178, W05391, W25005, W45393, W68815, X65965, X76732 and Z45051.

- 5 56. The method according to claim 54 or 55 wherein the ovarian cancer is an epithelial ovarian cancer.
  - 57. The method according to any one of claims 54 to 56 wherein the ovarian cancer is selected from the group consisting of serous ovarian cancer, non-invasive ovarian cancer, mixed phenotype ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer, clear cell ovarian cancer, papillary serous ovarian cancer, Brenner cell and undifferentiated adenocarcinoma.

10

20

25

30

35

- 58. The method according to claim 57 wherein the ovarian cancer is selected from the group consisting of serous ovarian cancer, mucinous ovarian cancer, endometrioid ovarian cancer and clear cell ovarian cancer.
  - 59. The method according to any one of claims 35 to 58 wherein the biological sample is contacted with a plurality of nucleic acid probes and/or antibodies and/or polypeptides.

60. A method for identifying a compound that modulates an ovarian cancerassociated polypeptide, the method comprising:

- (i) contacting the compound with a ovarian cancer-associated polypeptide, the polypeptide encoded by a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3; and
- (ii) determining the functional effect of the compound upon the polypeptide.
- 61. A method for determining a candidate compound for the treatment of ovarian cancer comprising:
- (i) administering a test compound to a mammal having ovarian cancer or a cell isolated therefrom;
  - (ii) comparing the level of gene expression of a polynucleotide that selectively hybridizes to a sequence at least 80% identical to a sequence as shown in Tables 1-3 in a treated cell or mammal with the level of gene expression of the polynucleotide in a control cell or mammal, wherein a test compound that

modulates the level of expression of the polynucleotide is a candidate for the treatment of ovarian cancer.

- 62. An assay device for use in the diagnosis or prognosis of ovarian cancer, said device comprising a plurality of polynucleotides immobilized to a solid phase, wherein each of said polnucleotides consists of a gene as listed in any one of Tables 1-3.
  - 63. The device of claim 62 consisting of a substantially planar chip.
- 10 64. An assay device for use in the diagnosis or prognosis of ovarian cancer, said device comprising a plurality of different antibodies immobilized to a solid phase, wherein each of said antibodies binds to a polypeptide listed in Tables 1-3.
  - 65. The device of claim 64 consisting of a substantially planar chip.

15

30

- 66. Use of a polynucleotide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.
- 20 67. Use of a vector comprising a polynucleotide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.
- 68. Use of an isolated polypeptide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.
  - 69. Use of an antibody that binds to an isolated polypeptide as set forth in any one of Tables 1-3 in the diagnosis or prognosis of ovarian cancer or for the preparation of a medicament for the treatment of ovarian cancer.
  - 70. A method of diagnosing an ovarian cancer in a human or animal subject being tested said method comprising determining aberrant methylation in a promoter sequence that regulates expression of a tumor suppressor gene in a biological sample from said subject compared to the methylation of the promoter in nucleic acid obtained for a control

subject not having ovarian cancer wherein said aberrant methylation indicates that the subject being tested has an ovarian ovarian cancer.

- 71. The method of claim 70 wherein hypermethylation of the promoter sequence is determined.
  - 72. The method of claim 70 or 71 wherein the methylation is determined in the promoter region that regulates expression of an MCC gene comprising a sequence selected from the group consisting of:
- 10 (i) the nucleotide sequence set forth as SEQ ID NO: 3;
  - (ii) a sequence that hybridizes under at least low stringency hybridization conditions to the nucleotide sequence set forth as SEQ ID NO: 3;
  - (iii) a sequence that is at least about 80% identical to (i) or (ii);
  - (iv) a sequence that encodes a polypeptide encoded by a nucleotide sequence set forth as SEQ ID NO: 3; and
    - (v) a sequence that is complementary to any one of the sequences set forth in (i) or(ii) or (iii) or (iv).
- 73. The method according to any one of claims 70 to 72 wherein the ovarian cancer that is diagnosed is an epithelial ovarian cancer.




FIGURE 1




FIGURE 2

WO 2004/022778 PCT/AU2003/001166




FIGURE 3




FIGURE 4a




FIGURE 4b

## SEQUENCE LISTING

<110>	Garv	an II	nstit	tute	of N	1edio	cal I	Résea	arch						
<120>	Meth	ods i	Eor t	he o	liagn	nosis	s and	i pro	ognos	sis (	o£ o	varia	an ca	ancer	
<130>	5017	31/MI	RO					•							
<150> <151>	AU20 2002														
<160>	84														
<170>	PatentIn version 3.1														
<210> <211> <212> <213>	3432 DNA														,
<220> <221> CDS <222> (94)(2850) <223>															
<400> 1 gtcagtccct tcgcgctcct gagccgccgg cgcgccgggc gccccgggaa tttataagta 60														60	
	ccgta (														114
		٠.	,	, ,										r Leu	
	tg gat eu Asp 10														162
	tc aag eu Lys 5														210
	ac aga sp Arg														258
	tt aca le Thr														306
aag t Lys Pl	tc cgc he Arg	cca Pro 75	gtg Val	aag Lys	cgg Arg	gtt Val	tcg Ser 80	cca Pro	ctg Leu	aaa Lys	cat His	cag Gln 85	cca Pro	gag Glu	354
act c	tg gag eu Glu 90	aac Asn	aat Asn	gaa Glu	agt Ser	gat Asp 95	gac Asp	caa Gln	aag Lys	aac Asn	cag Gln 100	aaa Lys	gtg Val	gtt Val	402
Glu T	ac cag yr Gln 05														450
ctt go Leu G 120	gc cct ly Pro	gga Gly	gat Asp	gga Gly 125	gtg Val	ggc Gly	ggc Gly	cca Pro	cca Pro 130	ggt Gly	aag Lys	agc Ser	tct Ser	gag Glu 135	498
ccc aq Pro Se	gc aca er Thr	tcg Ser	ctg Leu 140	ggt Gly	gaa Glu	ctg Leu	gag Glu	cac His 145	tac Tyr	gac Asp	ctc Leu	gac Asp	atg Met 150	gat Asp	546

•	ga Gl	g at u Il	t ci e Le	eu As	at gt sp Va 55	g co il Pr	t ta o Ty	ıt at r Il	t aa e Ly 16	s Se	c ag r Se	t ca r Gl	g ca n Gl	g ct n Le 16	u Āl	c tct a Ser	594
	tt Ph	t ac e Th	rЬ	ag gt ys Va 70	g ac	t to r Se	a ga r Gl	a aa u Ly 17	s Ar	a at g Il	t tt e Le	g gg u Gl	c tt y Le 18	u Cy	c ac	a acc r Thr	642
٠	at Il	c aa e As 18	n Gl	jc ct ly L∈	t to u Se	t gg r Gl	с аа у Lу 19	s Al	c tg a Cy	c tc s Se	t ac	a gg r Gl 19	y Se	t to r Se	t ga r Gl	g ago u Ser	690
	tc: Se: 200	c Se	a to r Se	c aa er As	c at n Me	g gc t Al 20	a Pr	a tt o Ph	t tgʻ e Cy:	t gti s Val	t ct l Le	u Se	t cc r Pr	c gt o Va	g aa l Ly	a agc s Ser 215	
	cct Pro	ca Hi	c tt s Le	g ag u Ar	a aa g Ly 22	s Al	a tc a Se	a gc r Al	t gto a Vai	225	e Hi	c ga s As	c ca p Gl:	g ca n Hi	c aas s Ly: 23	g ctg s Leu 0	786
	Ser	Thi	t ga r Gl	a ga u G1 23	u Th	c ga r Gl	g ato u Ilo	c tc. e Se:	a cct r Pro 240	Pro	cto Lei	g gti 1 Va.	t aaa l Lya	a tg s Cy: 24:	s Gl	c tct y Ser	834
	gca Ala	tai Tyi	ga G1 25	u Pr	t gaa o Gl	a aad u Asi	c caq n Gl:	g agi n Se: 25!	r Lys	gad Asp	tto Phe	cta Lei	a aad a Asi 260	ı Ly	g aca	ttt Phe	882
•	agt Ser	gat Asp 265	Pr	t ca o Hi	t ggi s Gl	t cga y Arg	a aaa g Lys 270	s Val	gag L Glu	ı aag	aca Thr	aca Thi	: Pro	a gad Asp	c tgo Cys	cag Gln	930
	ctc Leu 280	Arg	gc Al	c tto a Pho	c cad e His	cta Leu 285	ı Glr	tco Sei	c tca Ser	gca Ala	gca Ala 290	Gli	tco Ser	aaa Lys	a cca Pro	gaa Glu 295	978
	gag Glu	cag Gln	gt: Va:	c agi	300 300	, Leu	aac Asn	cgg Arg	acc Thr	agc Ser 305	Ser	caa Gln	ggc Gly	e cca	gaa Glu 310	gaa Glu	1026
	agg Arg	agt Ser	gaq Glu	y tat ı Tyı 315	: Leu	aaa Lys	aaa Lys	gtg Val	aaa Lys 320	agc Ser	atc Ile	ttg Leu	aac Asn	att Ile 325	Val	aaa Lys	1074
	gaa Glu	gga Gly	caç Glr 330	) ITE	tct Ser	ctc Leu	ctg Leu	Pro 335	cac His	cta Leu	gct Ala	gca Ala	gac Asp 340	Asn	cta Leu	gac Asp	1122
	aaa Lys	att Ile 345	Cac	gac Asp	gaa Glu	aat Asn	gga Gly 350	aac Asn	aat Asn	cta Leu	tta Leu	cat His 355	att Ile	gcg Ala	gcg Ala	tca Ser	1170
	cag Gln 360	gga Gly	Cac	gca Ala	gag Glu	tgt Cys 365	cta Leu	cag Gln	cac His	ctc Leu	act Thr 370	tct Ser	ttg Leu	atg Met	gga Gly	gaa Glu 375	1218
	gac Asp	tgc Cys	ctc	aat Asn	gag Glu 380	cgc Arg	aac Asn	act Thr	gag Glu	aag Lys 385	ttg Leu	act Thr	cca Pro	gca Ala	ggc Gly 390	ctg Leu	1266
•	gcc Ala	att Ile	aag Lys	aat Asn 395	ggt Gly	cag Gln	ttg Leu	gag Glu	tgc Cys 400	gta Val	cgc Arg	tgg Trp	atg Met	gtg Val 405	agc Ser	gaa Glu	1314
	aca Thr	gaa Glu	gcc Ala 410	att Ile	gca Ala	gaa Glu	ctg Leu	agt Ser 415	tgt Cys	tct Ser	aag Lys	gat Asp	ttt Phe 420	cca Pro	agc Ser	ctt Leu	1362

						tat Tyr 430										1410
	Gln		_		-	cag Gln				_	_	_	_	-	_	1458
						cac His										1506
						gaa Glu										1554
						tcc Ser										1602
						gtg Val 510										1650
						acc Thr										1698
		-	_			caa Gln						_	_	-		1746
						cct Pro									cct Pro.	1794
						tgg Trp										1842
						gga Gly 590	_					_	_			1890
agc Ser 600	ctg Leu	tca Ser	gcc Ala	tcc Ser	agc Ser 605	cgg Arg	gct Ala	aga Arg	ccc Pro	aaa Lys 610	gca Ala	aaa Lys	gat Asp	gaa Glu	gat Asp 615	1938
					_	cag Gln		_		_				_		1986
						ctg Leu										2034
						aag Lys										2082
						ctg Leu 670	_	-			_	_			-	2130
	_					gag Glu	-						-		-	2178

			cca Pro													2226
gac Asp	agc Ser	gca Ala	gaa Glu 715	agc Ser	ctg Leu	cac His	ctg Leu	atg Met 720	att Ile	aag Lys	aaa Lys	cac His	acc Thr 725	ttg Leu	gca Ala	2274
tca Ser	G] À aaa	gga Gly 730	cgc Arg	agg Arg	ttt Phe	cct Pro	ttc Phe 735	agc Ser	atc Ile	aag Lys	gcc Ala	tcc Ser 740	aaa Lys	tcc Ser	ctg Leu	2322
gat Asp	ggc Gly 745	cac His	agc Ser	cca Pro	tct Ser	ccc Pro 750	acc Thr	tca Ser	gag Glu	agc Ser	agc Ser 755	gaa Glu	cca Pro	gac Asp	tta Leu	2370
			tat Tyr													2418
			cag Gln													2466
gcc Ala	aca Thr	agt Ser	ccc Pro 795	aag Lys	agt Ser	gcc Ala	ctc Leu	aag Lys 800	tct Ser	cca	tct Ser	tcc Ser	aag Lys 805	cgt Arg	agg Arg	2514
aca Thr	tct Ser	cag Gln 810	aac Asn	tta Leu	aaa Lys	ctg Leu	aga Arg 815	gtt Val	acc Thr	ttt Phe	gag Glu	gag Glu 820	cct Pro	gtg Val	gtg Val	2562
cag Gln	atg Met 825	gag Glu	cag Gln	cct Pro	agc Ser	ctt Leu 830	gaa Glu	ctg Leu	aat Asn	gga Gly	gaa Glu 835	aaa Lys	gac Asp	aaa Lys	gat Asp	2610
aag Lys 840	ggc Gly	agg Arg	act Thr	ctc Leu	cag Gln 845	cgg Arg	acc Thr	tcc Ser	aca Thr	agt Ser 850	aac Asn	gaa Glu	tcg Ser	Gly ggg	gat Asp 855	2658
caa Gln	ctg Leu	aaa Lys	agg Arg	cct Pro 860	ttt Phe	gga Gly	gcc Ala	Phe	cga Arg 865	tct Ser	atc Ile	atg Met	gag Glu	aca Thr 870	cța Leu	2706
agt Ser	ggc Gly	aac Asn	caa Gln 875	aac Asn	aat Asn	aat Asn	aat Asn	aac Asn 880	tac Tyr	cag Gln	gca Ala	gcc Ala	aac Asn 885	cag Gln	ctg Leu	2754
aaa Lys	Thr	tct Ser 890	aca Thr	ttg Leu	ccc Pro	Leu	acc Thr 895	tca Ser	ctt Leu	GJĀ āāā	agg Arg	aag Lys 900	aca Thr	gat Asp	gcc Ala	2802
aag Lys	gga Gly 905	aac Asn	cct Pro	gcc Ala	agc Ser	tcc Ser 910	gct Ala	agc Ser	aaa Lys	gga Gly	aag Lys 915	aat Asn	aag Lys	gca Ala	gca Ala	2850
taat	gaca	tc a	atag	aaaa	a tg	aaga	aatc	cta	cage	ata	aagc	acat	tg c	tgag	ccaga	2910
gtca	aaag	aa c	tctt	cttg	t aa	atca	cttt	tta	aatt	ttc	tctc	actg	at g	ccct	ttgga	2970
aatt	attg	ga a	attt	ctgg	a ct	atcc	tctt	tgg	aaag	aga	acça	tgaa	aa c	aatg	cctca	3030
ccag	caga	ag a	acag	aata	t ca	ggat	gcct	taa	attt	ata	gtag	taga	ct g	taaa	agatt	3090
catt	ttgg	gg t	gata	tctg	t at	atat	aact	tgt	tttt	tta	aaag	atgc	cg t	ttaa	aagca	3150
tgati	tggg	aa a	atgt	acgt	t tt	ttaa	gagt	aga	ttga	ttc	accc	tacc	ca c	agga	cattc	3210

<210> 2

<211> 919

<212> PRT

<213> NM\_005460 SNCAIP, synphilin

<400> 2

Met Glu Ala Pro Glu Tyr Leu Asp Leu Asp Glu Ile Asp Phe Ser Asp 1 5 10 15

Asp Ile Ser Tyr Ser Val Thr Ser Leu Lys Thr Ile Pro Glu Leu Cys 20 25 30

Arg Arg Cys Asp Thr Gln Asn Glu Asp Arg Ser Ala Ser Ser Ser Ser 35 40 45

Trp Asn Cys Gly Ile Ser Thr Leu Ile Thr Asn Thr Gln Lys Pro Thr 50 55 60

Gly Ile Ala Asp Val Tyr Ser Lys Phe Arg Pro Val Lys Arg Val Ser 65 70 75 80

Pro Leu Lys His Gln Pro Glu Thr Leu Glu Asn Asn Glu Ser Asp Asp 85 90 95

Gln Lys Asn Gln Lys Val Val Glu Tyr Gln Lys Gly Glu Ser Asp 100 105 110

Leu Gly Pro Gln Pro Gln Glu Leu Gly Pro Gly Asp Gly Val Gly Gly 115 120 125

Pro Pro Gly Lys Ser Ser Glu Pro Ser Thr Ser Leu Gly Glu Leu Glu 130 135 140

His Tyr Asp Leu Asp Met Asp Glu Ile Leu Asp Val Pro Tyr Ile Lys 145 150 155 160

Ser Ser Gln Gln Leu Ala Ser Phe Thr Lys Val Thr Ser Glu Lys Arg 165 170 175

Ile Leu Gly Leu Cys Thr Thr Ile Asn Gly Leu Ser Gly Lys Ala Cys 180 185 190 Ser Thr Gly Ser Ser Glu Ser Ser Ser Ser Asn Met Ala Pro Phe Cys 195 200 205

Val Leu Ser Pro Val Lys Ser Pro His Leu Arg Lys Ala Ser Ala Val 210 215 220

Ile His Asp Gln His Lys Leu Ser Thr Glu Glu Thr Glu Ile Ser Pro 225 230 235 240

Pro Leu Val Lys Cys Gly Ser Ala Tyr Glu Pro Glu Asn Gln Ser Lys 245 250 255

Asp Phe Leu Asn Lys Thr Phe Ser Asp Pro His Gly Arg Lys Val Glu 260 265 270

Lys Thr Thr Pro Asp Cys Gln Leu Arg Ala Phe His Leu Gln Ser Ser 275 280 285

Ala Ala Glu Ser Lys Pro Glu Glu Gln Val Ser Gly Leu Asn Arg Thr 290 295 300

Ser Ser Gln Gly Pro Glu Glu Arg Ser Glu Tyr Leu Lys Lys Val Lys 305 310 315 320

Ser Ile Leu Asn Ile Val Lys Glu Gly Gln Ile Ser Leu Leu Pro His 325 330 335

Leu Ala Ala Asp Asn Leu Asp Lys Ile His Asp Glu Asn Gly Asn Asn 340 345 350

Leu Leu His Ile Ala Ala Ser Gln Gly His Ala Glu Cys Leu Gln His 355 360 365

Leu Thr Ser Leu Met Gly Glu Asp Cys Leu Asn Glu Arg Asn Thr Glu 370 375 380

Lys Leu Thr Pro Ala Gly Leu Ala Ile Lys Asn Gly Gln Leu Glu Cys 385 390 395 400

Val Arg Trp Met Val Ser Glu Thr Glu Ala Ile Ala Glu Leu Ser Cys
405 410 415

Ser Lys Asp Phe Pro Ser Leu Ile His Tyr Ala Gly Cys Tyr Gly Gln 420 425 430

Glu Lys Ile Leu Leu Trp Leu Leu Gln Phe Met Gln Glu Gln Gly Ile 435 440 445

Ser Leu Asp Glu Val Asp Gln Asp Gly Asn Ser Ala Val His Val Ala 450 455 460

。 (1.24 · 1.25 · 1.15 · 1.45

Market V

Ser Gln His Gly Tyr Leu Gly Cys Ile Gln Thr Leu Val Glu Tyr Gly 465 475 480

in the section of the section of the contract of the contract

Ala Asn Val Thr Met Gln Asn His Ala Gly Glu Lys Pro Ser Gln Ser 485 490 495

Ala Glu Arg Gln Gly His Thr Leu Cys Ser Arg Tyr Leu Val Val
500 505 510

Glu Thr Cys Met Ser Leu Ala Ser Gln Val Val Lys Leu Thr Lys Gln 515 520 525

Leu Lys Glu Gln Thr Val Glu Arg Val Thr Leu Gln Asn Gln Leu Gln 530 540

Gln Phe Leu Glu Ala Gln Lys Ser Glu Gly Lys Ser Leu Pro Ser Ser 545 550 555 560

Pro Ser Ser Pro Ser Ser Pro Ala Ser Arg Lys Ser Gln Trp Lys Ser 565 570 575

Pro Asp Ala Asp Asp Ser Val Ala Lys Ser Lys Pro Gly Val Gln 580 585 590

Glu Gly Ile Gln Val Leu Gly Ser Leu Ser Ala Ser Ser Arg Ala Arg 595 600 605

Pro Lys Ala Lys Asp Glu Asp Ser Asp Lys Ile Leu Arg Gln Leu Leu 610 620

Gly Lys Glu Ile Ser Glu Asn Val Cys Thr Gln Glu Lys Leu Ser Leu 625 630 635 · 640

Glu Phe Gln Asp Ala Gln Ala Ser Ser Arg Asn Ser Lys Lys Ile Pro 645 650 655

Leu Glu Lys Arg Glu Leu Lys Leu Ala Arg Leu Arg Gln Leu Met Gln 660 665 670

Arg Ser Leu Ser Glu Ser Asp Thr Asp Ser Asn Asn Ser Glu Asp Pro 675 680 685

Lys Thr Thr Pro Val Arg Lys Ala Asp Arg Pro Arg Pro Gln Pro Ile 690 695 700

Val Glu Ser Val Glu Ser Met Asp Ser Ala Glu Ser Leu His Leu Met 705 710 715 720

Ile Lys Lys His Thr Leu Ala Ser Gly Gly Arg Arg Phe Pro Phe Ser 725 730 735

Ile Lys Ala Ser Lys Ser Leu Asp Gly His Ser Pro Ser Pro Thr Ser 740 745 750

Glu Ser Ser Glu Pro Asp Leu Glu Ser Gln Tyr Pro Gly Ser Gly Ser 755 760 765

Ile Pro Pro Asn Gln Pro Ser Gly Asp Pro Gln Gln Pro Ser Pro Asp 770 780

Ser Thr Ala Ala Gln Lys Val Ala Thr Ser Pro Lys Ser Ala Leu Lys 785 790 795 800

Ser Pro Ser Ser Lys Arg Arg Thr Ser Gln Asn Leu Lys Leu Arg Val 805 810 815

Thr Phe Glu Glu Pro Val Val Gln Met Glu Gln Pro Ser Leu Glu Leu 820 825 830

Asn Gly Glu Lys Asp Lys Asp Lys Gly Arg Thr Leu Gln Arg Thr Ser 835 840 845

Thr Ser Asn Glu Ser Gly Asp Gln Leu Lys Arg Pro Phe Gly Ala Phe 850 855 860

Arg Ser Ile Met Glu Thr Leu Ser Gly Asn Gln Asn Asn Asn Asn Asn 865 870 870 880

Tyr Gln Ala Ala Asn Gln Leu Lys Thr Ser Thr Leu Pro Leu Thr Ser 885 890 895

Leu Gly Arg Lys Thr Asp Ala Lys Gly Asn Pro Ala Ser Ser Ala Ser 900 905 910

Lys Gly Lys Asn Lys Ala Ala 915

<210> 3

<211> 4181

<212> DNA

<213> NM\_002387 MCC, mutated in colorectal cancers

<220>

<221> CDS

<222> (221)..(2707)

<223>

-1	00>	3														
			j caa	atggo	tcg	tccg	tgaa	ac g	gcgag	rccad	g go	etgct	cttt	: tta	agagt	gc 6(
ct	gcat	ccto	cgt	ttgc	gct	tcgc	aact	gt c	ctgg	gtga	a aa	tggc	tgto	: tag	actaa	aa 120
tg	tggc	agaa	a ggg	gacca	agc	agtg	gata	ıtt g	agco	tgtg	ga ag	tcca	acto	tta	agctc	cg 180
ag	acct	gggg	gac	tgag	agc	ccag	cțct	ga a	aạġt	gcat					a gtt y Val 5	235
gc Ala	c at a Me	g aa t Ly	a ta 's Ty	r Gl 10	y As	c ga n As	c to p Se	c tc r Se	g gc r Al 15	a Gl	g ct u Le	g ag u Se	t ga r Gl	g ct u Le 20	c cat u His	283
tc: Se:	a gc r Al	a go a Al	c ct a Le 25	u AL	a tc. a Se:	a ct. r Le	a aa u Ly	g gg s Gl 30	a ga y As	t at p Il	a gt e Va	g ga 1 Gl	a ct u Le 35	t aa u As:	t aaa n Lys	331
cgt Arg	t cta g Le	c ca u Gl 40	n GI	a ac n Th	a gaq r Gli	g ago	g ga g Gl 45	a cg	g ga g As	c ct p Le	t ct u Le	g ga u Gl 50	a aa u Ly:	g aa s Ly:	a ttg s Leu	379
gco	aaq Lys 55	g gc s Al	a ca a Gl	g tg n Cy	c gaq s Gli	g caq u Gl: 60	tc Se:	c cad	c cto s Le	c at u Me	g ag t Ar 65	a gad g Gli	g cat u Hi:	t gaq s Gli	g gat 1 Asp	427
gto Val 70	caq Glr	g ga n Gli	g cga	a ac	g acc Thr 75	g ctt Lei	: cg	c tat	t gaq	g gaa 1 Gli 80	a cgo	c ato	c aca	a gaq Glu	ctc Leu 85	475
Cac His	ago Ser	gto Val	c att	t gcg a Ala 90	g gag a Glu	cto Lev	aad Asr	c aag n Lys	g aag Lys 95	g ata	a gad e Ası	cgt Arg	cto J Lei	g caa Glr 100	ggc Gly	523
acc Thr	acc Thr	ato Ile	agg Arg 105	f GII	gaa Glu	gat Asp	gag Glu	tac Tyr 110	Ser	gaa Glu	t cto	cga Arg	tca Ser 115	: Glu	ctc Leu	571
agc Ser	cag Gln	Ser 120	GLE	cac His	gag Glu	gtc Val	aac Asn 125	Glu	gac Asp	tct Ser	. cga	ago Ser 130	Met	gac Asp	caa Gln	619
gac Asp	cag Gln 135	Tur	tct Ser	gtc Val	tct Ser	atc Ile 140	Pro	gaa Glu	aac Asn	cag Gln	tct Ser 145	Thr	atg Met	gtt Val	act Thr	667
gct Ala 150	gac Asp	atg Met	gac Asp	aac Asn	tgc Cys 155	agt Ser	gac Asp	ctg Leu	aac Asn	tca Ser 160	Glu	ctg Leu	cag Gln	agg Arg	gtg Val 165	715
ctg Leu	aca Thr	Gly	ctg Leu	gag Glu 170	aat Asn	gtt Val	gtc Val	tgc Cys	ggc Gly 175	agg Arg	aag Lys	aag Lys	agc Ser	agc Ser 180	tgc Cys	763
agc Ser	ctc Leu	tcc Ser	gtg Val 185	gcc Ala	gag Glu	gtg Val	gac Asp	agg Arg 190	cac His	att Ile	gag Glu	cag Gln	ctc Leu 195	acc Thr	aca Thr	811
gcc Ala	agc Ser	gag Glu 200	cac His	tgt Cys	gac Asp	ctg Leu	gct Ala 205	att Ile	aag Lys	aca Thr	gtc Val	gag Glu 210	gag Glu	att Ile	gag Glu	859
GTĀ	gtg Val 215	ctt Leu	ggc Gly	cgg Arg	gac Asp	ctg Leu 220	tat Tyr	ccc Pro	aac Asn	ctg Leu	gct Ala 225	gaa Glu	gag Glu	agg Arg	tct Ser	907

						gct Ala										955
						aaa Lys										1003
	_		_			gaa Glu			_						-	1051
						cta Leu										1099
_			_			tcc Ser 300			-	_				•		1147
		_	_		-	agc Ser	-	_		-				_	_	1195
						cta Leu										1243
	_			-	_	ggc Gly		_		_	_			-		1291
						ctg Leu										1339
						aca Thr 380										1387
						ttt Phe										1435
						tcc Ser					_					1483
						gct Ala										1531
						gcg Ala										1579
						gaa Glu 460										1627
						cag Gln										1675
cct Pro	gga Gly	gac Asp	cag Gln	tcg Ser 490	G1 Å aaa	gat Asp	gaa Glu	aac Asn	atc Ile 495	act Thr	cag Gln	atg Met	ctc Leu	aag Lys 500	cga Arg	1723

and the state of the contract of

gct Ala	cat His	gac Asp	tgc Cys 505	cgg Arg	aag Lys	aca Thr	gct Ala	gag Glu 510	aac Asn	gct Ala	gcc Ala	aag Lys	gcc Ala 515	ctg Leu	ctc Leu	1771
					agc Ser											1819
		_			gag Glu	_					_				_	. 1867
					agt Ser 555											1915
	_		_	_	gat Asp			_	_		_		_			1963
	_	-	_		atg Met	_	-	_	_	_				_		2011
					aag Lys											2059
					atg Met											2107
	Glu				cag Gln 635											2155
					agc Ser											2203
					ctg Leu	_						-	-	-	_	2251
					agc Ser						Ser					2299
					gat Asp											2347
			_	_	agc Ser 715				_	_	-	-				2395
					aag Lys											2443
					ctc Leu											2491
					aat Asn											2539

gct gcc tat gag aaa gca aag aaa aag cat caa aac aaa ctg aag aag Ala Ala Tyr Glu Lys Ala Lys Lys Lys His Gln Asn Lys Leu Lys Lys 775 780 785	2587
tta gag tcg cag atg atg gcc atg gtg gag aga cat gag acc caa gtg Leu Glu Ser Gln Met Met Ala Met Val Glu Arg His Glu Thr Gln Val 790 800 805	2635
agg atg ctc aag caa aga ata gct ctg cta gag gag gag aac tcc agg Arg Met Leu Lys Gln Arg Ile Ala Leu Leu Glu Glu Glu Asn Ser Arg 810 815 820	2683
cca cac acc aat gaa act tog ott taatcagcac toacgcaccg gagttotgcc Pro His Thr Asn Glu Thr Ser Leu 825	2737
catgggaagt aaactgcagc aggccactgg ggacagaagg gcccatgtac ttgttgggag	2797
gaggaggaaa gggaaggctg gcaggtaggt cggcacttgg acaatggagt gccccaactc	2857
aaccettggg gtgactggcc atggtgacat tgtggactgt atccagaggt gcccgctctt	2917
ccctcctggg cccacaacag cgtgtaaaca catgttctgt gcctgctcag cagagcctcg	2977
tttctgcttt cagcactcac tctccccctc ctcttctggt ctggcggctg tgcatcagtg	3037
ggatcccaga cattigittc igtaagatti tccattgtat cctctititg gtagatgctg	3097
ggctcatctt ctagaatctc gtttctcctc tttcctcctg cttcatggga aaacagacct	3157
gtgtgtgcct ccagcattta aaaggactgc tgatttgttt actacagcaa ggctttggtt	3217
tecaagteec gggteteaac tttaagatag aggeggeeat aagaggtgat etetgggagt	3277
tataggtcat gggaagagcg tagacaggtg ttacttacag tcccagatac actaaagtta	3337
caaacagacc accaccagga ctgtgcctga acaattttgt attgagagaa taaaaacttc	3397
cttcaatctt cattttggag gcagggctgg gaagggagcg ctctcttgat tctgggattt	3457
ctccctctca gtggagcctt attaatatcc aagacttaga gctgggaatc tttttgatac	3517
ctgtagtgga actaaaattc tgtcaggggt ttcttcaaga gctgagaaac attattagca	3577
cttcccgccc cagggcacta cataattgct gttctgctga atcaaatctc ttccacatgg	3637
gtgcatttgt agctctggac ctgtctctac ctaaggacaa gacactgagg agatactgaa	3697
cattttgcaa aacttatcac gcctacttaa gagtgctgtg taacccccag ttcaagactt	375.7
ageteetgtt gteatgaegg ggaeagagtg agggaatggt agttaagget tettititge	3817
ccccagatac atggtgatgg ttagcatatg gtgcttaaaa ggttaaattt caagcaaaat	3877
gcttacaggg ctaggcagta ccaaagtaac tgaattattt caggaaggtc ttcaatctta	3937
aaacaaatto attattottt ttoagtttta ootottotot otoagttota cactgataca	3997
cttgaaggac catttactgt ttttttctgt agcaccagag aatccatcca aagttcccta	4057
tgaaaaatgt gttccattgc catagctgac tacaaattaa agttgaggag gtttctgcat	4117
agagtettta tgtecataag etaegggtag gtetatttte agageatgat acaaatteea	4177
cagg	4181

<211> 829

.<212> PRT

<213> NM\_002387 MCC, mutated in colorectal cancers

<400> 4

Met Asn Ser Gly Val Ala Met Lys Tyr Gly Asn Asp Ser Ser Ala Glu 1 5 10 15

Leu Ser Glu Leu His Ser Ala Ala Leu Ala Ser Leu Lys Gly Asp Ile 20 25 30

Val Glu Leu Asn Lys Arg Leu Gln Gln Thr Glu Arg Glu Arg Asp Leu 35 40 45

Leu Glu Lys Lys Leu Ala Lys Ala Gln Cys Glu Gln Ser His Leu Met 50 55 60

Arg Glu His Glu Asp Val Gln Glu Arg Thr Thr Leu Arg Tyr Glu Glu 65 70 75 80

Arg Ile Thr Glu Leu His Ser Val Ile Ala Glu Leu Asn Lys Lys Ile 85 90 95

Asp Arg Leu Gln Gly Thr Thr Ile Arg Glu Glu Asp Glu Tyr Ser Glu 100 105 110

Leu Arg Ser Glu Leu Ser Gln Ser Gln His Glu Val Asn Glu Asp Ser 115 120 125

Arg Ser Met Asp Gln Asp Gln Thr Ser Val Ser Ile Pro Glu Asn Gln 130 135 140

Ser Thr Met Val Thr Ala Asp Met Asp Asn Cys Ser Asp Leu Asn Ser 145 150 155 160

Glu Leu Gln Arg Val Leu Thr Gly Leu Glu Asn Val Val Cys Gly Arg 165 170 175

Lys Lys Ser Ser Cys Ser Leu Ser Val Ala Glu Val Asp Arg His Ile 180 185 190

Glu Gln Leu Thr Thr Ala Ser Glu His Cys Asp Leu Ala Ile Lys Thr 195 200 205

Val Glu Glu Ile Glu Gly Val Leu Gly Arg Asp Leu Tyr Pro Asn Leu 210 215 220

4. ...

Ala Glu Glu Arg Ser Arg Trp Glu Lys Glu Leu Ala Gly Leu Arg Glu 230 . 235 Glu Asn Glu Ser Leu Thr Ala Met Leu Cys Ser Lys Glu Glu Glu Leu Asn Arg Thr Lys Ala Thr Met Asn Ala Ile Arg Glu Glu Arg Asp Arg 260 265 270 Leu Arg Arg Val Arg Glu Leu Gln Thr Arg Leu Gln Ser Val Gln . 280 ... Ala Thr Gly Pro Ser Ser Pro Gly Arg Leu Thr Ser Thr Asn Arg Pro Ile Asn Pro Ser Thr Gly Glu Leu Ser Thr Ser Ser Ser Asn Asp 310 315 Ile Pro Ile Ala Lys Ile Ala Glu Arg Val Lys Leu Ser Lys Thr Arg . 330 Ser Glu Ser Ser Ser Asp Arg Pro Val Leu Gly Ser Glu Ile Ser 345 Ser Ile Gly Val Ser Ser Ser Val Ala Glu His Leu Ala His Ser Leu . 360 Gln Asp Cys Ser Asn Ile Gln Glu Ile Phe Gln Thr Leu Tyr Ser His 375 Gly Ser Ala Ile Ser Glu Ser Lys Ile Arg Glu. Phe Glu Val Glu Thr 385 390 395 Glu Arg Leu Asn Ser Arg Ile Glu His Leu Lys Ser Gln Asn Asp Leu . 415 405 Leu Thr Ile Thr Leu Glu Glu Cys Lys Ser Asn Ala Glu Arg Met Ser Met Leu Val Gly Lys Tyr Glu Ser Asn Ala Thr Ala Leu Arg Leu Ala 440 Leu Gln Tyr Ser Glu Gln Cys Ile Glu Ala Tyr Glu Leu Leu Ala Leu Ala Glu Ser Glu Gln Ser Leu Ile Leu Gly Gln Phe Arg Ala Ala

Gly Val Gly Ser Ser Pro Gly Asp Gln Ser Gly Asp Glu Asn Ile Thr

490

- Gln Met Leu Lys Arg Ala His Asp Cys Arg Lys Thr Ala Glu Asn Ala 500 505 510
- Ala Lys Ala Leu Leu Met Lys Leu Asp Gly Ser Cys Gly Gly Ala Phe 515 520 525
- Ala Val Ala Gly Cys Ser Val Gln Pro Trp Glu Ser Leu Ser Ser Asn 530 535 540
- Ser His Thr Ser Thr Thr Ser Ser Thr Ala Ser Ser Cys Asp Thr Glu 545 550 555 560
- Phe Thr Lys Glu Asp Glu Gln Arg Leu Lys Asp Tyr Ile Gln Gln Leu 565 570 575
- Lys Asn Asp Arg Ala Ala Val Lys Leu Thr Met Leu Glu Leu Glu Ser 580 585 590
- Ile His Ile Asp Pro Leu Ser Tyr Asp Val Lys Pro Arg Gly Asp Ser 595 600 605
- Gln Arg Leu Asp Leu Glu Asn Ala Val Leu Met Gln Glu Leu Met Ala 610 615 620
- Met Lys Glu Glu Met Ala Glu Leu Lys Ala Gln Leu Tyr Leu Leu Glu 625 630 635 640
- Lys Glu Lys Lys Ala Leu Glu Leu Lys Leu Ser Thr Arg Glu Ala Gln 645 650 655
- Glu Gln Ala Tyr Leu Val His Ile Glu His Leu Lys Ser Glu Val Glu 660 665 670
- Glu Gln Lys Glu Gln Arg Met Arg Ser Leu Ser Ser Thr Ser Ser Gly 675 680 685
- Ser Lys Asp Lys Pro Gly Lys Glu Cys Ala Asp Ala Ala Ser Pro Ala 690 695 700
- Leu Ser Leu Ala Glu Leu Arg Thr Thr Cys Ser Glu Asn Glu Leu Ala 705 710 715 720
- Ala Glu Phe Thr Asn Ala Ile Arg Arg Glu Lys Lys Leu Lys Ala Arg 725 730 735
- Val Gln Glu Leu Val Ser Ala Leu Glu Arg Leu Thr Lys Ser Ser Glu 740 745 750
- Ile Arg His Gln Gln Ser Ala Glu Phe Val Asn Asp Leu Lys Arg Ala 755 760 765

Asn Ser Asn Leu Val Ala Ala Tyr Glu Lys Ala Lys Lys His Gln 775 780 Asn Lys Leu Lys Lys Leu Glu Ser Gln Met Met Ala Met Val Glu Arg 790 795 His Glu Thr Gln Val Arg Met Leu Lys Gln Arg Ile Ala Leu Leu Glu 810 Glu Glu Asn Ser Arg Pro His Thr Asn Glu Thr Ser Leu 820 825 <210> 5 <211> 2830 <212> DNA <213> NM 022117 SE20-4 <220> <221> CDS <222> (130)..(2208) <223> <400> 5 aatteggeac gaggagaget ggttgegtga gteteeteag etetgettae eggtgegaet 60 agcggcagcg acgcggctaa aagcgaaggg gcgagtgcga gtcccctgag ctgtacgaac 120 geggtegee atg gae ege eea gat gag ggg eet eeg gee aag ace ege ege 171 Met Asp Arg Pro Asp Glu Gly Pro Pro Ala Lys Thr Arg Arg 1 10 ctg agc agc tcc gag tct cca cag cgc gac ccg ccc ccg ccg ccg 219 Leu Ser Ser Ser Glu Ser Pro Gln Arg Asp Pro Pro Pro Pro Pro 20 25 ccg ccg ccc ctc ctc cga ctg ccg ctg cct cca ccc cag cag cgc ccg 267 Pro Pro Pro Leu Leu Arg Leu Pro Leu Pro Pro Pro Gln Gln Arg Pro 40 agg ctc cag gag gaa acg gag gcg gca cag gtg ctg gcc gat atg agg 315 Arg Leu Gln Glu Glu Thr Glu Ala Ala Gln Val Leu Ala Asp Met Arg 55 ggg gtg gga ctg ggc ccc gcg ctg ccc ccg ccg cct ccc tat gtc att 363 Gly Val Gly Leu Gly Pro Ala Leu Pro Pro Pro Pro Pro Tyr Val Ile 70 ctc gag gag ggg ggg atc cgc gca tac ttc acg ctc ggt gct gag tgt 411 Leu Glu Glu Gly Gly Ile Arg Ala Tyr Phe Thr Leu Gly Ala Glu Cys 80 85 ccc ggc tgg gat tct acc atc gag tcg ggg tat ggg gag gcg ccc ccg 459

Pro Gly Trp Asp Ser Thr Ile Glu Ser Gly Tyr Gly Glu Ala Pro Pro

95	•			100			105				110	
						act Thr 120	Pro					507
						cag Gln						555
						gtg Val						603
						gcg Ala						651
						agg Arg						699
						gtg Val 200						747
_		_	 	_	 -	ctg Leu	_	_	_	 -		795
						aag .Lys						843
						cga Arg						891
-	_			_		ggc Gly			_	_		939
						atc Ile 280						987
						gta Val						1035
						ttc Phe						1083
						cag Gln						1131
						tgg Trp						1179
						gcg Ala 360						1227
						gct Ala						1275

			370					375	. •				380			
	aag Lys							Pro								1323
	ggc Gly 400															1371
	agg Arg															1419
	gca Ala															1467
	cat His															1515
	acc Thr		-					_						_	-	1563
	gag Glu 480															1611
	aac Asn		_	-	_	-		_			-					1659
_	gat Asp	-						_	-			_				1707
-	aac Asn	_														1755
	ttc Phe															1803
	gac Asp 560															1851
	ggc Gly															1899
ggt Gly	gac Asp	aat Asn	gaa Glu	ggc Gly 595	agc Ser	gat Asp	gat Asp	gac Asp	gac Asp 600	aga Arg	gac Asp	att Ile	gag Glu	tac Tyr 605	tat Tyr	1947
	aaa Lys															1995
	ata Ile															2043
gaa Glu	ggc Gly	atc Ile	cag Gln	caa Gln	gat Asp	gag Gļu	gac Asp	atc Ile	tat Tyr	gag Glu	gaa Glu	gga Gly	aac Asn	tat Tyr	gag Glu	2091

6	40		6	545				650					
			a gat g u Asp V 660										2139
			t gtg c p Val I 5										2187
			a acc g s Thr G	_	gggt	ttt (	ccct	ttt	gg gg	gato	acct		2238
tctgt	atccc	ccaccca	cta tcc	catttg	c cc	tcct	cctc	agc	tagg	gcc a	acgc	ggcccc	2298
acatt	gcact 1	tctgggg	ggt gac	cgactt	c gt	acac	gggt	tta	aagti	ta	tttt	tatggt	2358
ttagt	cattg (	cagagtt	ctt att	ttgggg	g ga	gggaa	aagg	ggg	ctagi	cc (	cctt	cttttg	2418
gccct	ccgcc (	cccgcag	gct tct	gtgtgc	t gc	taaci	tgta	ttt	attgi	tga '	tgcci	ttggtc	2478
agggc	ccctc 1	tacccac	tte tee	cagtca	g tt	gtgg	cccc	agc	ccct	etc (	cctg	tgctgt	2538
gtgga	gtgga (	caccctg	acc ccc	gaagcg	g gg	agggo	ccgc	tgt	ggcct	tc (	gtca	cagccg	2598
cgcagt	tgccc a	atggagg	cgc tgc	tgccac	c tt	cctct	ccc.	aag	ttctt	tc t	tccat	tccctc	2658
tcctc	ttccc	geegege	cgc tag	cccgcc	t cg	gtgto	ctat	gca	aggco	egc 1	ttcg	ccattg	2718
cggtai	ttctt 1	tgcggta	ttc ttg	tccccg	t cc	ccca	gaag	gct	egeet	ct (	cccç	gtggac	2778
cctgt	taatc o	ccaataa	aat tct	gagcaa	g tt	caaaa	aaa	aaaa	aaaa	aaa a	aa		2830
<210>	6				•								
<211>					•								
<212>													
<213>		22117 S	E20-4				•						
12.50	0.		20 1									•	
<400>	6												
Met As	sp Arg	Pro As 5	p Glu G	ly Pro	Pro	Ala 10	Lys	Thr	Arg	Arg	Leu 15	Ser	
Ser Se	er Glu	Ser Pr 20	o Gln A	rg Asp	Pro 25	Pro	Pro	Pro	Pro	Pro 30	Pro	Pro	
Pro Le	eu Leu 35	Arg Le	u Pro L	eu Pro 40	Pro	Pro	Gln	Gln	Arg 45	Pro	Arg	Leu	
Gln Gl 50		Thr Gl	u Ala A 5		Val	Leu	Ala	Asp 60	Met	Arg	Gly	Val	
Gly Le 65	eu Gly	Pro Al	a Leu P 70	ro Pro	Pro	Pro	Pro 75	Tyr.	Val	Ile	Leu	Glu 80	

Glu Gly Gly Ile Arg Ala Tyr Phe Thr Leu Gly Ala Glu Cys Pro Gly

85 90 95

Trp Asp Ser Thr Ile Glu Ser Gly Tyr Gly Glu Ala Pro Pro Pro Thr 100 105 110

Glu Ser Leu Glu Ala Leu Pro Thr Pro Glu Ala Ser Gly Gly Ser Leu 115 120 125

Glu Ile Asp Phe Gln Val Val Gln Ser Ser Ser Phe Gly Gly Glu Gly 130 135 140

Ala Leu Glu Thr Cys Ser Ala Val Gly Trp Ala Pro Gln Arg Leu Val 145 150 155 160

Asp Pro Lys Ser Lys Glu Glu Ala Ile Ile Ile Val Glu Asp Glu Asp 165 170 175

Glu Asp Glu Arg Glu Ser Met Arg Ser Ser Arg Arg Arg Arg Arg 180 185 190

Arg Arg Arg Lys Gln Arg Lys Val Lys Arg Glu Ser Arg Glu Arg Asn 195 200 205

Ala Glu Arg Met Glu Ser Ile Leu Gln Ala Leu Glu Asp Ile Gln Leu 210 215 220

Asp Leu Glu Ala Val Asn Ile Lys Ala Gly Lys Ala Phe Leu Arg Leu 225 230 235 240

Lys Arg Lys Phe Ile Gln Met Arg Arg Pro Phe Leu Glu Arg Arg Asp 245 250 255

Leu Ile Ile Gln His Ile Pro Gly Phe Trp Val Lys Ala Phe Leu Asn 260 265 270

His Pro Arg Ile Ser Ile Leu Ile Asn Arg Arg Asp Glu Asp Ile Phe 275 280 285

Arg Tyr Leu Thr Asn Leu Gln Val Gln Asp Leu Arg His Ile Ser Met 290 295 300

Gly Tyr Lys Met Lys Leu Tyr Phe Gln Thr Asn Pro Tyr Phe Thr Asn 305 310 315 320

Met Val Ile Val Lys Glu Phe Gln Arg Asn Arg Ser Gly Arg Leu Val 325 330 335

Ser His Ser Thr Pro Ile Arg Trp His Arg Gly Gln Glu Pro Gln Ala 340 345 350

Arg Arg His Gly Asn Gln Asp Ala Ser His Ser Phe Phe Ser Trp Phe

355 360 365

Ser Asn His Ser Leu Pro Glu Ala Asp Arg Ile Ala Glu Ile Ile Lys 370 375 380

Asn Asp Leu Trp Val Asn Pro Leu Arg Tyr Tyr Leu Arg Glu Arg Gly 385 390 395 400

Ser Arg Ile Lys Arg Lys Lys Gln Glu Met Lys Lys Arg Lys Thr Arg 405 410 415

Gly Arg Cys Glu Val Val Ile Met Glu Asp Ala Pro Asp Tyr Tyr Ala 420 425 430

Val Glu Asp Ile Phe Ser Glu Ile Ser Asp Ile Asp Glu Thr Ile His 435 440 445

Asp Ile Lys Ile Ser Asp Phe Met Glu Thr Thr Asp Tyr Phe Glu Thr 450 455 460

Thr Asp Asn Glu Ile Thr Asp Ile Asn Glu Asn Ile Cys Asp Ser Glu 465 470 475 480

Asn Pro Asp His Asn Glu Val Pro Asn Asn Glu Thr Thr Asp Asn Asn 485 490 495

Glu Ser Ala Asp Asp His Glu Thr Thr Asp Asn Asn Glu Ser Ala Asp 500 505 510

Asp Asn Asn Glu Asn Pro Glu Asp Asn Asn Lys Asn Thr Asp Asp Asn 515 520 525

Glu Glu Asn Pro Asn Asn Asn Glu Asn Thr Tyr Gly Asn Asn Phe Phe 530 535 540

Lys Gly Gly Phe Trp Gly Ser His Gly Asn Asn Gln Asp Ser Ser Asp 545 550 550 560

Ser Asp Asn Glu Ala Asp Glu Ala Ser Asp Asp Glu Asp Asn Asp Gly
565 570 575

Asn Glu Gly Asp Asn Glu Gly Ser Asp Asp Gly Asn Glu Gly Asp 580 585 590

Asn Glu Gly Ser Asp Asp Asp Asp Asp Ile Glu Tyr Tyr Glu Lys 595 600 . 605

Val Ile Glu Asp Phe Asp Lys Asp Gln Ala Asp Tyr Glu Asp Val Ile 610 615 620

Glu Ile Ile Ser Asp Glu Ser Val Glu Glu Glu Gly Ile Glu Glu Gly

60

720

780

840

625 630 635 640

Ile Gln Gln Asp Glu Asp Ile Tyr Glu Glu Gly Asn Tyr Glu Glu Glu 645 650 655

Gly Ser Glu Asp Val Trp Glu Glu Gly Glu Asp Ser Asp Asp Ser Asp 660 665 670

Leu Glu Asp Val Leu Gln Val Pro Asn Gly Trp Ala Asn Pro Gly Lys 675 680 685

Arg Gly Lys Thr Gly 690

<210> 7

<211> 2632

<212> DNA

<213> NM\_016463 HSPC195

<220>

<221> CDS

<222> (931)..(1611)

<223>

<400> 7

gagcagacge cegageegag egagaagage ggeagageet tateceetga ageegggeee 120 cgcgtcccag ccctgcccag cccgcgccca gccatgcgcg ccgcctgctg agtccgggcg ecgeacgetg ageceteege ecgegageeg egeteagete gggggtgatt agttgetttt 240 tgttgttttt taatttgggc cgcggggagg gggaggaggg gcaggtgctg caggctcccc 300 cccctccccg cctcgggcca gccgcggcgg cgcgactcgg gctccggacc cgggcactgc 360 tggcggctgg agcggagcgc accgcggcgg tggtgcccag agcggagcgc agctccctgc 420 cccgcccctc cccctcggcc tcgcggcgac ggcggcggtg gcggcttgga cgactcggag 480 agccgagtga agacatttcc acctggacac ctgaccatgt gcctgccctg agcagcgagg 540 cccaccaggc atctctgttg tgggcagcag ggccaggtcc tggtctgtgg accctcggca 600 gttggcaggc tecetetgca gtggggtetg ggceteggce ccaccatgte gagectegge 660

ggtggctccc aggatgccgg cggcagtagc agcagcagca ccaatggcag cggtggcagt

ggcagcagtg gcccaaaggc aggagcagca gacaagagtg cagtggtggc tgccgccgca

ccagcctcag tggcagatga cacaccaccc cccgagcgtc ggaacaagag cggtatcatc

ggcggctgag cctgagcggg gatgtagagg cggcggcagc agaggcggca ctggcggcaa

agtgageece teaacaagag eetgegeege teeegeeege teteecacta etettettt 900													
ggcagcagtg gtggtagtgg cggtggcagc atg atg ggc gga gag tct gct gac 95  Met Met Gly Gly Glu Ser Ala Asp  1 5													
aag gcc act gcg gct gca gcc gct gcc tcc ctg ttg gcc aat ggg cat Lys Ala Thr Ala Ala Ala Ala Ala Ser Leu Leu Ala Asn Gly His 10 15 20	1002												
gac ctg gcg gcg gcc atg gcg gtg gac aaa agc aac cct acc tca aag Asp Leu Ala Ala Met Ala Val Asp Lys Ser Asn Pro Thr Ser Lys 25 30 35 40	1050												
cac aaa agt ggt gct gtg gcc agc ctg ctg agc aag gca gag cgg gcc His Lys Ser Gly Ala Val Ala Ser Leu Leu Ser Lys Ala Glu Arg Ala 45 50 55	1098												
acg gag ctg gca gcc gag gga cag ctg acg ctg cag cag ttt gcg cag Thr Glu Leu Ala Ala Glu Gly Gln Leu Thr Leu Gln Gln Phe Ala Gln 60 65 70	1146												
tcc aca gag atg ctg aag cgc gtg gtg cag gag cat ctc ccg ctg atg Ser Thr Glu Met Leu Lys Arg Val Val Gln Glu His Leu Pro Leu Met 75 80 85	1194												
agc gag gcg ggt gct ggc ctg cct gac atg gag gct gtg gca ggt gcc Ser Glu Ala Gly Ala Gly Leu Pro Asp Met Glu Ala Val Ala Gly Ala 90 95 100	1242												
gaa gcc ctc aat ggc cag tcc gac ttc ccc tac ctg ggc gct ttc ccc Glu Ala Leu Asn Gly Gln Ser Asp Phe Pro Tyr Leu Gly Ala Phe Pro 105 110 115 120	1290												
atc aac cca ggc ctc ttc att atg acc ccg gca ggt gtg ttc ctg gcc Ile Asn Pro Gly Leu Phe Ile Met Thr Pro Ala Gly Val Phe Leu Ala 125 130 135	1338												
gag agc gcg ctg cac atg gcg ggc ctg gct gag tac ccc atg cag gga Glu Ser Ala Leu His Met Ala Gly Leu Ala Glu Tyr Pro Met Gln Gly 140 145 150	1386												
gag ctg gcc tct gcc atc agc tcc ggc aag aag cgg aaa cgc tgc Glu Leu Ala Ser Ala Ile Ser Ser Gly Lys Lys Lys Arg Lys Arg Cys 155 160 165	1434												
ggc atg tgc gcg ccc tgc cgg cgc atc aac tgc gag cag tgc agc Gly Met Cys Ala Pro Cys Arg Arg Ile Asn Cys Glu Gln Cys Ser 170 175 180	1482												
agt tgt agg aat cga aag act ggc cat cag att tgc aaa ttc aga aaa Ser Cys Arg Asn Arg Lys Thr Gly His Gln Ile Cys Lys Phe Arg Lys 185 190 195 200	1530												
tgt gag gaa ctc aaa aag aag cct tcc gct gct ctg gag aag gtg atg Cys Glu Glu Leu Lys Lys Lys Pro Ser Ala Ala Leu Glu Lys Val Met 205 210 215	1578												
ctt ccg acg gga gcc gcc ttc cgg tgg ttt cag tgacggcggc ggaacccaaa Leu Pro Thr Gly Ala Ala Phe Arg Trp Phe Gln 220 225	1631												
gctgccctct ccgtgcaatg tcactgctcg tgtggtctcc agcaagggat tcgggcgaag	1691												
acaaacggat gcacccgtct ttagaaccaa aaatattctc tcacagattt cattcctgtt 175													
tttatatata tattttttgt tgtcgtttta acatctccac gtccctagca taaaaagaaa	1811												

aagaaaaaa	tttaaactgc	tttttcggaa	gaacaacaac	aaaaagagg	taaagacgaa	1871
tctataaagt	accgagactt	cctgggcaaa	gaatggacaa	tcagtttcct	tcctgtgtcg	1931
atgtcgatgt	tgtctgtgca	ggagatgcag	tttttgtgta	gagaatgtaa	attttctgta	1991
accttttgaa	atctagttac	taataagcac	tactgtaatt	tagcacagtt	taactccacc	2051
ctcatttaaa	cttcctttga	ttctttccga	ccatgaaata	gtgcatagtt	tgcctggaga	2111
atccactcac	gttcataaag	agaatgttga	tggcgccgtg	tagaagccgc	tctgtatcca	2171
tccacgcgtg	cagagetgee	agcagggagc	tcacagaagg	ggagggagca	ccaggccagc	2231
tgagctgcac	ccacagtccc	gagactggga	tecceacee	caacagtgat	tttggaaaaa	2291
aaaatgaaag	ttctgttcgt	ttatccattg	cgatctgggg	agccccatct	cgatatttcc	2351
aatcctggct	acttttctta	gagaaaataa	gtccttttt	tctggccttg	ctaatggcaa	2411
cagaagaaag	ggcttctttg	cgtggtcccc	tgctggtggg	ggtgggtccc	cagggggccc	2471
cctgcggcct	gggcccccct	gcccacggcc	agcttcctgc	tgatgaacat	gctgtttgta	2531
ttgttttagg	aaaccaggct	gttttgtgaa	taaaacgaat	gcatgtttgt	gtcacgaaaa	2591
aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	aaaaaaaa	a		2632

<211> 227

<212> PRT

<213> NM\_016463 HSPC195

<400> 8

Met Met Gly Gly Glu Ser Ala Asp Lys Ala Thr Ala Ala Ala Ala Ala 1 10 15

Ala Ser Leu Leu Ala Asn Gly His Asp Leu Ala Ala Ala Met Ala Val 20 25 30

Asp Lys Ser Asn Pro Thr Ser Lys His Lys Ser Gly Ala Val Ala Ser 35 40 45

Leu Leu Ser Lys Ala Glu Arg Ala Thr Glu Leu Ala Ala Glu Gly Gln 50 55 60

Leu Thr Leu Gln Gln Phe Ala Gln Ser Thr Glu Met Leu Lys Arg Val 65 70 75 80

Val Gln Glu His Leu Pro Leu Met Ser Glu Ala Gly Ala Gly Leu Pro  $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$ 

Pho	e Pro	Tyr 115		Gly	Ala	Phe	Pro 120		Asn	Pro	Gly	Leu 125		Ile	Met	
Th:	r Pro 130	Ala	Gly	Val	Phe	Leu 135		Glu	Ser	Ala	Leu 140	His	Met	Ala	Gly	
Le:		Glu	Tyr	Pro	Met 150		Gly	Glu	Leu	Ala 155		Ala	Ile	Ser	Ser 160	
Gl	y Lys	Lys	Lys	Arg 165		Arg	Суз	Gly	Met 170	Суз	Ala	Pro	Cys	Arg 175	Arg	
Ar	J Ile	Asn	Cys 180	Glu	Gln	Cys	Ser	Ser 185	_	Arg	Asn	Arg	Lys 190		Gly	
His	s Glm	11e 195		Lys	Phe	Arg	Lys 200		Glu	Glu	Leu	Lys 205	Lys	Lys	Pro	
Sea	210	Ala	Leu	Glu	Lys	Val .215	Met	Leu	Pro	Thr	Gly 220	Ala	Ala	Phe	Arg	
Trp 225	Phe	Gln														
<21	.0>	9			•				•				-			
<21	.1>	2140								•						
<21	.2>	DNA														
<21	.3>	NM_0:	17697	7 FL	J201	71										
<22	:0>															
<22	:1>	CDS					•									
<22	2>	(58)	(11	131)					-	,						
<22	3>						٠				٠					
<40 gaa		9 aga a	aatgt	tgeo	c to	gtto	acct	: gat	atto	raca	aact	ggag	gt t	tqcca	ıca	57
		gag Glu														105
gga Gly	gcc Ala	tct Ser	caa Gln 20	gtt Val	gaa Glu	gat Asp	atg Met	ggg Gly 25	aat Asn	ata Ile	att Ile	tta Leu	gca Ala 30	atg Met	att Ile	153
		cct Pro 35														201

															aac Asn	249
acc Thr 65	gta Val	gto Val	agg Arg	gca Ala	cga Arg 70	ggt Gly	tta Leu	cca Pro	tgg Trp	cag Gln 75	tct Ser	tca Ser	gat Asp	caa Gln	gat Asp 80	297
att Ile	gca Ala	aga Arg	ttc Phe	tto Phe 85	aaa Lys	gga Gly	ctc	aat Asn	att Ile 90	gcc Ala	aag Lys	gga Gly	ggt Gly	gca Ala 95	gca Ala	345
				Ala					Asn					Val	agg Arg	393
ttt Phe	gta Val	agt Ser 115	Glu	gag Glu	cac His	cga Arg	gac Asp 120	Leu	gca Ala	cta Leu	cag Gln	agg Arg 125	His	aaa Lys	cat His	441
cac His	atg Met 130	Gly	acc Thr	cgg Arg	tat Tyr	att Ile 135	gag Glu	gtt Val	tac Tyr	aaa Lys	gca Ala 140	Thr	ggt Gly	gaa Glu	gat Asp	489
ttc Phe 145	Leu	aaa Lys	att Ile	gct Ala	ggt Gly 150	ggt Gly	act Thr	tcc Ser	aat Asn	gag Glu 155	gta Val	gcc	cag Gln	ttt Phe	ctc Leu 160	537
tcc Ser	aag Lys	gaa Glu	aat Asn	caa Gln 165	gtc Val	att Ile	gtt Val	cgc Arg	atg Met 170	cgg	Gly	ctc Leu	cct Pro	ttc Phe 175	acg Thr	585
					gtg Val											633
					Gly											681
cca Pro	aca Thr 210	Gly	gac Asp	gct Ala	ttt Phe	gtc Val 215	ctc Leu	ttt Phe	gcc Ala	tgt Cys	gag Glu 220	gaa Glu	tat Tyr	gca Ala	cag Gln	729
aat Asn 225	gcg Ala	ttg Leu	agg Arg	aag Lys	cat His 230	aaa Lys	gac Asp	ttg Leu	ttg Leu	ggt Gly 235	aaa Lys	aga Arg	tac Tyr	att Ile	gaa Glu 240	777
ctc Leu	ttc Phe	agg Arg	agc Ser	aca Thr 245	gca Ala	gct Ala	gaa Glu	gtt Val	cag Gln 250	cag Gln	gtg Val	ctg Leu	aat Asn	cga Arg 255	ttc Phe	825
tcc Ser	tcg Ser	gcc Ala	cct Pro 260	ctc Leu	att Ile	cca Pro	ctt Leu	cca Pro 265	acc Thr	cct Pro	ccc Pro	att Ile	att Ile 270	cca Pro	gta Val	873
cta Leu	cct Pro	cag Gln 275	caa Gln	ttt Phe	gtg Val	Pro	cct Pro 280	aca Thr	aat Asn	gtt Val	aga Arg	gac Asp 285	tgt Cys	ata Ile	cgc Arg	921
Leu	cga Arg 290	ggt Gly	ctt Leu	ccc Pro	tat Tyr	gca Ala 295	gcc Ala	aca Thr	att Ile	gag Glu	gac Asp 300	atc Ile	ctg Leu	gat Asp	ttc Phe	969
ctg Leu 305	ggg ggg	gag Glu	ttc Phe	gcc Ala	aca Thr 310	gat Asp	att Ile	cgt Arg	Thr	cat His 315	GJ Y GGG	gtt Val	cac His	atg Met	gtt Val 320	1017

ttg aat cac cag ggc cgc cca tca gga gat gcc ttt atc cag atg aag Leu Asn His Gln Gly Arg Pro Ser Gly Asp Ala Phe Ile Gln Met Lys 325 330 335	1065
tct gcg gac aga gca ttt atg gct gca cag aag tgt cat aaa aaa aaa Ser Ala Asp Arg Ala Phe Met Ala Ala Gln Lys Cys His Lys Lys 340 345 350	1113
cat gaa gga cag ata tgt tgaagtettt cagtgtteag etgaggagat His Glu Gly Gln Ile Cys 355	1161
gaactttgtg ttaatggggg gcactttaaa tcgaaatggc ttatccccac cgccatgtaa	1221
gttaccatgc ctgtctcctc cctcctacac atttccagct cctgctgcag ttattcctac	1281
agaagetgee atttaccage cetetgtgat tttgaateca egageaetge agecetecae	1341
agcgtactac ccagcaggca ctcagctctt catgaactac acagcgtact atcccagccc	1401
cccaggttcg cctaatagtc ttggctactt ccctacagct gctaatctta gcggtgtccc	1461
tccacagcct ggcacggtgg tcagaatgca gggcctggcc tacaatactg gagttaagga	1521
aattettaac ttetteeaag gttaeeagtg tttgaaagat gtatggtgat ettgaaacet	1581
ccagacacaa gaaaacttct agcaaattca ggggaagttt gtctacactc aggctgcagt	1641
attttcagca aacttgattg gacaaacggg cctgtgcctt atcttttggt ggagtgaaaa	1701
agtttgagct agtgaagcca aatcgtaact tacagcaagc agcatgcagc atacctggct	1761
ctttgctgat tgcaaatagg catttaaaat gtgaatttgg aatcagatgt ctccattact	1821
tccagttaaa gtggcatcat aggtgtttcc taagttttaa gtcttggata aaaactccac	1881
cagtgtctac catctccacc atgaactctg ttaaggaagc ttcatttttg tatattcccg	1941
ctcttttctc ttcatttccc tgtcttctgc ataatcatgc cttcttgcta agtaattcaa	2001
gcataagatc ttggaataat aaaatcacaa tcttaggaga aagaataaaa ttgttatttt	2061
cccagtetet tggccatgat gatatettat gattaaaaac aaattaaatt ttaaaacacc	2121
tgaaaaaaaa aaaaaaaa	2140

<211> 358

<212> PRT

<213> NM\_017697 FLJ20171

<400> 10

Met Thr Glu Tyr Leu Asn Phe Glu Lys Ser Ser Ser Val Ser Arg Tyr 1 5 10 15 .

Gly Ala Ser Gln Val Glu Asp Met Gly Asn Ile Ile Leu Ala Met Ile 20 25 30

Ser Glu Pro Tyr Asn His Arg Phe Ser Asp Pro Glu Arg Val Asn Tyr 35 40 45

Lys Phe Glu Ser Gly Thr Cys Ser Lys Met Glu Leu Ile Asp Asp Asn 50 55 60

Thr Val Val Arg Ala Arg Gly Leu Pro Trp Gln Ser Ser Asp Gln Asp 65 70 75 80

Ile Ala Arg Phe Phe Lys Gly Leu Asn Ile Ala Lys Gly Gly Ala Ala 85 90 95

Leu Cys Leu Asn Ala Gln Gly Arg Arg Asn Gly Glu Ala Leu Val Arg 100 105 110

Phe Val Ser Glu Glu His Arg Asp Leu Ala Leu Gln Arg His Lys His 115 120 125

His Met Gly Thr Arg Tyr Ile Glu Val Tyr Lys Ala Thr Gly Glu Asp 130 135 140

Phe Leu Lys Ile Ala Gly Gly Thr Ser Asn Glu Val Ala Gln Phe Leu 145 150 155 160

Ser Lys Glu Asn Gln Val Ile Val Arg Met Arg Gly Leu Pro Phe Thr 165 170 175

Ala Thr Ala Glu Glu Val Val Ala Phe Phe Gly Gln His Cys Pro Ile 180 185 190

Thr Gly Gly Lys Glu Gly Ile Leu Phe Val Thr Tyr Pro Asp Gly Arg 195 200 205

Pro Thr Gly Asp Ala Phe Val Leu Phe Ala Cys Glu Glu Tyr Ala Gln 210 215 220

Asn Ala Leu Arg Lys His Lys Asp Leu Leu Gly Lys Arg Tyr Ile Glu 225 230 235 240

Leu Phe Arg Ser Thr Ala Ala Glu Val Gln Gln Val Leu Asn Arg Phe 245 250 255

Ser Ser Ala Pro Leu Ile Pro Leu Pro Thr Pro Pro Ile Ile Pro Val 260 265 270

Leu Pro Gln Gln Phe Val Pro Pro Thr Asn Val Arg Asp Cys Ile Arg 275 280 285

Leu Arg Gly Leu Pro Tyr Ala Ala Thr Ile Glu Asp Ile Leu Asp Phe 290 295 300

Leu Gl	/ Glu	Phe	Ala	Thr 310	Asp	Ile	Arg	Thr	His 315	Gly	Val	His	Met	Val 320	
Leu As	n His	Gln	Gly 325	Arg	Pro	Ser	Gly	Asp 330	Ala	Phe	Ile	Gln	Met 335	Lys	
Ser Al	a Asp	Arg 340	Ala	Phe	Met	Ala	Ala 345	Gln	Ъуs	Суз	His	Lys 350	Lys	ГÀЗ	
His Gl	355	Gln	Ile	Суз											
<210>	11														
<211>	2808														
<212>	DNA									•					
<213>	NM_0	5288	6 MAI	L2				•	-		•				
<220>															
<221>	CDS														
<222>	(80)	(6	07)						•						
<223>															
<400>	11 ggcg	gcag	gageo	ec go	ggagg	gegga	a ggo	eggga	iggc	ggco	igeg	jcg (	egegg	, Jagacg	60
<400> ggcggc cagcag	ggcg			c ato	te	g gci	gge	e gga	a gc	g tca	gto	e eeg	gcc		60 112
ggcggc	egge .	agcg:	gcago	c ato Met 1 tcc	g tog Sea	g gcd r Ala	c ggo	c gga y Gly 5	a gcg	g tca a Sei gtc	gto Val	c ccq	g ccc Pro 10	g ccc Pro	
ggcggc cagcag ccg aa	ggcg cggc	gcc Ala 15	gtg Val ctg	tcc Ser	ttc Phe	ccg Pro	ccg Pro 20	c gga y Gly 5 ccc Pro	egg Arg	g toa g Ser gtc Val	acc Thr	ctg Leu 25	g ccc 10 ccc Pro	gcc Ala	112
ggcggc cagcag ccg aa Pro As	eggc eggc ecce Pro egac Asp 30	gcc Ala 15 atc Ile	gtg Val ctg Leu	tcc Ser cgg Arg	ttc Phe acc Thr	ccg Pro tac Tyr 35	ccg Pro 20 tcg Ser	c gga y Gly 5 ccc Pro ggc Gly	cgg Arg gcc Ala	g toa Ser gtc Val ttc Phe	acc Thr gtc Val 40	ctg Leu 25 tgc Cys	ccc Pro ctg	gcc Ala gag Glu	112
cagcag ccg aa Pro As ggc cc Gly Pr	eggc cggc cggc Asp 30 ttc Phe	gcc Ala 15 atc Ile ggg Gly	gtg Val ctg Leu ggt Gly	tcc Ser cgg Arg	ttc Phe acc Thr gtc Val 50	ccg Pro tac Tyr 35 tgg	ccg Pro 20 tcg Ser att Ile	ggc ggc ggc Gly ttg Leu	cgg Arg gcc Ala gtt Val	g toa gtc Val ttc Phe gcc Ala 55	acc Thr gtc Val 40 tcc Ser	ctg Leu 25 tgc Cys tcc Ser	ccc Pro ctg Leu aat Asn	gcc Ala gag Glu gtt Val	112 160 208
cagcagc cagcag ccg aa Pro As ggc cc Gly Pr att ct Ile Le 45 cct ct Pro Le	eggc eggc ccc n Pro egac o Asp 30 ttc n Phe	gcc Ala 15 atc Ile ggg Gly cta Leu	gtg Val ctg Leu ggt Gly	c atg Met 1 tcc Ser cgg Arg ctt Leu caa Gln 65	ttc Phe acc Thr gtc Val 50 gga Gly	ccg Pro tac Tyr 35 tgg Trp tggg	ccg Pro 20 tcg Ser att Ile gtc Val	c ggg Gly 5 ccc Pro ggc Gly ttg Leu atg	cgg gcc Ala gtt Val ttt Phe 70 ttc	gtc Val ttc Phe gcc Ala 55 gtg Val	acc Thr gtc Val 40 tcc Ser tcc	ctg Leu 25 tgc Cys tcc Ser gtg Val	ccc Pro ctg Leu aat Asn aca Thr	gcc Ala gag Glu gtt Val gcg Ala 75	112 160 208 256

aca gta tt Thr Val Ph 11	e Val Phe T	at ttt gga Cyr Phe Gly 115	gcc ttt tta Ala Phe Leu	ttg gaa gca gca gcc Leu Glu Ala Ala Ala 120	448
				ata acc ggg cag cca Ile Thr Gly Gln Pro 135	496
	r Āsp Asn G			gca gcc tca att ttt Ala Ala Ser Ile Phe 155	544
				ttg ggt ctg gct tta Leu Gly Leu Ala Leu 170	592
cga aga tg Arg Arg Tr		aacactcct t	agaaactgg ca	agtcgtatg ttagtttcac	647
ttgtctactt	tatatgtctg	atcaatttgg	ataccatttt	gtccagatgc aaaaacattc	707
caaaagtaat	gtgtttagta	gagagagact	ctaagctcaa	gttctggttt atttcatgga	767
tggaatgtta	attttattat	gatattaaag	aaatggcctt	ttattttaca tctctcccct	827
ttttcccttt	cccctttat	tttectcctt	ttctttctga	aagtttcctt ttatgtccat	887
aaaatacaaa	tatattgttc	ataaaaaatt	agtatccctt	ttgtttggtt gctgagtcac	947
ctgaacctta	attttaattg	gtaattacag	cccctaaaaa	aaacacattt caaataggct	1007
tcccactaaa	ctctatattt	tagtgtaaac	caggaattgg	cacacttttt ttagaatggg	1067
ccagatggta	aatatttatg	cttcacggtc	catacagtct	ctgtcacaac tattcagttc	1127
tgctagtata	gcgtgaaagc	agetatacac	aatacagaaa	tgaatgagtg tggttatgtt	1187
ctaataaaac	ttatttataa	aaacaagggg	aggctgggtt	tagcctgtgg gccatagttt	1247
gtcaaccact	ggtgtaaaac	cttagttata	tatgatctgc	attttcttga actgatcatt	1307
gaaaacttat	aaacctaaca	gaaaagccac	ataatattta	gtgtcattat gcaataatca	1367
cattgccttt	gtgttaatag	tcaaatactt	acctttggag	aatacttacc tttggaggaa	1427
tgtataaaat	ttctcaggca	gagtcctgga	tataggaaaa	agtaatttat gaagtaaact	1487
tcagttgctt	aatcaaacta	atgatagtct	aacaactgag	caagatcctc atctgagagt	1547
gcttaaaatg	ggatccccag	agaccattaa	ccaatactgg	aactggtatc tagctactga	1607
tgtcttactt	tgagtttatt	tatgcttcag	aatacagttg	tttgccctgt gcatgaatat	1667
acccatattt	gtgtgtggat	atgtgaagct	tttccaaata	gageteteag aagaattaag	1727
tttttacttc	taattattt	gcattacttt	gagttaaatt	tgaatagagt attaaatata	1787
aagttgtaga	ttcttatgtg	tttttgtatt	agcccagaca	tctgtaatgt ttttgcactg	1847
gtgacagaca	aaatctgttt	taaaatcata	tccagcacaa	aaactatttc tggctgaata	1907
gcacagaaaa	gtattttaac	ctacctgtag	agatectegt	catggaaagg tgccaaactg	1967
ttttgaatgg	aaggacaagt	aagagtgagg	ccacagttcc	caccacacga gggcttttgt	2027
attgttctac	tttttcagcc	ctttactttc	tggctgaagc	atccccttgg agtgccatgt	2087

ataagttggg	ctattagagt	tcatggaaca	tagaacaacc	atgaatgagt	ggcatgatcc	214
gtgcttaatg	atcaagtgtt	acttatctaa	taatcctcta	gaaagaaccc	tgttagatct	220
tggtttgtga	taaaaatata	aagacagaag	acatgaggaa	aaacaaaagg	tttgaggaaa	226
tcaggcatat	gactttatac	ttaacatcag	atcttttcta	taatatccta	ctactttggt	2327
tttcctagct	ccataccaca	cacctaaacc	tgtattatga	attacatatt	acaaagtcat	2387
aaatgtgcca	tatggatata	cagtacattc	tagttggaat	cgtttactct	gctagaattt	2447
aggtgtgaga	ttttttgttt	cccaggtata	gcaggcttat	gtttggtggc	attaaattgg	2507
tttctttaaa	atgctttggt	ggcacttttg	taaacagatt	gcttctagat	tgttacaaac	2567
caagcctaag	acacatctgt	gaatacttag	atttgtagct	taatcacatt	ctagacttgt	2627
gagttgaatg	acaaagcagt	tgaacaaaaa	ttatggcatt	taagaattta	acatgtctta	2687
gctgtaaaaa	tgagaaagtg	ttggttggtt	ttaaaatctg	gtaactccat	gatgaaaaga	2747
aatttattt	atacgtgtta	tgtctctaat	aaagtättca	tttgataaaa	aaaaaaaaa	2807
a				•		2808

<211> 176

<212> PRT

<213> NM\_052886 MAL2

<400> 12

Met Ser Ala Gly Gly Ala Ser Val Pro Pro Pro Pro Asn Pro Ala Val 1 5 15

Ser Phe Pro Pro Pro Arg Val Thr Leu Pro Ala Gly Pro Asp Ile Leu 20 25 30

Arg Thr Tyr Ser Gly Ala Phe Val Cys Leu Glu Ile Leu Phe Gly Gly 35 40 45

Leu Val Trp Ile Leu Val Ala Ser Ser Asn Val Pro Leu Pro Leu Leu 50 60

Gln Gly Trp Val Met Phe Val Ser Val Thr Ala Phe Phe Phe Ser Leu 70 75 80

Leu Phe Leu Gly Met Phe Leu Ser Gly Met Val Ala Gln Ile Asp Ala 85 90 95

Asn Trp Asn Phe Leu Asp Phe Ala Tyr His Phe Thr Val Phe Val Phe 100 105 110

Tyr Phe Gly Ala Phe Leu Leu Glu Ala Ala Ala Thr Ser Leu His Asp

32

120 125 115 Leu His Cys Asn Thr Thr Ile Thr Gly Gln Pro Leu Leu Ser Asp Asn 135 140 Gln Tyr Asn Ile Asn Val Ala Ala Ser Ile Phe Ala Phe Met Thr Thr Ala Cys Tyr Gly Cys Ser Leu Gly Leu Ala Leu Arg Arg Trp Arg Pro 165 170 <210> 13 <211> 4171 <212> DNA <213> NM 015238 KIAA0869, KIBRA <220> <221> CDS (67) . . (3405) <222> <223> <400> 13 ctggagccgc tgagcccccg ctgcggccgg gagctgcatg ggggagcgcc ggcagcgctt 60 108 gggaag atg ccc cgg ccg gag ctg ccc ctg ccg gag ggc tgg gag gag Met Pro Arg Pro Glu Leu Pro Leu Pro Glu Gly Trp Glu Glu 5 10 gcg cgc gac ttc gac ggc aag gtc tac tac ata gac cac acg aac cgc 156 Ala Arg Asp Phe Asp Gly Lys Val Tyr Tyr Ile Asp His Thr Asn Arg 20 25 204 acc acc age tgg atc gac eeg egg gae agg tac acc aaa eeg etc acc Thr Thr Ser Trp Ile Asp Pro Arg Asp Arg Tyr Thr Lys Pro Leu Thr 35 40 ttt gct gac tgc att agt gat gag ttg ccg cta gga tgg gaa gag gca 252 Phe Ala Asp Cys Ile Ser Asp Glu Leu Pro Leu Gly Trp Glu Glu Ala 50 tat gac cca cag gtt gga gat tac ttc ata gac cac aac acc aaa acc 300 Tyr Asp Pro Gln Val Gly Asp Tyr Phe Ile Asp His Asn Thr Lys Thr act cag att gag gat cct cga gta caa tgg cgg cgg gag cag gaa cat Thr Gln Ile Glu Asp Pro Arg Val Gln Trp Arg Arg Glu Gln Glu His 85 atg ctg aag gat tac ctg gtg gtg gcc cag gag gct ctg agt gca caa

Met Leu Lys Asp Tyr Leu Val Val Ala Gln Glu Ala Leu Ser Ala Gln

aag gag atc tac cag gtg aag cag cag cgc ctg gag ctt gca cag cag

105

444

Lys	Glu	Ile	Tyr	Gln 115	Val	Lys	Gln		Arg 120	Leu	Glu	Leu	Ala	Gln 125	Gln	
						gcc Ala			Glu							492
						tca Ser										540
						gcc Ala 165										588
						ctc Leu										636
		_		-	-	aaa Lys		-			•		_	-	_	684
	_			_	_	gaa Glu	_	_	_	-	_	-	-			732
						acc Thr										, 780
						ttg Leu 245										828
			-	_		tcc Ser	-	-	_		_		-	_	-	876
	_		_		-	tac Tyr	-	-		_		_		-		924
		_				aac Asn	_			_	-	_		_	_	972
						gaa Glu										1020
						ctt Leu 325										1068
						ctg Leu										1116
						atc Ile										1164
				Glu		gcc Ala										1212

											٠.						
	Ala	a gc a Al	c cg a Ar 38	g As	c ac p Th	c ca r Gl	g ag n Se:	c aa r Ly 39	s Ala	g ct	g ac	g ga r Gl	g age u Are 39	g Le	a aa u Ly	g tta s Leu	1260
	aad Asi	n Se 40	г ГА	g ag s Ar	g aa g As	c cae	g cti n Lei 40!	u Va	g aga	a gaa g Gli	a cte	g ga u Glu 410	u Glı	a gc	c ac a Th	c cgg r Arg	1308
	Caq Glr 415	n Va.	g gc l Al	a ac a Th	t cto	g ca u Hi: 420	s Se	c cad	g cto n Lei	g aaa 1.Lys	a agi s.Se: 425	r Lei	c tca u Seu	a age	c ag	c atg r Met 430	1356
	Glr	g tco n Sen	c cte	g tco u Se:	c tca c Sea 435	c Gly	c ago y Sei	c ago	c ccc	gga Gly 440	/ Sei	cto Lei	acç ı Thi	tco Sei	2 age 3 Se: 44!	c cgg r Arg	1404
	Gl y	tco Sei	c cto	g gtt u Val 450	L Ala	tco Sei	ago Ser	cto Lev	g gad 1 Asp 455	Ser	tco Ser	act Thr	tca Ser	gco Ala 460	. Se	ttc r Phe	1452
	act Thr	gac Asp	Let 465	з Туз	tat Tyr	gac Asp	ccc Pro	ttt Phe 470	≥ Glu	cag Gln	ctg Leu	gac Asp	tca Ser 475	Glu	cto Lev	g cag ı Gln	1500
	ago Ser	aag Lys 480	val	g gaç L Glu	tto Phe	ctg Leu	Leu 485	Leu	gag Glu	ggg	gcc Ala	acc Thr 490	Gly	tto Phe	cgg Arg	g ccc	1548
	tca Ser 495	GTA	tgo Cys	ato Ile	acc Thr	Thr 500	Ile	cac His	gag Glu	gat Asp	gag Glu 505	Val	gcc Ala	aag Lys	acc Thr	cag Gln 510	1596
-	aag Lys	gca Ala	gag Glu	gga Gly	ggt Gly 515	Gly	cgc Arg	ctg Leu	cag Gln	gct Ala 520	ctg Leu	cgt Arg	tcc Ser	ctg Leu	tct Ser 525	ggc	1644
	acc Thr	cca Pro	aag Lys	Ser 530	Met	acc Thr	tcc Ser	cta Leu	tcc Ser 535	cca	cgt Arg	tcc Ser	tct Ser	ctc Leu 540	tcc Ser	tcc Ser	1692
	ccc Pro	tcc Ser	cca Pro 545	Pro	tgt Cys	tcc Ser	cct Pro	ctc Leu 550	atg Met	gct Ala	gac Asp	ccc Pro	ctc Leu 555	ctg Leu	gct Ala	ggt Gly	1740
	gat Asp	gcc Ala 560	ttc Phe	ctc Leu	aac Asn	tcc Ser	ttg Leu 565	gag Glu	ttt Phe	gaa Glu	gac Asp	ccg Pro 570	gag Glu	ctg Leu	agt Ser	gcc Ala	1788
	act Thr 575	ctt Leu	tgt Cys	gaa Glu	ctg Leu	agc Ser 580	ctt Leu	ggt Gly	aac Asn	Ser	gcc Ala 585	cag Gln	gaa Glu	aga Arg	tac Tyr	cgg Arg 590	1836
	ctg Leu	gag Glu	gaa Glu	cca Pro	gga Gly 595	acg Thr	gag Glu	ggc	aag Lys	cag Gln 600	ctg Leu	ggc Gly	caa Gln	gct Ala	gtg Val 605	aat Asn	1884
	acg Thr	gcc Ala	cag Gln	ggg Gly 610	tgt Cys	ggc Gly	ctg Leu	aaa Lys	gtg Val 615	gcc Ala	tgt Cys	gtc Val	tca Ser	gcc Ala 620	gcc Ala	gta Val	1932
	tcg Ser	Asp	gag Glu 625	tca Ser	gtg Val	gct Ala	Gly	gac Asp 630	agt Ser	ggt Gly	gtg Val	tac Tyr	gag Glu 635	gct Ala	tcc Ser	gtg Val	1980
	cag	aga	ctg	ggt	gct	tca	gaa	gct	gct	gca	tţt	gac	agt	gac	gaa	tcg	2028

		-	~-		<b>C</b> -	<b>~</b> 3	7.7			ъ.		<u> </u>		٠.	~	
Gin	Arg 640		Gly	Ala	Ser	645	Ala	Ala	Ala	Phe	Asp 650	Ser	Asp	Glu	Ser	
	Ala				acc Thr 660											2076
					gca Ala											2124
					caa Gln											2172
				_	agc Ser			_	_						_	2220
					cta Leu											2268
					cac His 740											2316
					ctg Leu											2364
					cgg Arg											2412
					tac Tyr											2460
					cct Pro											2508
tct Ser 815	Ala	ctg Leu	Leu	Glu	cag Gln 820	Thr	gca Ala	gtg Val	Glu	ctg Leu 825	Glu	aag Lys	agg Arg	cag Gln	gag Glu 830	2556
					cag Gln											2604
					gca Ala											2652
					gag Glu											2700
					gca Ala											2748
	-	-			ccc Pro 900		-		_					_	-	2796

ggc Gly	acc Thr	ccg Pro	tcc Ser	cag Gln 915	ggg Gly	cca Pro	ttt Phe	ctt Leu	cga Arg 920	Gly	agc Ser	acc Thr	atc Ile	atc Ile 925	cgc Arg	2844	Ī
								_	_	_		gtg Val	_		_	2892	?
												aag Lys 955				2940	)
	_			_	-	_	-	_	_			aag Lys				2988	}
_	_	-	_	_	_			_			_	acc Thr	_	_		3036	5
										Tr		e ago s Ser			u Th		Į
				Val					ı - Ly			ag ct In Le	u G		caa Gln	3129	•
			cac His 1025	Gly					1 P:			gg tt cp Le	u Ai		gag Glu	3174	ł
				Arg					j Me			ig aa lu Ly	's Aı		cag Gln	3219	<b>,</b>
				Glu					ı Le			ca ga nr As	p Ly		atg Met	3264	ļ
			gct Ala 1070	Ala					A			ga gg gg Gl	y Gl		agc Ser	3309	,
				Pro					Pl			ig aa Lu Ly	s Me		gca Ala	3354	ŧ
				Pro					Pı			c to eu Se	r Al		gat Asp	3399	,
gac Asp		taat	cgcc	ag a	aaag	tatt	t cc	tttg	ttcc	act	gacc	agg	ctgt	gaa	cat	3455	,
tgac	tgtg	gc t	aaag	ttat	t ta	tgtg	gtġt	tat	atga	agg	tact	gagt	ca c	caagt	tcctc	t 3515	,
agtg	ctct	tg t	tggt	ttga	a ga	tgaa	ccga	ctt	ttta	igtt	tggg	rtcct	ac t	gtt	gttat	t 3575	
aaaa	acag	aa c	aaaa	acaa	a ac	acac	acac	aca	caaa	aac	agaa	acaa	aa a	aaac	ccago	a 3635	
ttaa	aata	at a	agat	tgta	t ag	tttg	tata	ttt	agga	igtg	tatt	tttg	gg a	aaga	aaat	t 3695	
taaa	tgaa	ct a	aagc	agta	t tg	agtt	gctg	ctc	ttct	taa	aato	gttt	ag a	tttt	tttt	g 3755	

gtttgtacag ctcc	accttt tagaggtctt	actgcaataa	gaagtaatgc	ctgggggacg	3815
gtaatcctaa tagga	acgtcc cgcacttgtc	acagtacagc	taatttttcc	tagttaacat	3875
attttgtaca atati	taaaaa aatgcacaga	aaccattggg	ggggattcag	aggtgcatcc	3935
acggatcttc ttgag	gctgtg acgtgtttt	atgtggctgc	ccaacgtgga	gcgggcagtg	3995
tgataggctg ggtgg	ggctaa gcagcctagt	ctatgtgggt	gacaggccac	gctggtctca	4055
gatgcccagt gaage	ccacta acatgagtga	ggggagggct	gtggggaact	ccattcagtt	4115
ttatctccat caata	aaagtg gcctttcaaa	aagaaaaaaa	aaaaaaaaa	aaaaaa	4171

<211> 1113

<212> PRT

<213> NM\_015238 KIAA0869, KIBRA

<400> 14

Met Pro Arg Pro Glu Leu Pro Leu Pro Glu Glu Glu Glu Glu Ala Arg 1 5 10 15

Asp Phe Asp Gly Lys Val Tyr Tyr Tle Asp His Thr Asn Arg Thr Thr 20 25 30

Ser Trp Ile Asp Pro Arg Asp Arg Tyr Thr Lys Pro Leu Thr Phe Ala 35 40 45

Asp Cys Ile Ser Asp Glu Leu Pro Leu Gly Trp Glu Glu Ala Tyr Asp 50 55 60

Pro Gln Val Gly Asp Tyr Phe Ile Asp His Asn Thr Lys Thr Thr Gln 65 70 75 80

Ile Glu Asp Pro Arg Val Gln Trp Arg Arg Glu Gln Glu His Met Leu 85 90 95

Lys Asp Tyr Leu Val Val Ala Gln Glu Ala Leu Ser Ala Gln Lys Glu
100 105 110

Ile Tyr Gln Val Lys Gln Gln Arg Leu Glu Leu Ala Gln Gln Glu Tyr 115 120 125

Gln Gln Leu His Ala Val Trp Glu His Lys Leu Gly Ser Gln Val Ser 130 135 140

Leu Val Ser Gly Ser Ser Ser Ser Eys Tyr Asp Pro Glu Ile Leu 145 150 155 160

Lys	Ala	Glu	Ile	Ala	Thr	Ala	Lys	Ser	Arg	Val	Asn	Lys	Leu	Lys	Arg
				165					170					175	
Glu	Met	Va1	His 180	Leu	Gln	His	Glu	Leu 185	Gln	Phe	Lys	Glu	Arg 190	Gly	Phe
Gln	Thr	Leu 195	Lys	Lys	Ile	Asp	Lys 200	Lys	Met	Ser		Ala 205	Gln	Gly	Ser
Tyr	Lys 210		Asp	Glu	Ala	Gln 215	Ala	Val	Leu	Arg	Glu 220	Thr	Lys	Ala	Ile
Lys 225	Lys	Ala	Ile	Thr	Cys 230	Gly	Glu	Ļys	Glu	Lys 235	Gln	Asp	Leu	Ile	Lys 240
Ser	Leu	Ala	Met	Leu 245	Lys	Asp	Gly	Phe	Arg 250	Thr	Asp	Arg	-	Ser 255	His
Ser	Asp	Leu	Trp 260	Ser	Ser	Ser	Ser	Ser 265	Leu	Glu	Ser	Ser	Ser 270	Phe	Pro
Leu	Pro	Lys 275	Gln	Tyr	Leu	Asp	Val 280	Ser	Ser	Gln	Thr	Asp 285	Ile	Ser	Gly
Ser	Phe 290	Gly	Ile	Asn	Ser	Asn 295	Asn	Gln	Leu	Ala	Glu 300	Lys	Val	Arg	Leu
Arg 305	Leu	Arg	Tyr	Glu	Glu 310	Ala	Lys	Arg	Arg	Ile 315	Ala	Asn	Leu	Lys	Ile 320
Gln	Leu	Ala	Lys	Leu 325		Ser	Glu	Ala	Trp 330	Pro	Gly	Val	Leu	Asp 335	Ser
Glu	Arg	Asp	Arg 340	Leu	Ile	Leu	Ile	Asn 345	Glu	Lys	Glu	Glu	Leu 350	Leu	Lys
Glu	Met	Arg 355	Phe	Ile	Ser	Pro	Arg 360	Lys	Ţrp	Thr	Gln	Gly 365	Glu	Val	Glu
Gln	Leu 370	Glu	Met	Ala	Arg	Lys 375	Arg	Leu	Glu	Lys	Asp 380	Leu	Gln	Ala	Ala
Arg 385	Asp	Thr	Gln	Ser	390	Ala	Leu	Thr	Glu	Arg 395	Leu	Lys	Leu	Asn	Ser 400
Lys	Arg	Asn	Gln	Leu 405	Val	Arg	Glu	Leu	Glu 410	Glu	Ala	Thr	Arg	Gln 415	Val
Ala	Thr	Leu	His 420	Ser	Gln	Leu	Lys	Ser 425	Leu	Ser	Ser	Ser	Met 430	Gln	Ser

Leu Ser Ser Gly Ser Ser Pro Gly Ser Leu Thr Ser Ser Arg Gly Ser 435 440 445

Leu Val Ala Ser Ser Leu Asp Ser Ser Thr Ser Ala Ser Phe Thr Asp 450 455 460

Leu Tyr Tyr Asp Pro Phe Glu Gln Leu Asp Ser Glu Leu Gln Ser Lys 465 470 475 480

Val Glu Phe Leu Leu Glu Gly Ala Thr Gly Phe Arg Pro Ser Gly 485 490 495

Cys Ile Thr Thr Ile His Glu Asp Glu Val Ala Lys Thr Gln Lys Ala 500 505 510

Glu Gly Gly Arg Leu Gln Ala Leu Arg Ser Leu Ser Gly Thr Pro 515 520 525

Lys Ser Met Thr Ser Leu Ser Pro Arg Ser Ser Leu Ser Ser Pro Ser 530 535 540

Pro Pro Cys Ser Pro Leu Met Ala Asp Pro Leu Leu Ala Gly Asp Ala 545 550 560

Phe Leu Asn Ser Leu Glu Phe Glu Asp Pro Glu Leu Ser Ala Thr Leu 565 570 575

Cys Glu Leu Ser Leu Gly Asn Ser Ala Gln Glu Arg Tyr Arg Leu Glu 580 585 590

Glu Pro Gly Thr Glu Gly Lys Gln Leu Gly Gln Ala Val Asn Thr Ala 595 600 605

Gln Gly Cys Gly Leu Lys Val Ala Cys Val Ser Ala Ala Val Ser Asp 610 615 620

Glu Ser Val Ala Gly Asp Ser Gly Val Tyr Glu Ala Ser Val Gln Arg 625 630 635 640

Leu Gly Ala Ser Glu Ala Ala Ala Phe Asp Ser Asp Glu Ser Glu Ala 645 650 655

Val Gly Ala Thr Arg Ile Gln Ile Ala Leu Lys Tyr Asp Glu Lys Asn 660 665 670

Lys Gln Phe Ala Ile Leu Ile Ile Gln Leu Ser Asn Leu Ser Ala Leu 675 680 685

Leu Gln Gln Gln Asp Gln Lys Val Asn Ile Arg Val Ala Val Leu Pro 690 695 700

Cys Ser Glu Ser Thr Thr Cys Leu Phe Arg Thr Arg Pro Leu Asp Ala 710 715 Ser Asp Thr Leu Val Phe Asn Glu Val Phe Trp Val Ser Met Ser Tyr .730 725 Pro Ala Leu His Gln Lys Thr Leu Arg Val Asp Val Cys Thr Thr Asp 740 745 Arg Ser His Leu Glu Glu Cys Leu Gly Gly Ala Gln Ile Ser Leu Ala 760 Glu Val Cys Arg Ser Gly Glu Arg Ser Thr Arg Trp Tyr Asn Leu Leu 775 Ser Tyr Lys Tyr Leu Lys Lys Gln Ser Arg Glu Leu Lys Pro Val Gly 790 Val Met Ala Pro Ala Ser Gly Pro Ala Ser Thr Asp Ala Val Ser Ala 805 810 Leu Leu Glu Gln Thr Ala Val Glu Leu Glu Lys Arg Gln Glu Gly Arg 825 820 Ser Ser Thr Gln Thr Leu Glu Asp Ser Trp Arg Tyr Glu Glu Thr Ser 835 840 845 Glu Asn Glu Ala Val Ala Glu Glu Glu Glu Glu Glu Val Glu Glu Glu 855 Glu Gly Glu Glu Asp Val Phe Thr Glu Lys Ala Ser Pro Asp Met Asp 875 870 Gly Tyr Pro Ala Leu Lys Val Asp Lys Glu Thr Asn Thr Glu Thr Pro 885 890 Ala Pro Ser Pro Thr Val Val Arg Pro Lys Asp Arg Arg Val Gly Thr 900 -905 Pro Ser Gln Gly Pro Phe Leu Arg Gly Ser Thr Ile Ile Arg Ser Lys 920 Thr Phe Ser Pro Gly Pro Gln Ser Gln Tyr Val Cys Arg Leu Asn Arg 935 Ser Asp Ser Asp Ser Ser Thr Leu Ser Lys Lys Pro Pro Phe Val Arg 950 955 Asn Ser Leu Glu Arg Arg Ser Val Arg Met Lys Arg Pro Ser Ser Val

Lys Ser Leu Arg Ser Glu Arg Leu Ile Arg Thr Ser Leu Asp Leu Glu 980 985 990

Leu Asp Leu Gln Ala Thr Arg Thr Trp His Ser Gln Leu Thr Gln Glu 995  $\phantom{\bigg|}$  1000  $\phantom{\bigg|}$  1005

Ile Ser Val Leu Lys Glu Leu Lys Glu Gln Leu Glu Gln Ala Lys 1010 1015 1020

Ser His Gly Glu Lys Glu Leu Pro Gln Trp Leu Arg Glu Asp Glu 1025 1030 1035

Arg Phe Arg Leu Leu Arg Met Leu Glu Lys Arg Gln Met Asp 1040 1045 1050

Arg Ala Glu His Lys Gly Glu Leu Gln Thr Asp Lys Met Met Arg 1055 1060 1065

Ala Ala Lys Asp Val His Arg Leu Arg Gly Gln Ser Cys Lys 1070 1075 1080

Glu Pro Pro Glu Val Gln Ser Phe Arg Glu Lys Met Ala Phe Phe 1085  $\phantom{\bigg|}$  1090  $\phantom{\bigg|}$  1095

Thr Arg Pro Arg Met Asn Ile Pro Ala Leu Ser Ala Asp Asp Val 1100 1105 1110

<210> 15

<211> 1294

<212> DNA

<213> NM\_001306 CLDN3, claudin 3

<220>

<221> CDS

<222> (198)..(857)

<223>

<400> 15
caaagccaca ggcaggtgca ggcgcagccg cggcgagagc gtatggagcc gagccgttag
60
cgcggcgccgt cggtgagtca gtccgtccgt ccgtccgtcc gtcgggggcgc cgcagctccc
120
gccaggccca gcggccccgg cccctcgtct ccccgcaccc ggagccaccc ggtggagcgg
180
gcttgccgc ggcagcc atg tcc atg ggc ctg gag atc acg ggc acc gcg
Met Ser Met Gly Leu Glu Ile Thr Gly Thr Ala

	,				1			į	5					10		
						ctg Leu										278
						ttc Phe										326
	_				_	tgg Trp 50	_		_			-	_			374
cag Gln 60	atg Met	cag Gln	tgc Cys	aag Lys	gtg Val 65	tac Tyr	gac Asp	tcg Ser	ctg Leu	ctg Leu 70	gca Ala	ctg Leu	cca Pro	cag Gln	gac Asp 75	422
						ctc Leu										470
						ctg Leu										518
						gcc Ala										566
						ctc Leu 130										614
aac Asn 140	acc Thr	att Ile	atc Ile	cgg Arg	gac Asp 145	ttc Phe	tac Tyr	aac Asn	ccc Pro	gtg Val 150	gtg Val	ccc Pro	gag Glu	gcg Ala	cag Gln 155	662
						ggc										710
						gcg Ala										758
						acc Thr										806
						ctg Leu 210										854
gtc Val 220	taag	ggac	ag a	.cgca	.ggga	g ac	ccca	.ccac	cac	cacc	acc	acca	acac	ca		907
ccac	cacc	ac c	gcga	gctg	g ag	cgcg	cacc	agg	ccat	cca	gcgt	gcag	cc t	tgcc	tcgga	967
ggcc	agcc	ca c	cccc	agaa	g cc	agga	agcc	ccc	gcgc	tgg	actg	gggc	ag c	ttcc	ccagc	1027
agcc	acgg	ct t	tgcg	ggcc	g gg	cagt	cgac	ttc	aaaa	ccc	aggg	acca	ac c	tgca	tggac	1087
tgtg	aaac	ct c	accc	ttct	g ga	gcac	gggg	cct	gggt	gac	cgcc	aata	ct t	gacc	acccc	1147
gtcg	agcc	сс а	tcgg	gecg	c tg	cccc	catg	ctc	gcgc	tgg	gcag	ggac	cg g	cago	cctgg	1207

, aag	iggg (	act	tgat	attt	tt c	aata	aaag	c ct	ttcg	rttt	gca	aaaa	aaa	aaaa	aaaaaa
aaa	aaaa	aaa	aaaa	aaaa	aa a	aaaa.	aa								
<21	.0>	16				•			. ,			•		•	
<21	.1>	220							•						
<21	.2>	PRT													
<21	.3>	NM_C	0130	6 CL	DN3,	cla	udin	3				•			
							•		٠.						
<40	0>	16													
Met 1	Ser	Met	. Gly	Leu 5	Glu	Ile	Thr	Gly	Thr 10	Ala	Leu	Ala	Val	Leu 15	Glý
Trp	Leu	Gly	Thr 20	Ile	Val	Cys	Суз	Ala 25	Leu	Pro	Met	Trp	Arg 30	Val	Ser
Ala	Phe	Ile 35	Gly	Ser	Asn	Ile	Ile 40	Thr	Ser	Gln	Asn	Ile 45	Trp	Glu	Gly
Leu	Trp 50	Met	Asn	Cys	Val	Val 55	Gln	Ser	Thr	Gly	Gln 60	Met	Gln	Суз	Lys
Val 65	Tyr	Asp	Ser	Leu	Leu 70	Ala	Leu	Pro	Gln	Asp 75	Leu	Gln	Ala	Ala	Arg 80
Ala	Leu	Ile	Val	Val 85	Ala	Ile	Leu 	Leu	Ala 90	Ala	Phe	Gly	Leu	Leu 95	Val
Ala	Leu	Val	Gly 100	Ala	Gln	Cys	Thr	Asn 105	Суз	Val	Gln	Asp	Asp 110	Thr	Ala
Lys	Ala	Lys 115	Ile	Thr	Ile	Val	Ala 120	Gly	Val	Leu	Phe	Leu 125	Leu	Ala	Ala
Leu	Leu 130	Thr	Leu	Val	Pro	Val 135	Ser	Trp	Ser	Ala	Asn 140	Thr	Ile	Ile	Arg
Asp 145	Phe	Tyr	Asn	Pro	Val 150	Val	Pro	Glu	Ala	Gln 155	Lys	Arg	Glu	Met	Gly 160
Ala	Gly	Leu	Tyr	Val 165	Gly	Trp	Ala	Ala	Ala 170	Ala	Leu	Gln	Leu	Leu 175	Gly
Gly	Ala	Leu	Leu 180	Cys	Cys	Ser	Cys	Pro 185.		Arg	Glu :	Lys	Lys 190	Tyr	Thr
Ala	Thr	Lys 195	Val	Val	Tyr	Ser	Ala 200	Pro	Arg	Ser	Thr	Gly 205	Pro	Gly	Ala

Ser Leu Gly Thr Gly Tyr Asp Arg Lys Asp Tyr Val 210 215 220	
<210> 17	
<211> 1853	
<212> DNA	
<213> NM_022454 SOX17	
<220>	
<221> CDS	
<222> (205)(1446)	
<223>	
<400> 17	
gcagtgtcac taggccggct gggggccctg ggtacgctgt agaccagacc	60
agaacacggg cggcggcttc gggccgggag acccgcgcag ccctcggggc atctcagtgc	120
ctcactcccc accccctccc ccgggtcggg ggaggcggcg cgtccggcgg agggttgagg	180
ggagcggggc aggcctggag cgcc atg agc agc ccg gat gcg gga tac gcc Met Ser Ser Pro Asp Ala Gly Tyr Ala 1 5	231
agt gac gac cag agc cag acc cag agc gcg ctg ccc gcg gtg atg gcc Ser Asp Asp Gln Ser Gln Thr Gln Ser Ala Leu Pro Ala Val Met Ala 10 15 20 25	279
ggg ctg ggc ccc tgc ccc tgg gcc gag tcg ctg agc ccc atc ggg gac Gly Leu Gly Pro Cys Pro Trp Ala Glu Ser Leu Ser Pro Ile Gly Asp 30 35 40	327
atg aag gtg aag ggc gag gcg ccg gcg aac agc gga gca ccg gcc ggg Met Lys Val Lys Gly Glu Ala Pro Ala Asn Ser Gly Ala Pro Ala Gly 45 50 55	375
gcc gcg ggc cga gcc aag ggc gag tcc cgt atc cgg cgg ccg atg aac Ala Ala Gly Arg Ala Lys Gly Glu Ser Arg Ile Arg Arg Pro Met Asn 60 65 70	423
get ttc atg gtg tgg get aag gae gag ege aag egg etg geg eag eag	471
Ala Phe Met Val Trp Ala Lys Asp Glu Arg Lys Arg Leu Ala Gln Gln 75 80 85	
aat cca gac ctg cac aac gcc gag ttg agc aag atg ctg ggc aag tcg Asn Pro Asp Leu His Asn Ala Glu Leu Ser Lys Met Leu Gly Lys Ser 90 95 100 105	519
tgg aag gcg ctg acg ctg gcg gag aag cgg ccc ttc gtg gag gag gca Trp Lys Ala Leu Thr Leu Ala Glu Lys Arg Pro Phe Val Glu Glu Ala 110 115 120	567
gag cgg ctg cgc gtg cag cac atg cag gac cac ccc aac tac aag tac Glu Arg Leu Arg Val Gln His Met Gln Asp His Pro Asn Tyr Lys Tyr 125 130 135	615

				g Arg					Lys					Va.		ggc Gly	663
	Gl <sup>7</sup>	tto Phe	e Let	g cad	ggc Gly	ctg / Leu	get Ala 160	Glu	ccg Pro	cag Gln	g gcg Ala	g gcc Ala 165	a Ala	cto Lev	ı Gl	ccc Pro	711
	gag Glu 170	: Gl	Gly	cgc Arg	gtg Val	gcc Ala	Met	gac Asp	ggc	ctg Leu	ggc Gly	Let	caç Glr	tto Phe	ccc Pro	gag Glu 185	759
	cag Gln	GJ7	tto Phe	e ccc	gcc Ala 190	Gly	ccg Pro	ccg Pro	ctg Leu	ctg Leu 195	Pro	ccg Pro	cac His	ato Met	ggc : Gly 200	ggc	807
	cac	tac	cgc	gac	tgc:	cag	agt	ctg	ggc	gcg	cct	ccg	cto	gac	ggc	tac	855
	His	Туг	Arg	Asp 205		Gln	Ser	Leu	Gly 210		Pro	Pro	Leu	Asp 215	_	Tyr	
	ccg Pro	Leu	Pro 220	Thr	Pro	gac Asp	acg Thr	tcc Ser 225	Pro	ctg Leu	gac	ggc	gtg Val 230	Asp	ccc Pro	gac Asp	903
	ccg Pro	gct Ala 235	Phe	ttc Phe	gcc Ala	gcc Ala	ccg Pro 240	Met	ccc Pro	Gly	gac Asp	tgc Cys 245	Pro	gcg Ala	gcc Ala	ggc Gly	951
	acc Thr 250	Tyr	agc Ser	tac Tyr	gcg Ala	cag Gln 255	Val	tcg Ser	gac Asp	tac Tyr	gct Ala 260	Gly	ccc Pro	ccg Pro	gag Glu	cct Pro 265	999
	ccc Pro	gcc Ala	ggt Gly	ccc	atg Met 270	cac His	ccc Pro	cga Arg	ctc Leu	ggc Gly 275	cca Pro	gag Glu	ccc Pro	gcg Ala	ggt Gly 280	ccc Pro	1047
•	tcg Ser	att Ile	ccg Pro	ggc Gly 285	ctc Leu	ctg Leu	gcg Ala	cca Pro	ccc Pro 290	agc Ser	gcc Ala	ctt Leu	cac His	gtg Val 295	tac Tyr	tac Tyr	1095
	ggc Gly	gcġ Ala	atg Met 300	Gly	tcg Ser	ccc Pro	GJ À āāā	gcg Ala 305	ggc	ggc Gly	Gly ggg	cgc Arg	ggc Gly 310	ttc Phe	cag Gln	atg Met	1143
	cag Gln	ccg Pro 315	caa Gln	cac His	cag Gln	cac His	cag Gln 320	cac His	cag Gln	cac His	cag Gln	cac His 325	cac His	ccc Pro	ccg Pro	ggc Gly	1191
	ccc	gga	cag	ccg	tcg	ccc	cct	ccg	gag	gca	ctg	ccc	tgc	cgg	gac	ggc	1239
	Pro 330	Gly	Gln	Pro	Ser	Pro 335	Pro	Pro	Glu	Ala	Leu 340	Pro	Cys	Arg	Asp	Gly 345	•
	acg Thr	gac Asp	ccc Pro	agt Ser	cag Gln 350	ccc Pro	gcc Ala	gag Glu	ctc Leu	ctc Leu 355	G17 ggg	gag Glu	gtg Val	gac Asp	cgc Arg 360	acg Thr	1287
	gaa Glu	ttt Phe	gaa Glu	cag Gln 365	tat Tyr	ctg Leu	cac His	Phe	gtg Val 370	tgc Cys	aag Lys	cct Pro	gag Glu	atg Met 375	ggc Gly	ctc Leu	1335
	ccc Pro	tac Tyr	cag Gln 380	GJ Å āāā	cat His	gac Asp	Ser	ggt Gly 385	gtg Val	aat Asn	ctc Leu	Pro	gac Asp 390	agc Ser	cac His	ggg Gly	. 1383
	gcc	att	tcc	tcg	gtg	gtg	tcc	gac	gcc	agc	tcc	gcg	gta	tat	tac	tgc	1431

Ala Ile Ser Ser Val Val Ser Asp Ala Ser Ser Ala Val Tyr Tyr Cys 395 400 405
aac tat cct gac gtg tgacaggtcc ctgatccgcc ccagcctgca ggccagaagc Asn Tyr Pro Asp Val 410
agtgttacac acttcctgga ggagctaagg aaatcctcag actcctgggt ttttgttgtt
gctgttgttg ttttttaaaa ggtgtgttgg catataattt atggtaattt attttgtctg
ccacttgaac agtttggggg ggtgaggttt catttaaaat ttgttcagag atttgtttcc
catagttgga ttgtcaaaac cctatttcca agttcaagtt aactagcttt gaatgtgtcc
caaaacagct teetecattt eetgaaagtt tattgateaa agaaatgttg teetgggtgt
gttttttcaa tottotaaaa aataaaatot ggaatootga aaaaaaaaaa
aaaaaaa
<210> 18
<211> 414
<212> PRT
<213> NM_022454 SOX17
<400> 18
Met Ser Ser Pro Asp Ala Gly Tyr Ala Ser Asp Asp Gln Ser Gln Thr 1 5 10 15
Gln Ser Ala Leu Pro Ala Val Met Ala Gly Leu Gly Pro Cys Pro Trp 20 25 30
Ala Glu Ser Leu Ser Pro Ile Gly Asp Met Lys Val Lys Gly Glu Ala 35 40 45
Pro Ala Asn Ser Gly Ala Pro Ala Gly Ala Ala Gly Arg Ala Lys Gly 50 55 60
Glu Ser Arg Ile Arg Pro Met Asn Ala Phe Met Val Trp Ala Lys 65 70 75 80
Asp Glu Arg Lys Arg Leu Ala Gln Gln Asn Pro Asp Leu His Asn Ala 85 90 95
Glu Leu Ser Lys Met Leu Gly Lys Ser Trp Lys Ala Leu Thr Leu Ala 100 105 110
Glu Lys Arg Pro Phe Val Glu Glu Ala Glu Arg Leu Arg Val Gln His 115 120 125

Met Gln Asp His Pro Asn Tyr Lys Tyr Arg Pro Arg Arg Lys Gln 130 135 140

Val Lys Arg Leu Lys Arg Val Glu Gly Gly Phe Leu His Gly Leu Ala 145 155 160

Glu Pro Gln Ala Ala Ala Leu Gly Pro Glu Gly Gly Arg Val Ala Met 165 170 175

Asp Gly Leu Gly Leu Gln Phe Pro Glu Gln Gly Phe Pro Ala Gly Pro 180 185 190

Pro Leu Leu Pro Pro His Met Gly Gly His Tyr Arg Asp Cys Gln Ser 195 200 205

Leu Gly Ala Pro Pro Leu Asp Gly Tyr Pro Leu Pro Thr Pro Asp Thr 210 215 220

Ser Pro Leu Asp Gly Val Asp Pro Asp Pro Ala Phe Phe Ala Ala Pro 225 230 235 240

Met Pro Gly Asp Cys Pro Ala Ala Gly Thr Tyr Ser Tyr Ala Gln Val 245 250 255

Ser Asp Tyr Ala Gly Pro Pro Glu Pro Pro Ala Gly Pro Met His Pro 260 265 270

Arg Leu Gly Pro Glu Pro Ala Gly Pro Ser Ile Pro Gly Leu Leu Ala 275 280 285

Pro Pro Ser Ala Leu His Val Tyr Tyr Gly Ala Met Gly Ser Pro Gly 290 295 300

Ala Gly Gly Gly Arg Gly Phe Gln Met Gln Pro Gln His Gln His Gln 305 310 315 320

His Gln His Gln His His Pro Pro Gly Pro Gly Gln Pro Ser Pro Pro 325 330 335

Pro Glu Ala Leu Pro Cys Arg Asp Gly Thr Asp Pro Ser Gln Pro Ala 340 345 350

Glu Leu Leu Gly Glu Val Asp Arg Thr Glu Phe Glu Gln Tyr Leu His 355 360 365

Phe Val Cys Lys Pro Glu Met Gly Leu Pro Tyr Gln Gly His Asp Ser 370 375 380

Gly Val Asn Leu Pro Asp Ser His Gly Ala Ile Ser Ser Val Val Ser 385 390 395 400

Asp Ala Ser Ser Ala Val Tyr Tyr Cys Asn Tyr Pro Asp Val

<210>	19												
<211>	3702							•					
<212>	DNA												•
<213>	им_00	5682 GF	PR56										
				•									
<220>													
<221>	CDS												
<222>	(159)	(2237	)										
<223>								٠					
			ē										
<400>	19	actotat	a	aaata		+	~+~~						60
acctcca		gctctgt											. 60
													120
caggtg	giga ci	LCCdag	ag tga	cceg	e gg	agga		et T				er Leu	176
ctg caq Leu Glr	Thr I	ica ctg Thr Leu 10	ttc c	g cto eu Lei	agt Ser 15	ctg Leu	ctc Leu	ttc Phe	ctg Leu	gtc Val 20	caa Gln	ggt Gly	224
gcc cad Ala His	ggc a Gly A 25	agg ggc Arg Gly	cac ac	gg gaa g Glu 30	gac Asp	ttt Phe	cgc Arg	ttc Phe	tgc Cys 35	agc Ser	cag Gln	cgg Arg	272
aac cac Asn Glr 40	g aca c n Thr H	ac agg His Arg	agc ag Ser Se 45	er Leu	cac His	tac Tyr	aaa Lys	ccc Pro 50	aca Thr	cca Pro	gac Asp	ctg Leu	320
cgc atc Arg Ile 55	tcc a Ser I	itc gag :le Glu	aac to Asn Se 60	c gaa r Glu	gag Glu	gcc Ala	ctc Leu 65	aca Thr	gtc Val	cat His	gcc Ala	cct Pro 70	368
ttc cct Phe Pro					Arg	Ser							416
ctc tac Leu Tyr	His P	tc tgc he Cys 0	ctc ta Leu Ty	c tgg r Trp	aac Asn 95	cga Arg	cat His	gct Ala	ej gaa	aga Arg 100	tta Leu	cat His	464
ctt ctc Leu Leu	tat g Tyr G 105	gc aag ly Lys	cgt ga Arg As	c ttc p Phe 110	ttg Leu	ctg Leu	agt Ser	gac Asp	aaa Lys 115	gcc Ala	tct Ser	agc Ser	512
ctc ctc Leu Leu 120	Cys P	tc cag he Gln	cac ca His Gl 12	n Glu	gag Glu	agc Ser	ctg Leu	gct Ala 130	Gln	ggc Gly	ccc Pro	ccg Pro	560
ctg tta Leu Leu 135	gcc a	ct tct hr Ser	gtc ac Val Th 140	c tcc r Ser	tgg Trp	tgg Trp	agc Ser 145	cct Pro	cag Gln	aac Asn	atc Ile	agc Ser 150	608
ctg ccc	agt go	cc gcc	agc tt	c acc	ttc	tee	ttc	cac	agt	cct	ccc	cac	656

Leu	Pro	Ser	Ala	Ala 155	Ser	Phe	Thr	Phe	Ser 160	Phe	His	Ser	Pro	Pro 165	His	
				aat Asn												704
	_	_		agc Ser	_		_	-			_	_	_			. 752
				gcc Ala												800
				gtg Val												84,8
				gcc Ala 235												896
_	_	_		atc Ile				-				_	-			944
				gtg Val												992
				gag Glu												1040
				ttc Phe												1088
				att Ile 315						Lys						1136
-				ctc Leu			_		_		_	_	_			1184
	_		_	gtg Val			_	-	-			_				1232
				agt Ser												1280
				tgc Cys												1328
				gtg Val 395												1376
				tgt Cys												1424
gcc	gcc	tac	ctc	tgc	tcc	agg	gtg	ccc	ctg	ccg	tgc	agg	agg	aaa	cct	1472

Ala	Ala	Tyr 425	Leu	Суз	Ser	Arg	Val 430	Pro	Leu	Pro	Суз	Arg 435	Arg	Lys	Pro	•
														gtc Val		1520
ctg Leu 455	ctg Leu	gac Asp	acg Thr	agc Ser	ttc Phe 460	ctg Leu	ctc Leu	agc Ser	gag Glu	ccg Pro 465	gtg Val	gcc Ala	ctg Leu	aca Thr	ggc Gly 470	1568
tct Ser	gag Glu	gct Ala	ggc	tgc Cys 475	cga Arg	gcc Ala	agt Ser	gcc Ala	atc Ile 480	ttc Phe	ctg Leu	cac His	ttc Phe	tcc Ser 485	ctg Leu	1616
		-				_								tac Tyr		1664
														ctc Leu		1712
														ctg Leu		1760
														gtg Val		1808
														cgg Arg 565		1856
														gtg Val		1904
-			_	_	_				_		-	_		ctg Leu		1952
														ctg Leu		2000
ctc Leu 615	agc Ser	ctg Leu	gtc Val	ctt Leu	ggc Gly 620	ctg Leu	ccc Pro	tgg Trp	gcc Ala	ttg Leu 625	atc Ile	ttc Phe	ttc Phe	tcc Ser	ttt Phe 630	2048
														atc Ile 645		2096
														atg Met		2144
														gac Asp		2192
gcc Ala	agg Arg 680	ctc Leu	ccc Pro	atc Ile	agc Ser	tcg Ser 685	ggc Gly	agc Ser	acc Thr	tcg Ser	tcc Ser 690	agc Ser	cgc Arg	atc Ile		2237
tago	jecto	ca g	ICCCa	cctg	ic co	atgt	gato	g aag	gcaga	gat	gcgg	ccto	gt d	gcac	actgc	2297

ctgtggcccc	cgagccaggc	ccagccccag	gccagtcagċ	cgcagacttt	ggaaagccca	2357
acgaccatgg	agagatgggc	cgttgccatg	gtggacggac	tcccgggctg	ggcttttgaa	2417
ttggccttgg	ggactactcg	gctctcactc	agctcccacg	ggactcagaa	gtgcgccgcc	2477
atgetgeeta	gggtactgtc	cccacatctg	tcccaaccca	gctggaggcc	tggtctctcc	2537
ttacaaaccc	tgggcccagc	cctcattgct	gggggccagg	ccttggatct	tgagggtctg	2597
gcacatcctt	aatcctgtgc	ccctgcctgg	gacagaaatg	tggctccagt	tgctctgtct	2657
ctcgtggtca	ccctgagggc	actctgcatc	ctctgtcatt	ttaacctcag	gtggcaccca	2717
gggcgaatgg	ggcccagggc	agaccttcag	ggccagagcc	ctggcggagg	agaggccctt	2777
tgccaggagc	acagcagcag	ctcgcctacc	tetgagecca	ggccccctcc	ctccctcagc	2837
ccccagtcc	tccctccatc	ttccctgggg	ttetectect	ctcccagggc	ctccttgctc	2897
cttcgttcac	agctgggggt	ccccgattcc	aatgctgttt	tttggggagt	ggtttccagg	2957
agctgcctgg	tgtctgctgt	aaatgtttgt	ctactgcaca	agcctcggcc	tgcccctgag	3017
ccaggctcgg	taccgatgcg	tgggctgggc	taggtccctc	tgtccatctg	ggcctttgta	3077
tgagctgcat	tgcccttgct	caccctgacc	aagcacacgc	ctcagagggg	ccctcagcct	3137
ctcctgaagc	cctcttgtgg	caagaactgt	ggaccatgcc	agtcccgtct	ggtttccatc	3197
ccaccactcc	aaggactgag	actgacctcc	tctggtgaca	ctggcctaga	gcctgacact	3257
ctcctaagag	gttctctcca	agcccccaaa	tagctccagg	cgccctcggc	cgcccatcat	3317
ggttaattct	gtccaacaaa	cacacacggg	tagattgctg	gcctgttgta	ggtggtaggg	337
acacagatga	ccgacctggt	cactcctcct	gccaacattc	agtctggtat	gtgaggcgtg	343
cgtgaagcaa	gaactcctgg	agctacaggg	acagggagcc	atcattcctg	cctgggaatc	349
ctggaagact	tcctgcagga	gtcagcgttc	aatcttgacc	ttgaagatgg	gaaggatgtt	355
ctttttacgt	accaattctt	ttgtcttttg	atattaaaaa	gaagtacatg	ttcattgtag	361
agaatttgga	aactgtagaa	gagaatcaag	aagaaaaata	aaaatcagct	gttgtaatcg	367
cctagcaaaa	aaaaaaaaa	aaaaa				370
<210> 20						

<211> 693

<212> PRT

<213> NM\_005682 GPR56

<400> 20

Met Thr Pro Gln Ser Leu Leu Gln Thr Thr Leu Phe Leu Leu Ser Leu 15

Leu Phe Leu Val Gln Gly Ala His Gly Arg Gly His Arg Glu Asp Phe

20 25 Arg Phe Cys Ser Gln Arg Asn Gln Thr His Arg Ser Ser Leu His Tyr Lys Pro Thr Pro Asp Leu Arg Ile Ser Ile Glu Asn Ser Glu Glu Ala Leu Thr Val His Ala Pro Phe Pro Ala Ala His Pro Ala Ser Arg Ser 70 . 75 Phe Pro Asp Pro Arg Gly Leu Tyr His Phe Cys Leu Tyr Trp Asn Arg 85 His Ala Gly Arg Leu His Leu Leu Tyr Gly Lys Arg Asp Phe Leu Leu 105 Ser Asp Lys Ala Ser Ser Leu Leu Cys Phe Gln His Gln Glu Glu Ser 120 Leu Ala Gln Gly Pro Pro Leu Leu Ala Thr Ser Val Thr Ser Trp Trp 135 Ser Pro Gln Asn Ile Ser Leu Pro Ser Ala Ala Ser Phe Thr Phe Ser 145 150 Phe His Ser Pro Pro His Thr Ala Ala His Asn Ala Ser Val Asp Met 170 Cys Glu Leu Lys Arg Asp Leu Gln Leu Leu Ser Gln Phe Leu Lys His. 185 Pro Gln Lys Ala Ser Arg Arg Pro Ser Ala Ala Pro Ala Ser Gln Gln 200 Leu Gln Ser Leu Glu Ser Lys Leu Thr Ser Val Arg Phe Met Gly Asp 210 215 220 Met Val Ser Phe Glu Glu Asp Arg Ile Asn Ala Thr Val Trp Lys Leu 230 Gln Pro Thr Ala Gly Leu Gln Asp Leu His Ile His Ser Arg Gln Glu Glu Glu Gln Ser Glu Ile Met Glu Tyr Ser Val Leu Leu Pro Arg Thr 265

Leu Phe Gln Arg Thr Lys Gly Arg Ser Gly Glu Ala Glu Lys Arg Leu

Leu Leu Val Asp Phe Ser Ser Gln Ala Leu Phe Gln Asp Lys Asn Ser

280

A 100 to 1

295 300 290 Ser His Val Leu Gly Glu Lys Val Leu Gly Ile Val Val Gln Asn Thr 315 310 Lys Val Ala Asn Leu Thr Glu Pro Val Val Leu Thr Phe Gln His Gln 325 330 Leu Gln Pro Lys Asn Val Thr Leu Gln Cys Val Phe Trp Val Glu Asp 340 345 Pro Thr Leu Ser Ser Pro Gly His Trp Ser Ser Ala Gly Cys Glu Thr 360 Val Arg Arg Glu Thr Gln Thr Ser Cys Phe Cys Asn His Leu Thr Tyr 375 Phe Ala Val Leu Met Val Ser Ser Val Glu Val Asp Ala Val His Lys 390 His Tyr Leu Ser Leu Leu Ser Tyr Val Gly Cys Val Val Ser Ala Leu 405 410 Ala Cys Leu Val Thr Ile Ala Ala Tyr Leu Cys Ser Arg Val Pro Leu 425 Pro Cys Arg Arg Lys Pro Arg Asp Tyr Thr Ile Lys Val His Met Asn 440 Leu Leu Leu Ala Val Phe Leu Leu Asp Thr Ser Phe Leu Leu Ser Glu 455 460 Pro Val Ala Leu Thr Gly Ser Glu Ala Gly Cys Arg Ala Ser Ala Ile 470 475 Phe Leu His Phe Ser Leu Leu Thr Cys Leu Ser Trp Met Gly Leu Glu 485 490 Gly Tyr Asn Leu Tyr Arg Leu Val Val Glu Val Phe Gly Thr Tyr Val 505 Pro Gly Tyr Leu Leu Lys Leu Ser Ala Met Gly Trp Gly Phe Pro Ile 520 Phe Leu Val Thr Leu Val Ala Leu Val Asp Val Asp Asn Tyr Gly Pro 535 Ile Ile Leu Ala Val His Arg Thr Pro Glu Gly Val Ile Tyr Pro Ser 550 .. 555

Met Cys Trp Ile Arg Asp Ser Leu Val Ser Tyr Ile Thr Asn Leu Gly
565 570 575

Leu Phe Ser Leu Val Phe Leu Phe Asn Met Ala Met Leu Ala Thr Met 580 590

Val Val Gln Ile Leu Arg Leu Arg Pro His Thr Gln Lys Trp Ser His 595 600 605

Val Leu Thr Leu Leu Gly Leu Ser Leu Val Leu Gly Leu Pro Trp Ala 610 615 620

Leu Ile Phe Phe Ser Phe Ala Ser Gly Thr Phe Gln Leu Val Val Leu 625 630 635 640

Tyr Leu Phe Ser Ile Ile Thr Ser Phe Gln Gly Phe Leu Ile Phe Ile 645 650 655

Trp Tyr Trp Ser Met Arg Leu Gln Ala Arg Gly Gly Pro Ser Pro Leu 660 665 670

Lys Ser Asn Ser Asp Ser Ala Arg Leu Pro Ile Ser Ser Gly Ser Thr 675 680 . 685

Ser Ser Ser Arg Ile 690

<210> 21

<211> 1332

<212> DNA

<213> NM\_001307 claudin 7, CLDN7

<220>

<221> CDS

<222> (427)..(1059)

<223>

<400> 21
ctgctggctc acctccgagc cacctctgct gcgcaccgca gcctcggacc tacagcccag 60
gatactttgg gacttgccgg cgctcagaaa cgcgcccaga cggcccctcc accttttgtt 120
tgcctagggt cgccgagagc gcccggaggg aaccgcctgg ccttcgggga ccaccaattt 180
tgtctggaac caccctcccg gcgtatccta ctccctgtgc cgcgaggcca tcgcttcact 240
ggaggggtcg atttgtgtg agtttggtga caagatttgc attcacctgg cccaaaccct 300
ttttgtctct ttgggtgacc ggaaaactcc acctcaagtt ttctttgtg gggctgcccc 360

ccaa	gtgt	cg t	tttgt	ttta	ic tg	tagg	gtct	ccc	gee	ggc	gccc	cca	gtg t	ttt	ctgagg	420
gcgg	raa a N	1et 1	gcc a Ala A	at t Sn S	cg g Ser G	gc c ly I	tg c	ag t	tg d Leu I	Seu (	gc t Sly E LO	tc t	tcc a Ser N	atg ( Met .	gcc Ala	468
ctg Leu 15	ctg Leu	ggc Gly	tgg Trp	gtg Val	ggt Gly 20	ctg Leu	gtg Val	gcc Ala	tgc Cys	acc Thr 25	gcc Ala	atc Ile	ccg Pro	cag Gln	tgg Trp 30	51 <sub>.</sub> 6
cag Gln	atg Met	agc Ser	tcc Ser	tat Tyr 35	gcg Ala	ggt Gly	gac Asp	aac Asn	atc Ile 40	atc Ile	acg Thr	gcc Ala	cag Gln	gcc Ala 45	atg Met	564
tac Tyr	aag Lys	Gly	ctg Leu 50	tgg Trp	atg Met	gac Asp	tgc Cys	gtc Val 55	acg Thr	cag Gln	agc Ser	acg Thr	ggg Gly 60	atg Met	atg Met	612
agc Ser	tgc Cys	aaa Lys 65	atg Met	tac Tyr	gac Asp	tcg Ser	gtg Val 70	ctc Leu	gcc Ala	ctg Leu	tcc Ser	gcg Ala 75	gcc Ala	ttg Leu	cag Gln	660
gcc Ala	act Thr 80	cga Arg	gcc Ala	cta Leu	atg Met	gtg Val 85	gtc Val	tcc Ser	ctg Leu	gtg Val	ctg Leu 90	Gly	ttc Phe	ctg Leu	gcc Ala	708
atg Met 95	ttt Phe	gtg Val	gcc Ala	acg Thr	atg Met 100	G] À dàc	atg Met	aag Lys	tgc Cys	acg Thr 105	cgc Arg	tgt Cys	G1y ggg	gga Gly	gac Asp 110	756
gac Asp	aaa Lys	gtg Val	aag Lys	aag Lys 115	gcc Ala	cgt Arg	ata Ile	gcc Ala	atg Met 120	Gly	gga Gly	Gly	ata Ile	att Ile 125	ttc Phe	804
atc Ile	gtg Val	gca Ala	ggt Gly 130	ctt Leu	gcc Ala	acc Thr	ttg Leu	gta Val 135	Ala	tgc Cys	tcc Ser	tgg Trp	tat Tyr 140	GT	cat His	852
cag Gln	att	gto Val 145	Thr	gac Asp	ttt Phe	tat Tyr	aac Asn 150	cct Pro	ttg Leu	atc Ile	cct Pro	acc Thr 155	Asn	att	aag Lys	900
tat Tyr	gag Glu 160	Phe	ggc Gly	cct Pro	gcc Ala	atc Ile 165	ttt Phe	att Ile	ggc	tgg Trp	gca Ala 170	Gly	tct Ser	gcc	cta Leu	948
gtc Val 175	Ile	ctg Leu	gga Gly	ggt Gly	gca Ala 180	Leu	ctc Leu	tcc Ser	tgt Cys	Ser 185	Cys	Pro	Gly Gly	aat Asr	gag Glu 190	996
agc Ser	aag Lys	gct	Gly ggg	tac Tyr 195	Arg	gca Ala	ccc Pro	cgc	Ser 200	Tyr	cct Pro	aaq Lys	tcc Ser	aac Asr 205	tct Ser	1044
tcc Ser	aag Lys	gaç	tat Tyr 210	Val	tga	cctg	gga	tctc	cttg	ec c	cago	ctga	ıc ag	gcta	atggg	1099
agt	gtct	aga	tgcc	tgaa	ag g	gcct	gggg	c tg	agct	cago	ctg	rtggg	gcag	ggt	gccggac	1159
aaa	ggcc	tcc	tggt	cact	ct g	tece	tgca	c to	catg	rtata	gto	ctct	tgg	gtt	gggggtg	1219
ggg	gggt	gcc	gttg	gtgg	ga g	agac	aaaa	a ga	ggga	gagt	gtg	cttt	ttg	taca	agtaata	1279
aaa	aata	agt	attg	ggaa	aa a	aaaa	aaaa	a aa	aaaa	aaaa	aaa	aaaa	aaa	aaa		1332

<210> 22

<211> 211

<212> PRT

<213> NM\_001307 claudin 7, CLDN7

<400> 22

Met Ala Asn Ser Gly Leu Gln Leu Leu Gly Phe Ser Met Ala Leu Leu 1 5 10 15

Gly Trp Val Gly Leu Val Ala Cys Thr Ala Ile Pro Gln Trp Gln Met 20 25 30

Ser Ser Tyr Ala Gly Asp Asn Ile Ile Thr Ala Gln Ala Met Tyr Lys 35 40

Gly Leu Trp Met Asp Cys Val Thr Gln Ser Thr Gly Met Met Ser Cys 50 55 60

Lys Met Tyr Asp Ser Val Leu Ala Leu Ser Ala Ala Leu Gln Ala Thr 65 70 75 80

Arg Ala Leu Met Val Val Ser Leu Val Leu Gly Phe Leu Ala Met Phe 85 90 95

Val Ala Thr Met Gly Met Lys Cys Thr Arg Cys Gly Gly Asp Asp Lys 100 105 110

Val Lys Lys Ala Arg Ile Ala Met Gly Gly Gly Ile Ile Phe Ile Val 115 120 125

Ala Gly Leu Ala Thr Leu Val Ala Cys Ser Trp Tyr Gly His Gln Ile 130 135 140

Val Thr Asp Phe Tyr Asn Pro Leu Ile Pro Thr Asn Ile Lys Tyr Glu 145 150 155 160

Phe Gly Pro Ala Ile Phe Ile Gly Trp Ala Gly Ser Ala Leu Val Ile 165 170 175

Leu Gly Gly Ala Leu Leu Ser Cys Ser Cys Pro Gly Asn Glu Ser Lys 180 185 190

Ala Gly Tyr Arg Ala Pro Arg Ser Tyr Pro Lys Ser Asn Ser Ser Lys 195 200 205

Glu Tyr Val 210

ааааааааа аааааааааа а

888

52

<210> 24

<211> 111

<212> PRT

<213> NM\_014736 KIAA0101 gene product .

<400> 24

Met Val Arg Thr Lys Ala Asp Ser Val Pro Gly Thr Tyr Arg Lys Val

Val Ala Ala Arg Ala Pro Arg Lys Val Leu Gly Ser Ser Thr Ser Ala 20 25 30

Thr Asn Ser Thr Ser Val Ser Ser Arg Lys Ala Glu Asn Lys Tyr Ala 35 40 45

Gly Gly Asn Pro Val Cys Val Arg Pro Thr Pro Lys Trp Gln Lys Gly 50 55 60

Ile Gly Glu Phe Phe Arg Leu Ser Pro Lys Asp Ser Glu Lys Glu Asn 65 70 75 80

Gln Ile Pro Glu Glu Ala Gly Ser Ser Gly Leu Gly Lys Ala Lys Arg 85 90 95

Lys Ala Cys Pro Leu Gln Pro Asp His Thr Asn Asp Glu Lys Glu 100 105 110

<210> 25

<211> 598

<212> DNA

<213> NM\_003064 secretory leukocyte protease inhibitor, SLPI

<220>

<221> CDS

<222> (23)..(418)

<223>

<400> 25
cagagtcact cctgccttca cc atg aag tcc agc ggc ctc ttc ccc ttc ctg
Met Lys Ser Ser Gly Leu Phe Pro Phe Leu
10

gtg Val	ctg Leu	ctt Leu	gcc Ala	ctg Leu 15	gga Gly	act Thr	ctg Leu	gca Ala	cct Pro 20	tgg Trp	gct Ala	gtg Val	GTU	ggc Gly 25	tct Ser	100
gga Gly	aag Lys	tcc Ser	ttc Phe 30	aaa Lys	gct Ala	gga Gly	gtc Val	tgt Cys 35	cct Pro	cct Pro	aag Lys	aaa Lys	tct Ser 40	gcc Ala	cag Gln	148
tgc Cys	ctt Leu	aga Arg 45	tac Tyr	aag Lys	aaa Lys	cct Pro	gag Glu 50	tgc Cys	cag Gln	agt Ser	gac Asp	tgg Trp 55	cag Gln	tgt Cys	cca Pro	196
Gly ggg	aag Lys 60	aag Lys	aga Arg	tgt Cys	tgt Cys	cct Pro 65	gac Asp	act Thr	tgt Cys	GJ y ggc	atc Ile 70	aaa Lys	tgc Cys	ctg Leu	gat Asp	244
cct Pro 75	gtt Val	gac Asp	acc Thr	cca Pro	aac Asn 80	cca Pro	aca Thr	agg Arg	agg Arg	aag Lys 85	cct	ggg	aag Lys	tgc Cys	cca Pro 90	292
gtg Val	act Thr	tat Tyr	ggc Gly	caa Gln 95	tgt Cys	ttg Leu	atg Met	ctt Leu	aac Asn 100	Pro	ccc Pro	aat Asn	ttc Phe	tgt Cys 105	gag Glu	340
atg Met	gat Asp	ggc Gly	cag Gln 110	Cys	aag Lys	cgt Arg	gac Asp	ttg Leu 115	Lys	tgt Cys	tgc Cys	atg Met	ggc Gly 120	Met	tgt Cys	388
GJ A aaa	aaa Lys	tcc Ser	Cys	gtt Val	tcc Ser	cct Pro	gtg Val 130	Lys	gct Ala	tga	ttcc	tgc	cata	tgga	gg	438
agg	ctct			tact	.ct q	tata	gtco	a gg	tect	ttcc	aco	ctga	igac	ttgg	ctccac	498
															gccagt	558
					ac g							•				598
									•							
<21	.0>	26													•	
<21	.1>	132			:					•	٠	•				
<21		PRT														
<21	.3>	им_0	0306	54 se	ecret	ory	leu	cocyt	e pi	rotea	ase :	inhil	oitor	, SI	PLI	
<4(	00>	26														
Met · 1	: Lys	s Sei	c Sei	c Gly 5	, Let	ı Phe	e Pro	) Phe	e Let 10	ı Val	l Le	ı Let	ı Ala	1 Le:	ı Gly	
Thi	r Let	ı Ala	a Pro	o Trj	o Ala	a Val	L Gla	a Gly 25	y Se:	r Gl	у Гу	s Se:	r Phe	e Ly:	s Ala	
Gl	y Val	1 Cy: 35	s Pro	o Pr	o Lys	i Ly:	s Se:	r Ala	a Gl	n Cy:	s Le	u Arg	g Ty:	r Ly:	s Lys	
Pro	50 50	u Cy:	s Gl	n Se	r Ası	55	o Gl	n Cy	s Pr	o Gl	60 Y Ly	s Ly	s Aro	g Cy	s Cys	

546

Pro Asp Thr Cys Gly Ile Lys Cys Leu Asp Pro Val Asp Thr Pro Asn Pro Thr Arg Arg Lys Pro Gly Lys Cys Pro Val Thr Tyr Gly Gln Cys 85 Leu Met Leu Asn Pro Pro Asn Phe Cys Glu Met Asp Gly Gln Cys Lys 105 . 100 Arg Asp Leu Lys Cys Cys Met Gly Met Cys Gly Lys Ser Cys Val Ser . 125 120 115 Pro Val Lys Ala 130 <210> 27 <211> 3970 <212> DNA <213> NM 013994 DDR1 <220> <221> CDS (337)..(3093) <222> <223> <400> 27 ggcttaggaa gtattaactg atctctgccc tagttctcat gtgttaaata tggatagtaa 60 tagtatctac cttatgaagt gactgtgaag ataaaattat ggaftctgtt taagggttta 120 180 ggccagtgtc tggcacaggg gaagcattct aaaaatatag ctgatgctgt taaacaatga ctgttgttgt tgttttactg ttattatccc caaagcggcc cattetgtct gttgctgtca 240 300 gctatgactc agtcccctga ttaacttacg caccacccat tttatcccct gcagagatgc 354 tgcccccacc cccttaggcc cgagggatca ggagct atg gga cca gag gcc ctg Met Gly Pro Glu Ala Leu 402 tca tct tta ctg ctg ctc ttg gtg gca agt gga gat gct gac atg Ser Ser Leu Leu Leu Leu Leu Val Ala Ser Gly Asp Ala Asp Met 450 aag gga cat ttt gat cct gcc aag tgc cgc tat gcc ctg ggc atg cag Lys Gly His Phe Asp Pro Ala Lys Cys Arg Tyr Ala Leu Gly Met Gln

30

45

gac cgg acc atc cca gac agt gac atc tct gct tcc agc tcc tgg tca Asp Arg Thr Ile Pro Asp Ser Asp Ile Ser Ala Ser Ser Ser Trp Ser

gat tee act gee gee ege cac age agg ttg gag age agt gae ggg gat

Asp 55	Ser	Thr	Ala	Ala	Arg 60	His	Ser	Arg	Leu	Glu 65	Ser	Ser	Asp	Gly	Asp 70		
Gly	gcc Ala	tgg Trp	tgc Cys	ccc Pro 75	gca Ala	ejā āāā	tcg Ser	gtg Val	ttt Phe 80	ccc Pro	aag Lys	gag Glu	gag Glu	gag Glu 85	tac Tyr		594
ttg Leu	cag Gln	gtg Val	gat Asp 90	cta Leu	caa Gln	cga Arg	ctg Leu	cac His 95	ctg Leu	gtg Val	gct Ala	ctg Leu	gtg Val 100	ggc Gly	acc Thr		642
cag Gln	gga Gly	cgg Arg 105	cat His	gcc Ala	GJA GGA	ggc Gly	ctg Leu 110	ggc Gly	aag Lys	gag Glu	ttc Phe	tcc Ser 115	cgg Arg	agc Ser	tac Tyr		690
cgg Arg	ctg Leu 120	cgt Arg	tac Tyr	tcc Ser	cgg Arg	gat Asp 125	ggt Gly	cgc Arg	cgc Arg	tgg Trp	atg Met 130	ggc Gly	tgg Trp	aag Lys	gac Asp		738
cgc Arg 135	tgg Trp	ggt Gly	cag Gln	gag Glu	gtg Val 140	atc Ile	tca Ser	ggc Gly	aat Asn	gag Glu 145	gac Asp	cct Pro	gag Glu	gga Gly	gtg Val 150		786
gtg Val	ctg Leu	aag Lys	gac Asp	ctt Leu 155	Gly ggg	ccc Pro	ccç Pro	atg Met	gtt Val 160	gcc Ala	cga Arg	ctg Leu	gtt Val	cgc Arg 165	ttc Phe		834
tac Tyr	ccc Pro	cgg Arg	gct Ala 170	gac Asp	cgg Arg	gtc Val	atg Met	agc Ser 175	gtc Val	tgt Cys	ctg Leu	cgg Arg	gta Val 180	gag Glu	ctc Leu		882
tat Tyr	Gly	tgc Cys 185	ctc Leu	tgg Trp	agg Arg	gat Asp	gga Gly 190	ctc Leu	ctg. Leu	tct Ser	tac Tyr	acc Thr 195	gcc Ala	cct Pro	gtg Val		930
GJ Y GGG	cag Gln 200	aca Thr	atg Met	tat Tyr	tta Leu	tct Ser 205	gag Glu	gcc Ala	gtg Val	tac Tyr	ctc Leu 210	aac Asn	gac Asp	tcc Ser	acc Thr		978
tat Tyr 215	Asp	gga Gly	cat His	acc Thr	gtg Val 220	ggc Gly	gga Gly	ctg Leu	cag Gln	tat Tyr 225	Gly	ggt Gly	ctg Leu	Gly	cag Gln 230	:	1026
ctg Leu	gca Ala	gat Asp	ggt Gly	gtg Val 235	gtg Val	Gly	Leu	gat Asp	Asp	Phe	Arg	aag Lys	agt Ser	cag Gln 245	GLu	:	1074
ctg Leu	cgg Arg	gtc Val	tgg Trp 250	cca Pro	ggc Gly	tat Tyr	gac Asp	tat Tyr 255	Val	gga Gly	tgg Trp	agc Ser	aac Asn 260	His	agc Ser	:	1122
ttc Phe	tcc Ser	agt Ser 265	Gly	tat Tyr	gtg Val	gag Glu	atg Met 270	Glu	ttt Phe	gag Glu	ttt Phe	gac Asp 275	Arg	ctg Leu	agg Arg	:	1170
gcc Ala	ttc Phe 280	Gln	gct Ala	atg Met	cag Gln	gtc Val 285	cac His	tgt Cys	aac Asn	aac Asn	atg Met 290	cac His	acg Thr	ctg Leu	gga Gly		1218
gcc Ala 295	Arg	ctg Leu	cct Pro	ggc	300 Gly ggg	gtg Val	gaa Glu	tgt Cys	cgc Arg	ttc Phe 305	Arg	cgt Arg	ggc	cct Pro	gcc Ala 310		1266
atg Met	gcc Ala	tgg Trp	gag Glu	ggg Gly 315	Glu	CCC Pro	atg Met	cgc Arg	cac His 320	Asn	cta Leu	Gly	ggc	aac Asn 325	ctg Leu		1314

ejā aaa	gac Asp	ccc Pro	aga Arg 330	gcc Ala	cgg Arg	gct Ala	gtc Val	tca Ser 335	gtg Val	ccc Pro	ctt Leu	ggc Gly	ggc Gly 340	cgt Arg	gtg Val	1362
gct Ala	cgc Arg	ttt Phe 345	ctg Leu	cag Gln	tgc Cys	cgc Arg	ttc Phe 350	ctc Leu	ttt Phe	gcg Ala	Gly ggg	ccc Pro 355	tgg Trp	tta Leu	ctc Leu	1410
ttc Phe	agc Ser 360	gaa Glu	atc Ile	tcc Ser	ttc Phe	atc Ile 365	tct Ser	gat Asp	gtg Val	gtg Val	aac Asn 370	aat Asn	tcc Ser	tct Ser	ccg Pro	1458
gca Ala 375	ctg Leu	gga Gly	Gly	acc Thr	ttc Phe 380	ccg Pro	cca Pro	gcc Ala	ccc Pro	tgg Trp 385	tgg Trp	ccg Pro	cct Pro	ggc Gly	cca Pro 390	1506
cct Pro	ccc Pro	acc Thr	aac Asn	ttc Phe 395	agc Ser	agc Ser	ttg Leu	gag Glu	ctg Leu 400	gag Glu	ccc	aga Arg	ggc	cag Gln 405	cag Gln	1554
ccc	gtg Val	gcc Ala	aag Lys 410	gcc Ala	gag Glu	ggg Gly	agc Ser	ccg Pro 415	acc Thr	gcc Ala	atc Ile	ctc	atc Ile 420	ggc	tgc Cys	1602
ctg Leu	gtg Val	gcc Ala 425	atc Ile	atc Ile	ctg Leu	ctc Leu	ctg Leu 430	ctg Leu	ctc Leu	atc Ile	att Ile	gcc Ala 435	ctc Leu	atg Met	ctc Leu	1650
tgg Trp	cgg Arg 440	ctg Leu	cac His	tgg Trp	cgc	agg Arg 445	ctc Leu	ctc Leu	agc Ser	aag Lys	gct Ala 450	gaa Glu	cgg Arg	agg Arg	gtg Val	1698
ttg Leu 455	gaa Glu	gag Glu	gag Glu	ctg Leu	acg Thr 460	gtt Val	cac His	ctc Leu	tct Ser	gtc Val 465	cct	GJY	gac Asp	act Thr	atc Ile 470	1746
ctc Leu	atc Ile	aac Asn	aac Asn	cgc Arg 475	cca Pro	ggt Gly	CCĻ	aga Arg	gag :Glu 480	.Pro	ccc Pro	ccg Pro	tac Tyr	cag Gln 485	GIU	1794
ccc Pro	cgg	cct	cgt Arg 490	Gly	aat Asn	ccg Pro	ccc Pro	cac His 495	Ser	gct Ala	ccc Pro	tgt Cys	gtc Val 500	Pro	aat Asn	1842
ggc Gly	tct Ser	gcg Ala 505	Leu	ctg Leu	ctc Leu	tcc Ser	aat Asn 510	Pro	gcc Ala	tac Tyr	cgc Arg	cto Leu 515	Leu	ctg Leu	gcc Ala	1890
act	tac Tyr 520	Ala	cgt Arg	ccc Pro	cct Pro	cga Arg 525	Gly	e ccg	Gly ggc	Pro	Pro 530	) Tux	Pro	gcc Ala	tgg Trp	1938
gcc Ala 535	Lys	ccc Pro	acc	aac Asn	acc Thr 540	Gln	gcc Ala	tac Tyr	agt Ser	ggg Gly 545	/ Asp	tat Tyr	atg Met	gag Glu	r cct Pro 550	1986
gaç Ģlu	aag Lys	cca Pro	ggc Gly	gcc Ala 555	Pro	Lev	cto Lev	r ccc	cca Pro 560	Pro	Pro	caç Glr	g aac n Asr	ago Ser 565	gtc Val	2034
Pro	cat His	tat Tyr	gcc Ala 570	Glu	gct Ala	gac Asp	att Ile	gtt Val 575	LThi	cto Lev	g cag ı Glr	g ggo	gto Val 580	. Thi	: Gly	2082

ggc Gly	aac Asn	acc Thr 585	tat Tyr	gct Ala	gtg Val	cct Pro	gca Ala 590	ctg Leu	ccc Pro	cca Pro	gly ggg	gca Ala 595	gtç Val	GJ À āāā	gat Asp	2130
Gly	ccc Pro 600	ccc Pro	aga Arg	gtg Val	gat Asp	ttc Phe 605	cct Pro	cga Arg	tct Ser	cga Arg	ctc Leu 610	cgc Arg	ttc Phe	aag Lys	gag Glu	2178
aag Lys 615	ctt Leu	ggc Gly	gag Glu	ggc Gly	cag Gln 620	ttt Phe	GJ À Gàà	gag Glu	gtg Val	cac His 625	ctg Leu	tgt Cys	gag Glu	gtc Val	gac Asp 630	2226
agc Ser	cct Pro	caa Gln	gat Asp	ctg Leu 635	gtt Val	agt Ser	ctt Leu	gat Asp	ttc Phe 640	ccc Pro	ctt Leu	aat Asn	gtg Val	cgt Arg 645	aag Lys	2274
gga Gly	cac His	cct Pro	ttg Leu 650	ctg Leu	gta Val	gct Ala	gtc Val	aag Lys 655	atc Ile	tta Leu	cgg Arg	cca Pro	gat Asp 660	gcc Ala	acc Thr	2322
aag Lys	aat Asn	gcc Ala 665	agc Ser	ttc Phe	tcc Ser	ttg Leu	ttc Phe 670	tcc Ser	agg Arg	aat Asn	gat Asp	ttc Phe 675	ctg Leu	aaa Lys	gag Glu	2370
Val	aag Lys 680	atc Ile	atg Met	tcg Ser	agg Arg	ctc Leu 685	aag Lys	gac Asp	cca Pro	aac Asn	atc Ile 690	Ile	cgg Arg	ctg Leu	ctg Leu	2418
ggc Gly 695	gtg Val	tgt Cys	gtg Val	cag Gln	gac Asp 700	gac Asp	ccc Pro	ctc Leu	tgc Cys	atg Met 705	att Ile	act Thr	gac Asp	tac Tyr	atg Met 710	2466
gag Glu	aac Asn	ggc Gly	gac Asp	ctc Leu 715	aac Asn	cag Gln	ttc Phe	ctc Leu	agt Ser 720	gcc Ala	cac His	cag Gln	ctg Leu	gag Glu 725	gac Asp	2514
aag Lys	gca Ala	gcc Ala	gag Glu 730	G] À ààà	gcc Ala	cct Pro	Gly ggg	gac Asp 735	Gly	cag Gln	gct Ala	gcg Ala	cag Gln 740	GJ Å aaa	ccc Pro	2562
acc Thr	atc Ile	agc Ser 745	tac Tyr	cca Pro	atg Met	ctg Leu	ctg Leu 750	cat His	gtg Val	gca Ala	gcc Ala	cag Gln 755	atc Ile	gcc Ala	tcc Ser	2610
ggc Gly	atg Met 760	cgc Arg	tat Tyr	ctg Leu	gcc Ala	aca Thr 765	ctc Leu	aac Asn	ttt Phe	gta Val	cat His 770	cgg Arg	gac Asp	ctg Leu	gcc Ala	2658
acg Thr 775	cgg	aac Asn	tgc Cys	cta Leu	gtt Val 780	GJ À ààà	gaa Glu	aat Asn	ttc Phe	acc Thr 785	atc Ile	aaa Lys	atc Ile	gca Ala	gac Asp 790	2706
ttt Phe	ggc Gly	atg Met	agc Ser	cgg Arg 795	aac Asn	ctc Leu	tat Tyr	gct Ala	800 GJÀ âââ	gac Asp	tat Tyr	tac Tyr	cgt Arg	gtg Val 805	cag Gln	2754
Gly	cgg Arg	gca Ala	gtg Val 810	ctg Leu	ccc Pro	atc Ile	cgc Arg	tgg Trp 815	atg Met	gcc Ala	tgg Trp	gag Glu	tgc Cys 820	atc Ile	ctc Leu	2802
atg Met	ej aaa	aag Lys 825	ttc Phe	acg Thr	act Thr	gcg Ala	agt Ser 830	gac Asp	gtg ·Val	tgg Trp	gcc Ala	ttt Phe 835	ggt Gly	gtg Val	acc Thr	2850
ctg Leu	tgg Trp	gag Glu	gtg Val	ctg Leu	atg Met	ctc Leu	tgt Cys	agg Arg	gcc Ala	cag Gln	ccc Pro	ttt Phe	Gly ggg	tca Ser	gct Ala	2898

840			845	•				850	•				
cac cga c His Arg A 855	ga gca rg Ala	ggt cat Gly Hi:	s Arg	gaa Glu	cgc Arg	ggg Gly	gga Gly 865	gtt Val	ctt Leu	ccg Pro	gga Gly	cca Pro 870	2946
ggg ccg g Gly Pro A	gca gtg Mla Val	tac cto Tyr Let 875	g tcc u Ser	cgg.	ccg Pro	cct Pro 880	gcc Ala	tgc Cys	ccg Pro	cag Gln	ggc Gly 885	cta Leu	2994
tat gag o Tyr Glu I	tg atg Leu Met 890	Leu Ar	g tgc g Cys	tgg Trp	agc Ser 895	cgg Arg	gag Glu	tct Ser	gag Glu	cag Gln 900	cga Arg	cca . Pro	3042
ccc ttt t	cc cag Ser Gln 905	ctg ca Leu Hi	t cgg s Arg	ttc Phe 910	ctg Leu	gca Ala	gag Glu	gat Asp	gca Ala 915	ctc Leu	aac Asn	acg Thr	3090
gtg tgaat Val	tcacac	atccago	tgc c	cctc	ctc	a gg	gagc	gatc	cag	ggga	agc		3143
cagtgacad	ct aaaa	caagag	gacac	aatq	g ca	cctć	tgcc	ctt	cccc	tcc	cgac	agccca	3203
tcacctct													3263
cccttctc													3323
cccctgt													3383
tecettgg													3443
cctgggcc													3503
tccctgtc													3563
gaggaaaa													3623
tccatcac													3683
ttcccact													3743
tattggga													3803
gtgtagct													3863
agacacag													3923
taggcagg													3970
<210> 2	28												
<211> 9	919 .	•		•									

<212>. PRT

<213> NM\_013994 DDR1

<400> 28

Met Gly Pro Glu Ala Leu Ser Ser Leu Leu Leu Leu Leu Val Ala 5 10

Ser Gly Asp Ala Asp Met Lys Gly His Phe Asp Pro Ala Lys Cys Arg

20 25 30 Tyr Ala Leu Gly Met Gln Asp Arg Thr Ile Pro Asp Ser Asp Ile Ser 40 Ala Ser Ser Ser Trp Ser Asp Ser Thr Ala Ala Arg His Ser Arg Leu <sub>.</sub> 60 Glu Ser Ser Asp Gly Asp Gly Ala Trp Cys Pro Ala Gly Ser Val Phe Pro Lys Glu Glu Glu Tyr Leu Gln Val Asp Leu Gln Arg Leu His Leu 90 Val Ala Leu Val Gly Thr Gln Gly Arg His Ala Gly Gly Leu Gly Lys 105 110 Glu Phe Ser Arg Ser Tyr Arg Leu Arg Tyr Ser Arg Asp Gly Arg Arg 120 125 Trp Met Gly Trp Lys Asp Arg Trp Gly Gln Glu Val Ile Ser Gly Asn 135 Glu Asp Pro Glu Gly Val Val Leu Lys Asp Leu Gly Pro Pro Met Val 155 150 Ala Arg Leu Val Arg Phe Tyr Pro Arg Ala Asp Arg Val Met Ser Val 170 Cys Leu Arg Val Glu Leu Tyr Gly Cys Leu Trp Arg Asp Gly Leu Leu 180 185 190 Ser Tyr Thr Ala Pro Val Gly Gln Thr Met Tyr Leu Ser Glu Ala Val 200 Tyr Leu Asn Asp Ser Thr Tyr Asp Gly His Thr Val Gly Gly Leu Gln Tyr Gly Gly Leu Gly Gln Leu Ala Asp Gly Val Val Gly Leu Asp Asp Phe Arg Lys Ser Gln Glu Leu Arg Val Trp Pro Gly Tyr Asp Tyr Val 250 Gly Trp Ser Asn His Ser Phe Ser Ser Gly Tyr Val Glu Met Glu Phe

Glu Phe Asp Arg Leu Arg Ala Phe Gln Ala Met Gln Val His Cys Asn

Asn Met His Thr Leu Gly Ala Arg Leu Pro Gly Gly Val Glu Cys Arg

300 295 290 Phe Arg Arg Gly Pro Ala Met Ala Trp Glu Gly Glu Pro Met Arg His 315 310 Asn Leu Gly Gly Asn Leu Gly Asp Pro Arg Ala Arg Ala Val Ser Val 330 . 335 Pro Leu Gly Gly Arg Val Ala Arg Phe Leu Gln Cys Arg Phe Leu Phe Ala Gly Pro Trp Leu Leu Phe Ser Glu Ile Ser Phe Ile Ser Asp Val 365 360 Val Asn Asn Ser Ser Pro Ala Leu Gly Gly Thr Phe Pro Pro Ala Pro Trp Trp Pro Pro Gly Pro Pro Pro Thr Asn Phe Ser Ser Leu Glu Leu 395 390 . Glu Pro Arg Gly Gln Gln Pro Val Ala Lys Ala Glu Gly Ser Pro Thr 410 Ala Ile Leu Ile Gly Cys Leu Val Ala Ile Ile Leu Leu Leu Leu 425 . Ile Ile Ala Leu Met Leu Trp Arg Leu His Trp Arg Arg Leu Leu Ser 440 Lys Ala Glu Arg Arg Val Leu Glu Glu Glu Leu Thr Val His Leu Ser 450 Val Pro Gly Asp Thr Ile Leu Ile Asn Asn Arg Pro Gly Pro Arg Glu 475 Pro Pro Pro Tyr Gln Glu Pro Arg Pro Arg Gly Asn Pro Pro His Ser 490 Ala Pro Cys Val Pro Asn Gly Ser Ala Leu Leu Ser Asn Pro Ala 505 Tyr Arg Leu Leu Leu Ala Thr Tyr Ala Arg Pro Pro Arg Gly Pro Gly 515 Pro Pro Thr Pro Ala Trp Ala Lys Pro Thr Asn Thr Gln Ala Tyr Ser 535 540

Gly Asp Tyr Met Glu Pro Glu Lys Pro Gly Ala Pro Leu Leu Pro Pro

550

545

- Pro Pro Gln Asn Ser Val Pro His Tyr Ala Glu Ala Asp Ile Val Thr 565 570 575
- Leu Gln Gly Val Thr Gly Gly Asn Thr Tyr Ala Val Pro Ala Leu Pro 580 590
- Pro Gly Ala Val Gly Asp Gly Pro Pro Arg Val Asp Phe Pro Arg Ser 595 600 605
- Arg Leu Arg Phe Lys Glu Lys Leu Gly Glu Gly Gln Phe Gly Glu Val 610 620
- His Leu Cys Glu Val Asp Ser Pro Gln Asp Leu Val Ser Leu Asp Phe 625 630 635
- Pro Leu Asn Val Arg Lys Gly His Pro Leu Leu Val Ala Val Lys Ile 645 650 655
- Leu Arg Pro Asp Ala Thr Lys Asn Ala Ser Phe Ser Leu Phe Ser Arg 660 665 670
- Asn Asp Phe Leu Lys Glu Val Lys Ile Met Ser Arg Leu Lys Asp Pro 675 680 685
- Asn Ile Ile Arg Leu Leu Gly Val Cys Val Gln Asp Asp Pro Leu Cys 690 695 700
- Met Ile Thr Asp Tyr Met Glu Asn Gly Asp Leu Asn Gln Phe Leu Ser 705 710 715 720
- Ala His Gln Leu Glu Asp Lys Ala Ala Glu Gly Ala Pro Gly Asp Gly 725 730 735
- Gln Ala Ala Gln Gly Pro Thr Ile Ser Tyr Pro Met Leu His Val 740 745 . 750
- Ala Ala Gln Ile Ala Ser Gly Met Arg Tyr Leu Ala Thr Leu Asn Phe 755 760 765
- Val His Arg Asp Leu Ala Thr Arg Asn Cys Leu Val Gly Glu Asn Phe 770 780
- Thr Ile Lys Ile Ala Asp Phe Gly Met Ser Arg Asn Leu Tyr Ala Gly 785 790 795 800
- Asp Tyr Tyr Arg Val Gln Gly Arg Ala Val Leu Pro Ile Arg Trp Met 805 810 815
- Ala Trp Glu Cys Ile Leu Met Gly Lys Phe Thr Thr Ala Ser Asp Val 820 825

Trp Ala Phe Gly Val Thr Leu Trp Glu Val Leu Met Leu Cys Arg Ala 835 840 845	
Gln Pro Phe Gly Ser Ala His Arg Arg Ala Gly His Arg Glu Arg Gly 850 855 860	
Gly Val Leu Pro Gly Pro Gly Pro Ala Val Tyr Leu Ser Arg Pro Pro 865 870 875 880	
Ala Cys Pro Gln Gly Leu Tyr Glu Leu Met Leu Arg Cys Trp Ser Arg 885 890 895	
Glu Ser Glu Gln Arg Pro Pro Phe Ser Gln Leu His Arg Phe Leu Ala 900 905 910	
Glu Asp Ala Leu Asn Thr Val 915	
<210> 29	
<211> 5698	
<212> DNA	
<213> NM_001067 TOP2A	
<220>	
<221> CDS	
<222> (127)(4719)	
<223>	
<400> 29 aggttcaagt ggagctctcc taaccgacgc gcgtctgtgg agaagcggct tggtcggggg	60
tggtetegtg gggteetgee tgtttagteg ettteagggt tettgageee etteaegaee 13	20
gtcacc atg gaa gtg tca cca ttg cag cct gta aat gaa aat atg caa  Met Glu Val Ser Pro Leu Gln Pro Val Asn Glu Asn Met Gln  1 5 10	68
gtc aac aaa ata aag aaa aat gaa gat gct aag aaa aga ctg tct gtt Val Asn Lys Ile Lys Lys Asn Glu Asp Ala Lys Lys Arg Leu Ser Val 15 20 25 30	16
gaa aga atc tat caa aag aaa aca caa ttg gaa cat att ttg ctc cgc 2 Glu Arg Ile Tyr Gln Lys Lys Thr Gln Leu Glu His Ile Leu Leu Arg 35 40 45	64
cca gac acc tac att ggt tct gtg gaa tta gtg acc cag caa atg tgg Pro Asp Thr Tyr Ile Gly Ser Val Glu Leu Val Thr Gln Gln Met Trp 50 55 60	12
	~~

gtt tac gat gaa gat gtt ggc att aac tat agg gaa gtc act ttt gtt

Val	Tyr	Asp 65	Glu	Asp	Val	Glу	Ile 70	Asn	Tyr	Arg	Glu	Val 75	Thr	Phe	Val		
cct Pro	ggt Gly 80	ttg Leu	tac Tyr	aaa Lys	atc Ile	ttt Phe 85	gat Asp	gag Glu	att Ile	cta Leu	gtt Val 90	aat Asn	gct Ala	gcg Ala	gac Asp	40	
aac Asn 95	aaa Lys	caa Gln	agg Arg	gac Asp	cca Pro 100	aaa Lys	atg Met	tct Ser	tgt Cys	att Ile 105	aga Arg	gtc Val	aca Thr	att Ile	gat Asp 110	45	56
ccg Pro	gaa Glu	aac Asn	aat Asn	tta Leu 115	att Ile	agt Ser	ata Ile	tgg Trp	aat Asn 120	aat Asn	gga Gly	aaa Lys	ggt Gly	att Ile 125	cct Pro	50	04
gtt Val	gtt Val	gaa Glu	cac His 130	aaa Lys	gtt Val	gaa Glu	aag Lys	atg Met 135	tat Tyr	gtc Val	cca Pro	gct Ala	ctc Leu 140	ata Ile	ttt Phe	55	52
gga Gly	cag Gln	ctc Leu 145	cta Leu	act Thr	tct Ser	agt Ser	aac Asn 150	tat Tyr	gat Asp	gat Asp	gat Asp	gaa Glu 155	aag Lys	aaa Lys	gtg Val	60	00
aca Thr	ggt Gly 160	ggt Gly	cga Arg	aat Asn	ggc Gly	tat Tyr 165	gga Gly	gcc	aaa Lys	ttg Leu	tgt Cys 170	aac Asn	ata Ile	ttc Phe	agt Ser	64	48
acc Thr 175	aaa Lys	ttt Phe	act Thr	gtg Val	gaa Glu 180	aca Thr	gcc Ala	agt Ser	aga Arg	gaa Glu 185	tac Tyr	aag Lys	aaa Lys	atg Met	ttc Phe 190	6	96
aaa Lys	cag Gln	aca Thr	tgg Trp	atg Met 195	gat Asp	aat Asn	atg Met	gga Gly	aga Arg 200	Ala	ggt Gly	gag Glu	atg Met	gaa Glu 205	ctc Leu	7	44
aag Lys	ccc	ttc Phe	aat Asn 210	Gly	gaa Glu	gat Asp	tat Tyr	aca Thr 215	Cys	atc Ile	acc Thr	ttt Phe	cag Gln 220	cct Pro	gat Asp	7	92
ttg Leu	tct Ser	aag Lys 225	Phe	aaa Lys	atg Met	caa Gln	agc Ser 230	Leu	gac Asp	aaa Lys	gat Asp	att Ile 235	gtt Val	gca Ala	cta Leu	8	40
atg Met	gto Val 240	Arg	aga Arg	gca Ala	tat Tyr	gat Asp 245	Ile	gct Ala	gga Gly	tcc Ser	Thr 250	Lys	gat Asp	gtc Val	aaa Lys	8	88
gto Val 255	. Phe	ctt Lev	aat Asn	gga Gly	aat Asn 260	Lys	ctg Leu	cca Pro	gta Val	aaa Lys 265	G13	ttt Phe	cgt Arg	agt Ser	tat Tyr 270	9	36
gtg Val	gac Asp	atg Met	tat Tyr	ttg Lev 275	Lys	gac Asp	aag Lys	ttg Lev	gat Asp 280	Glu	act Thr	ggt Gly	aac Asn	Ser 285	ttg Leu	9	84
aaa Lys	ı gta ; Val	ata L Ile	cat His	Gli	caa Gln	gta Val	aac Asr	cac His 295	Aro	j tgg j Trp	gaa Glu	ı gtg ı Val	tgt Cys 300	тел	act Thr	10	32
ato Met	g agt	gaa Glu 305	ь Гр	ggo Gly	ttt Phe	cag Gln	caa Glr 310	ı Ile	ago Ser	ttt Phe	gto Val	aac L Asn 315	Ser	att Ile	gct Ala	10	080
aca Thi	tco Sei 320	: Буя	g ggt	ggo Gly	aga Arg	cat His 325	Va]	gat Asp	tat o Tyr	gta Val	gct Ala 330	ı Asp	cag Glr	g att	gtg Val	11	128

											•					
act Thr 335	aaa Lys	ctt Leu	gtt Val	gat Asp	gtt Val 340	gtg Val	aag Lys	aag Lys	aag Lys	aac Asn 345	aag Lys	ggt Gly	ggt Gly	gtt Val	gca Ala 350	1176
gta Val	aaa Lys	gca Ala	cat His	cag Gln 355	gtg Val	aaa Lys	aat Asn	His	atg Met 360	tgg Trp	att Ile	ttt Phe	gta Val	aat Asn 365	gcc Ala	1224
					acc Thr											1272
					ttt Phe											1320
atc Ile	aaa Lys 400	gct Ala	gcc Ala	att Ile	ggc Gly	tgt Cys 405	ggt Gly	att Ile	gta Val	gaa Glu	agc Ser 410	ata Ile	cta Leu	aac Asn	tgg Trp	1368
					caa Gln 420											1416
					aag Lys											1464
gca Ala	GJ À GGÀ	ggc Gly	cga Arg 450	aac Asn	tcc Ser	act Thr	gag Glu	tgt Cys 455	acg Thr	ctt Leu	atc Ile	ctg Leu	act Thr 460	gag Glu	gga Gly	1512
					ttg Leu											1560
					ttc Phe											1608
					cag Gln 500											1656
			Val		ctt Leu		Tyr		Lys	Asn						1704
tca Ser	ttg Leu	aag Lys	acg Thr 530	ctt Leu	cgt Arg	tat Tyr	Gly	aag Lys 535	ata Ile	atg Met	att Ile	atg Met	aca Thr 540	gat Asp	cag Gln	1752
					cac His											1800
					ctt Leu											1848
					gta Val 580											1896
agc	ctt	cct	gaa	ttt	gaa	gag	tgg	aag	agt	tct	act	cca	aat	cat	aaa	1944

Ser	Leu	Pro	Glu	Phe 595	Glu	Glu	Trp	Lys	Ser 600	Ser	Thr	Pro	Asn	ніs 605	Lys	
						tac Tyr										1992
						gca Ala										2040
						gat Asp 645										2088
						cga Arg										2136
						tta Leu										2184
						ctg Leu										2232
ctt Leu	atc Ile	ttg Leu 705	ttc Phe	tca Ser	aat Asn	tct Ser	gat Asp 710	aac Asn	gag Glu	aga Arg	tct Ser	atc Ile 715	cct Pro	tct Ser	atg Met	2280
gtg Val	gat Asp 720	ggt Gly	ttg Leu	aaa Lys	cca Pro	ggt Gly 725	cag Gln	aga Arg	aag Lys	gtt Val	ttg Leu 730	ttt Phe	act Thr	tgc Cys	ttc Phe	2328
aaa Lys 735	cgg Arg	aat Asn	gac Asp	aag Lys	cga Arg 740	gaa Glu	gta Val	aag Lys	gtt Val	gcc Ala 745	caa Gln	tta Leu	gct Ala	gga Gly	tca Ser 750	2376
						tat Tyr										2424
acc Thr	att Ile	atc Ile	aat Asn 770	ttg Leu	gct Ala	cag Gln	aat Asn	ttt Phe 775	gtg Val	ggt Gly	agc Ser	aat Asn	aat Asn 780	cta Leu	aac Asn	2472
ctc Leu	ttg Leu	cag Gln 785	ccc Pro	att Ile	ggt Gly	cag Gln	ttt Phe 790	ggt Gly	acc Thr	agg Arg	cta Leu	cat His 795	ggt Gly	ggc Gly	aag Lys	2520
gat Asp	tct Ser 800	gct Ala	agt Ser	cca Pro	cga Arg	tac Tyr 805	atc Ile	ttt Phe	aca Thr	atg Met	ctc Leu 810	agc Ser	tct Ser	ttg Leu	gct Ala	2568
cga Arg 815	ttg Leu	tta Leu	ttt Phe	cca Pro	cca Pro 820	aaa Lys	gat Asp	gat Asp	cac His	acg Thr 825	ttg Leu	aag Lys	ttt Phe	tta Leu	tat Tyr 830	2616
gat Asp	gac Asp	aac Asn	cag Gln	cgt Arg 835	gtt Val	gag Glu	cct Pro	gaa Glu	tgg Trp 840	tac Tyr	att Ile	cct Pro	att Ile	att Ile 845	ccc Pro	2664
atg Met	gtg Val	ctg Leu	ata Ile 850	aat Asn	ggt Gly	gct Ala	gaa Glu	gga Gly 855	atc Ile	ggt Gly	act Thr	ggg Gly	tgg Trp 860	tcc Ser	tgc Cys	2712

Lys	Ile	Pro . 865	aac t Asn 1	Phe	Asp	Val	Arg 870	Glu	Ile	Val	Asn	875	Ile	Arg	Arg .	2760
ttg Leu	atg Met 880	gat Asp	gga ( Gly (	gaa Glu	gaa Glu	cct Pro 885	ttg Leu	cca Pro	atg Met	ctt Leu	cca Pro 890	agt Ser	tac Tyr	aag Lys	aac Asn	2808
ttc Phe 895	aag Lys	ggt Gly	act a	att Ile	gaa Glu 900	gaa Glu	ctg Leu	gct Ala	cca Pro	aat Asn 905	caa Gln	tat Tyr	gtg Val	att Ile	agt Ser 910	2856
ggt Gly	gaa Glu	gta Val	gct ( Ala 1	att Ile 915	ctt Leu	aat Asn	tct Ser	aca Thr	acc Thr 920	Ile	gaa Glu	atc     Ile	tca Ser	gag Glu 925	ctt Leu	2904
ccc Pro	gtc Val	aga Arg	aca Thr 930	tgg Trp	acc Thr	cag Gln	aca Thr	tac Tyr 935	aaa Lys	gaa Glu	caa Gln	gtt Val	cta Leu 940	Glu	ccc Pro	2952
atg Met	ttg Leu	aat Asn 945	ggc	acc Thr	gag Glu	aag Lys	aca Thr 950	cct Pro	cct Pro	ctc Leu	ata Ile	aca Thr 955	Asp	tat Tyr	agg Arg	3000
gaa Glu	tac Tyr 960	cat His	aca Thr	gat Asp	acc Thr	act Thr 965	gtg Val	aaa Lys	ttt Phe	gtt Val	gtg Val 970	Lys	atç Met	act Thr	gaa Glu	3048
gaa Glu 975	aaa Lys	ctg Leu	gca Ala	gag Glu	gca Ala 980	gag Glu	aga Arg	gtt Val	gga Gly	cta Leu 985	His	aaa Lys	gto Val	ttc Phe	aaa Lys 990	3096
ctc Leu	caa Gln	act Thr	agt Ser	ctc Leu 995	aca Thr	tgc Cys	aac Asn	tct Ser	atg Met 100	Va	g ct l Le	t tt u Ph	t ga e As	p Hi	c gta s Val 05	3144
			aag Lys 1010	Lys	tai	t gad r Ası	e ac	g gt r Va 10	l. L	tg g eu A	at a sp I	tt c	eu F	iga Arg .020	gac Asp	3189
			ctc Leu 1025	Arg	a cti g Lei	t aaa 1 Ly:	a ta s Ty	t ta r Ty 10	r G	ga t ly I	ta a eu A	iga a irg I	ys (	gaa Slu 1035	tgg Trp	3234
		gga Gly	atg Met 1040	Let				a tc u Se 10	r A			tg a eu A	sn A		cag Gln	3279
gct Ala	cgc Arg	ttt Phe	atc Ile 1055	Let	a gaq ı Gli	g aa u Ly	a at s Il	a ga e As 10	рG	gc a	aa a ys l	ita a :le I	le l	tt [le [065	gaa Glu	3324
aat Asn	aag Lys	cct Pro	aag Lys 1070	Lys	a gaa s Gli	a tt u Le	a at u Il	t aa e Ly 10	s V	tt c	tg a eu 1	itt o	3ln 2	agg LO80	Gly gga	3369
tat Tyr	gat Asp	tcg Ser	gat Asp 1085	Pro	t gte	g aa l Ly	g gc s Al	a Tr	g a p I 90	aa g ys G	gaa ç Slu <i>R</i>	jec c	Sln (	caa Gln 1095	aag Lys	3414
gtt Val	cca	gat Asp	gaa Glu	gaa Gl	a ga u Gl	a aa u As	t ga n Gl	u Gl	u S	gt g Ser A	ac a	aac g Asn G	ilu l	.ys	gaa Glu	3459
	PLO	•	1100					11	05				•	L <b>11</b> 0		

Thr	Glu	Lys	Ser 1115	Asp	Ser	Val	Thr	Asp 1120	Ser	Gly	Pro	Thr	Phe 1125	Asn	
			gat Asp 1130											aaa Lys	3549
			tgc Cys 1145												3594
			aga Arg 1160						Leu					ttg Leu	3639
			att Ile 1175						Val					aaa Lys	3684
caa Gln	gat Asp	gaa Glu	caa Gln 1190	gtc Val	gga Gly	ctt Leu	cct Pro	999 Gly 1195	aaa Lys	GJ A GGG	Gly ggg	aag Lys	gcc Ala 1200	<b>aag</b>	3729
G1A aaa	aaa Lys	aaa Lys	aca Thr 1205	caa Gln	atg Met	gct Ala	gaa Glu	gtt Val 1210	ttg Leu	cct Pro	tct Ser	ccg Pro	cgt Arg 1215	ggt Gly	3774
	-	-	att Ile 1220					ata Ile 1225						gca Ala	3819
-	-		aat Asn 1235		_			_					gaa Glu 1245	gga Gly	3864
			gaa Glu 1250												3909
	-	-	aaa Lys 1265	_		_	_				_			aaa Lys	3954
			ttg Leu 1280											aga Arg	3999
aat Asn	ccc Pro	tgg Trp	tct Ser 1295	gat Asp	tca Ser	gaa Glu	tca Ser	gat Asp 1300	agg Arg	agc Ser	agt Ser	gac Asp	gaa Glu 1305	agt Ser	4044
			gtc Val 1310												4089
			aaa Lys 1325											ttc Phe	4134
tca Ser	gat Asp	ttt Phe	gat Asp 1340	gaa Glu	aaa Lys	act Thr	gat Asp	gat Asp 1345	gaa Glu	gat Asp	ttt Phe	gtc Val	cca Pro 1350	tca Ser	4179
gat Asp	gct Ala	agt Ser	cca Pro 1355	cct Pro	aag Lys	acc Thr	aaa Lys	act Thr 1360	tcc Ser	cca Pro	aaa Lys	ctt Leu	agt Ser 1365	aac Asn	4224

aaa Lys	gaa Glu	ctg Leu	aaa Lys 1370	cca Pro	cag Gln	aaa Lys	agt Ser	gtc Val 1375	Val	tca Ser	gac Asp	ctt Leu	gaa Glu 1380	gct Ala	4269
gat Asp	gat Asp	gtt Val	aag Lys 1385	ggc Gly	agt Ser	gta Val	cca Pro	ctg Leu 1390	tct Ser	tca Ser	agc Ser	cct Pro	cct Pro 1395	gct Ala	4314
aca Thr	cat His	ttc Phe	cca Pro 1400	Asp	gaa Glu	act Thr	gaa Glu	att Ile 1405	aca Thr	aac Asn	cca Pro	gtt Val	cct Pro 1410	aaa Lys	4359
aag Lys	aat Asn	gtg Val	aca Thr 1415	gtg Val	aag Lys	aag Lys	aca Thr	gca Ala 1420	gca Ala	aaa Lys	agt Ser	cag Gln	tct Ser 1425	tcc Ser	4404
acc Thr	tcc Ser	act Thr	acc Thr 1430	ggt Gly	gcc Ala	aaa Lys	aaa Lys	agg Arg 1435	gct Ala	gcc Ala	cca Pro	aaa Lys	gga Gly 1440	act Thr	4449
aaa Lys	agg Arg	gat Asp	cca Pro 1445	gct Ala	ttg Leu	aat Asn	tct Ser	ggt Gly 1450	Val	tct Ser	caa Gln	aag Lys	cct Pro 1455	gat Asp	449,4
			acc Thr 1460	Lys	aat Asn	cgc Arg	cgc Arg	aaa Lys 1465	Arg	aag Lys	cca Pro	tcc Ser	act Thr 1470	tct Ser	4539
			gac Asp 1475	Ser	aat Asn	ttt Phe	gag Glu	aaa Lys 1480	Ile	gtt Val	tcg Ser	aaa Lys	gca Ala 1485	gtc Val	4584
			aaa Lys 1490	Ser	aag Lys	G] À	gag Glu	agt Ser 1495	Asp	gac Asp	ttc Phe	cat His	atg Met 1500	Asp	4629
ttt Phe	gac Asp	tca Ser	gct Ala 1505	Val	gct Ala	cct Pro	cgg Arg	gca Ala 1510	Lys	tct Ser	gta Val	. cgg . Arg	gca Ala 1515	газ	4674
aaa Lys	cct Pro	ata Ile	aag Lys 1520	Tyr	ctg Leu	gaa Glu	gag Glu	tca Ser 1525	Asp	gaa Glu	gat Asp	gat Asp	ctg Leu 1530	Pne	4719
taa	aatg	tga	ggcga	ttat	t tt	aagt	aatt	atct	tacc	aa g	ccca	agac	t ggt	tttaaag	4779
tta	cctg	aag	ctctt	aact	t co	tccc	ctct	gaat	ttag	ıtt t	gggg	gaagg	ıt gtt	tttagta	4839
caa	igaca	tca	aagtg	aagt	a aa	geco	aagt	gtto	ttta	igc t	tttt	ataa	ıt act	gtctaaa	4899
tag	gtgac	cat	ctcat	gggc	a tt	gtt:	tct	ctct	gctt	tg t	ctgt	gttt	t gag	gtctgctt	4959
tct	tttç	ıtct	ttaaa	acct	g at	tttt	aagt	tctt	ctga	ac t	gtag	gaaat	a gct	atctgat	5019
cac	ttca	ıgcg	taaag	cagt	g to	ıttta	ttaa	ccat	ccac	ta a	igcta	aaaa	et aga	gcagttt	5079
gat	ttaa	aag	tgtca	ctct	t co	etcct	tttc	tact	ttca	agt a	agata	atgag	ga tag	gagcataa	5139
tta	atcto	gttt	tatct	tagt	t ti	atad	ataa	ttta	accat	ca ç	gata	gaact	t tat	ggttcta	5199
gta	acaga	atac	tctac	taca	c to	cagco	ctctt	: atgt	gcca	ag 1	tttt	ctti	a ago	caatgaga	5259
aat	tgct	cat	gttct	tcat	c ti	ctca	aato	atca	agago	gec a	aaaga	aaaa	ac act	tttggctg	5319
tgt	ctat	caac	ttgad	cacaç	gt ca	ata	gaato	g aaga	aaat	ta 🤈	gagta	agtta	at gt	gattattt	5379

cagetettga cetgteecet o	etggetgeet etgagtetg	a atctcccaaa gagagaaacc	5439
aatttctaag aggactggat t	gcagaagac teggggaca	a catttgatcc aagatcttaa	5499
atgttatatt gataaccatg o	ctcagcaatg agctattag	a ttcattttgg gaaatctcca	5559
taatttcaat ttgtaaactt t	gttaagacc tgtctacat	t gttatatgtg tgtgacttga	5619
gtaatgttat caacgttttt g	gtaaatattt actatgttt	t tctattagct aaattccaac	5679
aattttgtac tttaataaa		•	5698
· <01'0> 20			
<210> 30			
<211> 1531		· .	
<212> PRT			
<213> NM_001067 TOP2A			
<400> 20			
<400> 30	a) - p - m ) - a)		
Met Glu Val Ser Pro Leu 1 5	i GIn Pro Val Asn GI 10	ų Asn Met Gin Vai Asn 15	
	,		
Lys Ile Lys Lys Asn Glu 20	Asp Ala Lys Lys Ar 25	g Leu Ser Val Glu Arg 30	
<b>71</b>	a		
Ile Tyr Gln Lys Lys Thr 35	40	e Leu Leu Arg Pro Asp 45	
The Tie Clarker Wal	Clu Iou Val Mbm Cl	- Cla Mat Two Val Tyr	
Thr Tyr Ile Gly Ser Val 50	55	60	
han Clu han Val Clu Tla	. A	I Mbw Dho Wal Dwo Clu	
Asp Glu Asp Val Gly Ile	r Ash Tyr Arg Giu va 75	1 Thi Phe Val Plo Gly 80	
Y M Y T1 N			
Leu Tyr Lys Ile Phe Asp . 85	90 gru lie Leu val As	n Ala Ala Asp Asn Lys 95	
Ola North Day Day Torr Walt	d O 71 . 3	l mba tla bas Due Gla	
Gln Arg Asp Pro Lys Met 100	Ser Cys 11e Arg Va 105	1 Thr lie Asp Pro Giu	
Non-Non-You The Gov The		- Clar II- Due Wel Wel	
Asn Asn Leu Ile Ser Ile 115	120	s GIY IIE Pro Vai Vai 125	
Ole rile tue vel ole v	Mak Man W-1 Por- 22	- Iou Tle Dh- Clu Cl-	
Glu His Lys Val Glu Lys	-	_	
130	135	140	
Leu Leu Thr Ser Ser Asn			
145 150	. 15.	5 160	

Gly Arg Asn Gly Tyr Gly Ala Lys Leu Cys Asn Ile Phe Ser Thr Lys 165 170 175

Phe Thr Val Glu Thr Ala Ser Arg Glu Tyr Lys Lys Met Phe Lys Gln 180 185 . 190

Thr Trp Met Asp Asn Met Gly Arg Ala Gly Glu Met Glu Leu Lys Pro

Phe Asn Gly Glu Asp Tyr Thr Cys Ile Thr Phe Gln Pro Asp Leu Ser 210 215 220

Lys Phe Lys Met Gln Ser Leu Asp Lys Asp Ile Val Ala Leu Met Val 225 230 240

Arg Arg Ala Tyr Asp Ile Ala Gly Ser Thr Lys Asp Val Lys Val Phe 245 250 255

Leu Asn Gly Asn Lys Leu Pro Val Lys Gly Phe Arg Ser Tyr Val Asp 260 265 270

Met Tyr Leu Lys Asp Lys Leu Asp Glu Thr Gly Asn Ser Leu Lys Val 275 280 285

Ile His Glu Gln Val Asn His Arg Trp Glu Val Cys Leu Thr Met Ser 290 295 300

Glu Lys Gly Phe Gln Gln Ile Ser Phe Val Asn Ser Ile Ala Thr Ser 305 310 315 320

Lys Gly Gly Arg His Val Asp Tyr Val Ala Asp Gln Ile Val Thr Lys 325 330 335

Leu Val Asp Val Val Lys Lys Lys Asn Lys Gly Gly Val Ala Val Lys 340 345 350

Ala His Gln Val Lys Asn His Met Trp Ile Phe Val Asn Ala Leu Ile 355 360 365

Glu Asn Pro Thr Phe Asp Ser Gln Thr Lys Glu Asn Met Thr Leu Gln 370 375 380

Pro Lys Ser Phe Gly Ser Thr Cys Gln Leu Ser Glu Lys Phe Ile Lys 385 390 395

Ala Ala Ile Gly Cys Gly Ile Val Glu Ser Ile Leu Asn Trp Val Lys 405 410 415

Phe Lys Ala Gln Val Gln Leu Asn Lys Lys Cys Ser Ala Val Lys His
420 425 430

Asn Arg Ile Lys Gly Ile Pro Lys Leu Asp Asp Ala Asn Asp Ala Gly
435 440 445

Gly Arg Asn Ser Thr Glu Cys Thr Leu Ile Leu Thr Glu Gly Asp Ser 450 455 460

Ala Lys Thr Leu Ala Val Ser Gly Leu Gly Val Val Gly Arg Asp Lys
465 470 475 480

Tyr Gly Val Phe Pro Leu Arg Gly Lys Ile Leu Asn Val Arg Glu Ala 485 490 495

Ser His Lys Gln Ile Met Glu Asn Ala Glu Ile Asn Asn Ile Ile Lys 500 505 510

Ile Val Gly Leu Gln Tyr Lys Lys Asn Tyr Glu Asp Glu Asp Ser Leu 515 520 525

Lys Thr Leu Arg Tyr Gly Lys Ile Met Ile Met Thr Asp Gln Asp Gln 530 540

Asp Gly Ser His Ile Lys Gly Leu Leu Ile Asn Phe Ile His His Asn 545 550 555 560

Trp Pro Ser Leu Leu Arg His Arg Phe Leu Glu Glu Phe Ile Thr Pro 565 570 575

Ile Val Lys Val Ser Lys Asn Lys Gln Glu Met Ala Phe Tyr Ser Leu 580 585 590

Pro Glu Phe Glu Glu Trp Lys Ser Ser Thr Pro Asn His Lys Lys Trp 595 600 605

Lys Val Lys Tyr Tyr Lys Gly Leu Gly Thr Ser Thr Ser Lys Glu Ala 610 615 620

Lys Glu Tyr Phe Ala Asp Met Lys Arg His Arg Ile Gln Phe Lys Tyr 625 630 635 640

Ser Gly Pro Glu Asp Asp Ala Ala Ile Ser Leu Ala Phe Ser Lys Lys
645 650 655

Gln Ile Asp Asp Arg Lys Glu Trp Leu Thr Asn Phe Met Glu Asp Arg 660 665 670

Arg Gln Arg Lys Leu Leu Gly Leu Pro Glu Asp Tyr Leu Tyr Gly Gln 675 680 685

Thr Thr Tyr Leu Thr Tyr Asn Asp Phe Ile Asn Lys Glu Leu Ile 690 695 700

Leu Phe Ser Asn Ser Asp Asn Glu Arg Ser Ile Pro Ser Met Val Asp

705 710 715 720

Gly Leu Lys Pro Gly Gln Arg Lys Val Leu Phe Thr Cys Phe Lys Arg 725 730 735

Asn Asp Lys Arg Glu Val Lys Val Ala Gln Leu Ala Gly Ser Val Ala 740 745 750

Glu Met Ser Ser Tyr His His Gly Glu Met Ser Leu Met Met Thr Ile 755 760 . 765

Ile Asn Leu Ala Gln Asn Phe Val Gly Ser Asn Asn Leu Asn Leu France 770 780

Gln Pro Ile Gly Gln Phe Gly Thr Arg Leu His Gly Gly Lys Asp Ser 785 790 795 800

Ala Ser Pro Arg Tyr Ile Phe Thr Met Leu Ser Ser Leu Ala Arg Leu 805 810 815

Leu Phe Pro Pro Lys Asp Asp His Thr Leu Lys Phe Leu Tyr Asp Asp 820 825 830

Asn Gln Arg Val Glu Pro Glu Trp Tyr Ile Pro Ile Ile Pro Met Val 835 840 845

Leu Ile Asn Gly Ala Glu Gly Ile Gly Thr Gly Trp Ser Cys Lys Ile 850 855 860

Pro Asn Phe Asp Val Arg Glu Ile Val Asn Asn Ile Arg Arg Leu Met 865 870 875 880

Asp Gly Glu Glu Pro Leu Pro Met Leu Pro Ser Tyr Lys Asn Phe Lys 885 890 . 895

Gly Thr Ile Glu Glu Leu Ala Pro Asn Gln Tyr Val Ile Ser Gly Glu 900 905 910

Val Ala Ile Leu Asn Ser Thr Thr Ile Glu Ile Ser Glu Leu Pro Val 915 920 925

Arg Thr Trp Thr Gln Thr Tyr Lys Glu Gln Val Leu Glu Pro Met Leu 930 935 940

Asn Gly Thr Glu Lys Thr Pro Pro Leu Ile Thr Asp Tyr Arg Glu Tyr 945 950 955 960

His Thr Asp Thr Thr Val Lys Phe Val Val Lys Met Thr Glu Glu Lys 965 970 975

Leu Ala Glu Ala Glu Arg Val Gly Leu His Lys Val Phe Lys Leu Gln

980 985 990

Thr Ser Leu Thr Cys Asn Ser Met  $\mbox{Val}$  Leu Phe Asp His  $\mbox{Val}$  Gly Cys  $\mbox{995}$   $\mbox{1000}$ 

Leu Lys Lys Tyr Asp Thr Val Leu Asp Ile Leu Arg Asp Phe Phe 1010 1015 1020

Glu Leu Arg Leu Lys Tyr Tyr Gly Leu Arg Lys Glu Trp Leu Leu 1025 1030 1035

Gly Met Leu Gly Ala Glu Ser Ala Lys Leu Asn Asn Gln Ala Arg 1040 1045 1050

Phe Ile Leu Glu Lys Ile Asp Gly Lys Ile Ile Glu Asn Lys 1055 1060 1065

Pro Lys Lys Glu Leu Ile Lys Val Leu Ile Gln Arg Gly Tyr Asp 1070 1075 1080

Ser Asp Pro Val Lys Ala Trp Lys Glu Ala Gln Gln Lys Val Pro 1085 1090 1095

Asp Glu Glu Glu Asn Glu Glu Ser Asp Asn Glu Lys Glu Thr Glu 1100 1105 1110

Lys Ser Asp Ser Val Thr Asp Ser Gly Pro Thr Phe Asn Tyr Leu 1115 1120 1125

Leu Asp Met Pro Leu Trp Tyr Leu Thr Lys Glu Lys Lys Asp Glu 1130 1140

Leu Cys Arg Leu Arg Asn Glu Lys Glu Gln Glu Leu Asp Thr Leu 1145 1150 1155

Lys Arg Lys Ser Pro Ser Asp Leu Trp Lys Glu Asp Leu Ala Thr 1160 1165 1170

Phe Ile Glu Glu Leu Glu Ala Val Glu Ala Lys Glu Lys Gln Asp 1175 1180 1185

Glu Gln Val Gly Leu Pro Gly Lys Gly Gly Lys Ala Lys Gly Lys 1190 1195 1200

Lys Thr Gln Met Ala Glu Val Leu Pro Ser Pro Arg Gly Gln Arg 1205 1210 1215

Val Ile Pro Arg Ile Thr Ile Glu Met Lys Ala Glu Ala Glu Lys 1220 1225 1230

Lys Asn Lys Lys Lys Ile Lys Asn Glu Asn Thr Glu Gly Ser Pro

. 12	235					1240		٠			1245			
Gln Gl 12	Lu A 250	sp (	Gly	Val	G1u	Leu 1255	Glu	Gly	Leu	Lys	Gln 1260	Arg	Leu	Glu
Lys Ly	/s G 265	ln I	Гуs	Arg	Glu	Pro 1270	Gly	Thr.	Lys	Thr	Lys 1275	Lys ·	Gln <sub>.</sub>	Thr
Thr Le	eu A 280	la 1	Phe <sub>.</sub>	Lys	Pro	Ile 1285	Lys	·Lys	Gly	Lys	Lys 1290	Arg	Asn	Pro
Trp Se	er A 295	sp s	Ser	Glu	Ser	Asp 1300	Arg	Ser	Ser	Asp	Glu 1305	Ser	Asn	Phe
Asp Va	al P 310	ro l	Pro	Arg	Glu	Thr 1315	G1u	Pro	Arg	Arg	Ala 1320	Ala	Thr	Lys
Thr Ly	ys P 325	he :	Thr	Met	Asp	Leu 1330	Asp	Ser	Asp	Glu	Asp 1335	Phe	Ser	Asp
Phe As	sp G 340	lu 1	Lys	Thr	Asp	Asp 1345	Glu	Asp	Phe	Val	Pro 1350	Ser	Asp	Ala
Ser Pr	ro P 355	ro l	Lys	Thr	Lys	Thr 1360	Ser	Pro	Lys	Leu	Ser 1365	Asn	Lys	Glu
Leu Ly	ys P 370	ro (	Gln	Lys	Ser	Val 1375	Val	Ser	Asp	Leu	Glu 1380	Ala	Asp	Asp
Val Ly	ys G 385	ly:	Ser	Val		Leu 1390	Ser	Ser	Ser	Pro	Pro 1395	Ala	Thr	His
Phe Pr	ro A 400	sp (	Glu	Thr	Glu	Ile 1405	Thr	Asn	Pro	Val	Pro 1410	Lys	Lys	Asn
Val Th	nr V 415	al 1	Lys	Lys	Thr	Ala 1420	Ala	Lys	Ser	Gln	Ser 1425	Ser	Thr	Ser
Thr Th	nr G 430	ly i	Ala	Lys		Arg 1435	Ala	Ala	Pro	Lys	Gly 1440	Thr	Lys	Arg
Asp Pr	co A 145	la 1	Leu	Asn	Ser	Gly 1450	Val	Ser	Gln	Lys	Pro 1455	Asp	Pro	Ala
Lys Th	nr I 160	ys 1	Asn	Arg	Arg	Lys 1465	Arg	Lys	Pro	Ser	Thr 1470	Ser	Asp	Asp
Ser As	sp S 175	er 1	Asn	Phe	Glu	Lys 1480	Ile	Val	Ser	Lys	Ala 1485	Val	Thr	Ser

Lys Lys Ser Lys Gly Glu Ser Asp Asp Phe His Met Asp Phe Asp 1490 1495 1500	
Ser Ala Val Ala Pro Arg Ala Lys Ser Val Arg Ala Lys Lys Pro	
1505 1510 1515	
Ile Lys Tyr Leu Glu Glu Ser Asp Glu Asp Asp Leu Phe 1520 1525 1530	
<210> 31	
<211> 1369	
<212> DNA	
<213> NM_138410 chemokine-like factor super family 7, CKLFSF7	
<220>	
<221> CDS	
<222> (237)(761)	
<223>	•
<400> 31 agtotaactt cogogoatet acgagggoog ggaotgoogo tacetttotg gaaggogoog	60
gccggccagt caccggaggg atcccgccag aggcggctcc gcgtctccag cagcggggca	120
gggacccggg cgccccgccc tegecagcgc ccgcccccgc ccgcccccgg cccgccctct	180
gtatetggce eetgggeage tgeeegggga ggeggeeage gagetgggge egegea atg Met 1	239
tcg cac gga gcc ggg ctc gtc cgc acc acg tgc agc agc ggc agc gcg Ser His Gly Ala Gly Leu Val Arg Thr Thr Cys Ser Ser Gly Ser Ala 5 10	287
ctc gga ccc ggg gcc ggc gcg gcc cag ccc agc gcg agc ccc ttg gag	
Leu Gly Pro Gly Ala Gly Ala Ala Gln Pro Ser Ala Ser Pro Leu Glu 20 25 30	335
Leu Gly Pro Gly Ala Gly Ala Ala Gln Pro Ser Ala Ser Pro Leu Glu	335 383
Leu Gly Pro Gly Ala Gly Ala Ala Gln Pro Ser Ala Ser Pro Leu Glu 20 25 30  ggg ctg ctg gac ctc agc tac ccc cgc acc cac gcg gcc ctg ctg aaa Gly Leu Leu Asp Leu Ser Tyr Pro Arg Thr His Ala Ala Leu Leu Lys	
Leu Gly Pro Gly Ala Gly Ala Ala Gln Pro Ser Ala Ser Pro Leu Glu 20 25 30  ggg ctg ctg gac ctc agc tac ccc cgc acc cac gcg gcc ctg ctg aaa Gly Leu Leu Asp Leu Ser Tyr Pro Arg Thr His Ala Ala Leu Leu Lys 35 40 45  gtg gcg caa atg gtc acc ctg ctg att gcc ttc atc tgt gtg cgg agc Val Ala Gln Met Val Thr Leu Leu Ile Ala Phe Ile Cys Val Arg Ser	383
Leu Gly Pro Gly Ala Gly Ala Ala Gln Pro Ser Ala Ser Pro Leu Glu 20 25 30  ggg ctg ctg gac ctc agc tac ccc cgc acc cac gcg gcc ctg ctg aaa Gly Leu Leu Asp Leu Ser Tyr Pro Arg Thr His Ala Ala Leu Leu Lys 35 40 45  gtg gcg caa atg gtc acc ctg ctg att gcc ttc atc tgt gtg cgg agc Val Ala Gln Met Val Thr Leu Leu Ile Ala Phe Ile Cys Val Arg Ser 50 60 65  tcc ctg tgg acc aac tac agc gcc tac agc tac ttt gaa gtg gtc acc Ser Leu Trp Thr Asn Tyr Ser Ala Tyr Ser Tyr Phe Glu Val Val Thr	383 431

Arg Phe Tyr Arg Val Leu Thr	Cys Ile Ser Trp 105	Pro Leu Ser Glu Leu 110	•
ctg cac tat tta atc ggt acc Leu His Tyr Leu Ile Gly Thr 115 120	Leu Leu Leu Leu		623
gca gct tcc aag agt tac aac Ala Ala Ser Lys Ser Tyr Asn 130	cag agc gga ctg Gln Ser Gly Leu 140	gta gcc gga gcg atc Val Ala Gly Ala Ile 145	671
ttt ggt ttc atg gcc acc ttc Phe Gly Phe Met Ala Thr Phe 150			719
tat aag atc tcg tgt gta acc Tyr Lys Ile Ser Cys Val Thr 165			761
tgatgaggcc acaaccccta ggccc	ctcag gagetttgca	gagaggagga cgtgtactcc	821
aggcgaggcc tctggacctg tgttc	ctgtg ccaaagtcct	gtcaggctgg tgggcaccag	881
gaaaggcctg caccctcttc ctgct	ctccc aggaagccag	ctccctgagc tcctgagcca	941
gccggaaact cttcctccag ccttc	cgggg agaacatccc	teccattetg ggaaaggaaa	1001
gcagcctcca gggaaatgtt ttctg	ccttc ctgcttctag	aaccacctca ggtactgatg	1061
aaccccactt agcacagctg aaggg	gtttg tgaatactcc	cgcctaaatc ccttctactt	1121
cactcctcag gggagtgaag tgcct	taaga aacaaagccc	tgtcctaatt tatctagctt	1181
gtcagtccgg tcttagagat accct	ctttc ctgaagtgag	gcgtgcctgt agaaacacta	1241
tgtggtcagc ctgtccccaa ggaga	tettg tgteteetet	ccatctctgc ctttgttacc	1301
agtgtgcatg tgtttgtgtg ttttt	taata aaatattgac	tcggccagtt aaaaaaaaaa	1361
aaaaaaaa			1369
<210> 32	٠.	, .	
<211> 175			
<212> PRT		•	
<213> NM_138410 chemokine-	like factor super	r family 7, CKLFSF7	
<400> 32			
Met Ser His Gly Ala Gly Leu 1 5	Val Arg Thr Thr	Cys Ser Ser Gly Ser 15	•
Ala Leu Gly Pro Gly Ala Gly 20	Ala Ala Gln Pro 25	Ser Ala Ser Pro Leu 30	
Glu Gly Leu Leu Asp Leu Ser 35	Tyr Pro Arg Thr	His Ala Ala Leu Leu 45	

Lys Val Ala Gln Met Val Thr Leu Leu Ile Ala Phe Ile Cys Val Arg

60

55

50

PCT/AU2003/001166

Ser Ser Leu Trp Thr Asn Tyr Ser Ala Tyr Ser Tyr Phe Glu Val 80

Thr Ile Cys Asp Leu 85

He Met Ile Leu Ala Phe Tyr Leu Val His Leu 90

Phe Arg Phe Tyr Arg Val Leu Thr Cys Ile Ser Trp Pro Leu Ser Glu 100 105 110

Leu Leu His Tyr Leu Ile Gly Thr Leu Leu Leu Leu Ile Ala Ser Ile
115 120 125

Val Ala Ala Ser Lys Ser Tyr Asn Gln Ser Gly Leu Val Ala Gly Ala 130 135 140

Ile Phe Gly Phe Met Ala Thr Phe Leu Cys Met Ala Ser Ile Trp Leu 145 150 155 160

Ser Tyr Lys Ile Ser Cys Val Thr Gln Ser Thr Asp Ala Ala Val 165 170 170

<210> 33

<211> 3470

<212> DNA

<213> NM\_012112 C20orfl, chromosome 20 open reading frame 1

<220>

<221> CDS

<222> (458)..(2698)

<223>

<400> 33 ggctaggtga gccgtgggaa gaaaagaggg agcagctagg gcgcgggtct ccctcctccc 60 120 ggagtttgga acggctgaag ttcaccttcc agcccctagc gccgttcgcg ccgctaggcc 180 tggcttctga ggcggttgcg gtgctcggtc gccgcctagg cggggcaggg tgcgagcagg 240 ggcttcgggc cacgcttctc ttggcgacag gattttgctg tgaagtccgt ccgggaaacg 300 gaggaaaaaa agagttgcgg gaggctgtcg gctaataacg gttcttgata catatttgcc agacttcaag atttcagaaa aggggtgaaa gagaagattg caactttgag tcagacctgt 360 aggeetgata gaetgattaa accacagaag gtgacetget gagaaaagtg gtacaaatac 420 tgggaaaaac ctgctcttct gcgttaagtg ggagaca atg tca caa gtt aaa agc 475 Met Ser Gln Val Lys Ser 1

tct Ser	tat Tyr	tcc Ser	tat Tyr 10	gat Asp	gcc Ala	ccc Pro	tcg Ser	gat Asp 15	ttc Phe	atc Ile	aat Asn	ttt Phe	tca Ser 20	tcc Ser	ttg Leu	523
gat Asp	gat Asp	gaa Glu 25	gga Gly	gat Asp	act Thr	caa Gln	aac Asn 30	ata Ile	gat Asp	tca Ser	tgg Trp	ttt Phe 35	gag Glu	gag Glu	aag Lys	571
gcc Ala	aat Asn 40	ttg Leu	gag Glu	aat Asn	Lys	tta Leu 45	ctg Leu	GJ À GGG	aag Lys	aat Asn	gga Gly 50	act Thr	gga Gly	GJA āāā	ctt Leu	619
ttt Phe 55	cag Gln	ggc Gly	aaa Lys	act Thr	cct Pro 60	ttg Leu	aga Arg	aag Lys	gct Ala	aat Asn 65	ctt Leu	cag Gln	caa Gln	gct Ala	att Ile 70	667
gtc Val	aca Thr	cct Pro	ttg Leu	aaa Lys 75	cca Pro	gtt Val	gac Asp	aac Asn	act Thr 80	tac Tyr	tac Tyr	aaa Lys · .	gag Glu	gca Ala 85	gaa Glu	715
aaa Lys	gaa Glu	aat Asn	ctt Leu 90	gtg Val	gaa Glu	caa Gln	tcc Ser	att Ile 95	ccg Pro	tca Ser	aat Asn	gct Ala	tgt Cys 100	tct Ser	tcc Ser	763
ctg Leu	gaa Glu	gtt Val 105	gag Glu	gca Ala	gcc Ala	ata Ile	tca Ser 110	aga Arg	aaa Lys	act Thr	cca Pro	gcc Ala 115	cag Gln	cct Pro	cag Gln	811
aga Arg	aga Arg 120	tct Ser	ctt Leu	agg Arg	ctt Leu	tct Ser 125	gct Ala	cag Gln	aag Lys	gat Asp	ttg Leu 130	gaa Glu	cag Gln	aaa Lys	gaa Glu	859
aag Lys 135	cat His	cat His	gta Val	aaa Lys	atg Met 140	aaa Lys	gcc Ala	aag Lys	aga Arg	tgt Cys 145	gcc Ala	act Thr	cct Pro	gta Val	atc Ile 150	907
atc Ile	gat Asp	gaa Glu	att Ile	cta Leu 155	ccc Pro	tct Ser	aag Lys	aaa Lys	atg Met 160	aaa Lys	gtt Val	tct Ser	aac Asn	aac Asn 165	aaa Lys	955
aag Lys	aag Lys	cca Pro	gag Glu 170	gaa Glu	gaa Glu	ggc Gly	agt Ser	gct Ala 175	cat His	caa Gln	gat Asp	act Thr	gct Ala 180	gaa Glu	aag L <b>ỷ</b> s	1003
aat Asn	gca Ala	tct Ser 185	tcc Ser	cca Pro	gag Glu	aaa Lys	gcc Ala 190	aag Lys	ggt Gly	aga Arg	cat	act Thr 195	gtg Val	cct Pro	tgt Cys	1051
atg Met	cca Pro 200	cct Pro	gca Ala	aag Lys	cag Gln	aag Lys 205	ttt Phe	cta Leu	aaa Lys	agt Ser	act Thr 210	gag Glu	gag Glu	caa Gln	gag Glu	1099
ctg Leu 215	gag Glu	aag Lys	agt Ser	atg Met	aaa Lys 220	atg Met	cag Gln	caa Gln	gag Glu	gtg Val 225	gtg Val	gag Glu	atg Met	cgg Arg	aaa Lys 230	1147
aag Lys	aat Asn	gaa Glu	gaa Glu	ttc Phe 235	aag Lys	aaa Lys	ctt Leu	gct Ala	ctg Leu 240	gct Ala	gga Gly	ata Ile	GJÀ āāā	caa Gln 245	cct Pro	1195
gtg Val	aag Lys	aaa Lys	tca Ser 250	gtg Val	agc Ser	cag Gln	gtc Val	acc Thr 255	aaa Lys	tca Ser	gtt Val	gac Asp	ttc Phe 260	cac His	ttc Phe	1243
cgc Arg	aca Thr	gat Asp 265	gag Glu	cga Arg	atc Ile	aaa Lys	caa Gln 270	cat His	cct Pro	aag Lys	aac Asn	cag Gln 275	gag Glu	gaa Glu	tat Tyr	1291

															•		
		_			aca Thr		Glu			-		Pro				133	39
					gga Gly 300						Pro					138	37
		_		-	aca Thr		_	-		_						143	3.5
	_	_			gaa Glu	_				-						148	33
					aag Lys			Ile								153	31
			-		tgc Cys	_	_		-			_	_			157	19
					gct Ala 380											162	27
					aaa Lys											167	15
					ctt Leu											172	:3
					acc Thr			Ile								177	1
aaa Lys	aga Arg 440	atc Ile	cag Gln	gag Glu	cga Arg	gaa Glu 445	tca Ser	aag Lys	aag .Lys	aaa Lys	aca Thr 450	gag Glu	gat Asp	gaa Glu	cac His	181	.9
ttt Phe 455	gaa Glu	ttt Phe	cat His	tcc Ser	aga Arg 460	cct Pro	tgc Cys	cct Pro	act Thr	aag Lys 465	att Ile	ttg Leu	gaa Glu	gat Asp	gtt Val 470	186	7
gtg Val	ggt Gly	gtt Val	cct Pro	gaa Glu 475	aag Lys	aag Lys	gta Val	ctt Leu	cca Pro 480	atc Ile	acc Thr	gtc Val	ccc Pro	aag Lys 485	tca Ser	191	5
					aag Lys											196	3
gag Glu	gaa Glu	gag Glu 505	gac Asp	gaa Glu	ccg Pro	gta Val	gtg Val 510	ata Ile	aaa Lys	gct Ala	caa Gln	cct Pro 515	gtg Val	cca Pro	cat His	201	1
tat Tyr	ggg Gly 520	gtg Val	cct Pro	ttt Phe	aag Lys	ccc Pro 525	caa Gln	atc Ile	cca Pro	gag Glu	gca Ala 530	aga Arg	act Thr	gtg Val	gaa Glu	205	9
ata Ile 535	tgc Cys	cct Pro	ttc Phe	tcg Ser	ttt Phe 540	gat Asp	tct Ser	cga Arg	gac Asp	aaa Lys 545	gaa Glu	cgt Arg	cag Gln	tta Leu	cag Gln 550	210	7

•					•			
aag gag aag Lys Glu Lys	aaa ata Lys Ile 555	aaa gaa Lys Glu	ctg cag Leu Gln	aaa ggg Lys Gly 560	gag gtg Glu Val	ccc aag Pro Lys 565	ttc Phe	2155
aag gca ctt Lys Ala Leu	ccc ttg Pro Leu 570	cct cat Pro His	ttt gac Phe Asp 575	acc att Thr Ile	aac ctg Asn Leu	cca gag Pro Glu 580	aag Lys	2203
aag gta aag Lys Val Lys 585	Asn Val	acc cag Thr Gln	att gaa Ile Glu 590	cct ttc Pro Phe	tgc ttg Cys Leu 595	gag act Glu Thr	gac Asp	2251
aga aga ggt Arg Arg Gly 600	gct ctg Ala Leu	aag gca Lys Ala 605	cag act Gln Thr	tgg aag Trp Lys	cac cag His Gln 610	ctg gaa Leu Glu	gaa Glu	2299
gaa ctg aga Glu Leu Arg 615	cag cag Gln Gln	aaa gaa Lys Glu 620	gca gct Ala Ala	tgt ttc Cys Phe 625	aag gct Lys Ala	cgt cca Arg Pro	aac Asn 630	2347
acc gtc atc Thr Val Ile	tct cag Ser Gln 635	Glu Pro	ttt gtt Phe Val	ccc aag Pro Lys 640	aaa gag Lys Glu	aag aaa Lys Lys 645	Ser	2395
gtt gct gag Val Ala Glu	ggc ctt Gly Leu 650	tct ggt Ser Gly	tct cta Ser Leu 655	Val Gln	gaa cct Glu Pro	ttt cag Phe Gln 660	ctg Leu	2443
gct act gag Ala Thr Glu 665	Lys Arg							2491
gct gag gta Ala Glu Val 680	gaa gcc Glu Ala	cag aaa Gln Lys 685	gcc cag Ala Gln	cag ttg Gln Leu	gag gag Glu Glu 690	gcc aga Ala Arg	cta Leu	2539
cag gag gaa Gln Glu Glu 695 .	gag cag Glu Gln	aaa aaa Lys Lys 700	gag gag Glu Glu	ctg gcc Leu Ala 705	agg cta Arg Leu	cgg aga Arg Arg	gaa Glu 710	2587
ctg gtg cat Leu Val His	aag gca Lys Ala 715	Asn Pro	ata cgc Ile Arg	aag tac Lys Tyr 720	cag ggt Gln Gly	ctg gag Leu Glu 725	Ile	2635
aag tca agt Lys Ser Ser	gac cag Asp Gln 730	cct ctg Pro Leu	act gtg Thr Val 735	Pro Val	tct ccc Ser Pro	aaa tto Lys Phe 740	tcc Ser	2683
act cga ttc Thr Arg Phe 745	His Cys		agc tgtg	agctgc g	gataccgc	c cggcaa	tggg	2738
acctgctctt	aacctcaa	ac ctagg	accgt ct	tgctttgt	cattggg	cat ggag	agaacc	2798
catttctcca	gactttta	cc taccc	gtgcc tg	agaaagca	tacttga	caa ctgt	ggactc	2858
cagttttgtt	gagaattg	tt ttctt	acatt ac	taaggcta	ataatga	gat gtaa	ctcatg	2918
· aatgtctcga	ttagactc	ca tgtag	ttact to	ctttaaac	catcagc	cgg cctt	ttatat	2978
gggtcttcac	tctgacta	ga attta	gtctc tg	tgtcagca	cagtgta	atc tcta	ttgcta	3038
ttgcccctta	cgactctc	ac cctct	cccca ct	ttttttaa	aaatttt	aac caga	aaataa	3098
agatagttaa	atcctaag	at agaga	ttaag to	atggttta	aatgagg	aac aatc	agtaaa	3158
tcagattctg	tcctcttc	tc tgcat	accgt ga	atttatag	ttaagga	tcc cttt	gctgtg	3218

agggtagaaa acctcaccaa ctgcaccagt gaggaagaag actgcgtgga ttcatgggga
gcctcacage agecacgcag caggetetgg gtggggetge egttaaggea egttetttee
ttactggtgc tgataacaac agggaaccgt gcagtgtgca ttttaagacc tggcctggaa
taaatacgtt ttgtctttcc ctcaaaaaaa aaaaaaaaaa
aaaaaaaaa aa
<210> 34
<211> 747
<212> PRT
<213> NM_012112 C20orf1, chromosome 20 open reading frame 1
•
<400> 34
Met Ser Gln Val Lys Ser Ser Tyr Ser Tyr Asp Ala Pro Ser Asp Phe 1 5 10 15
Ile Asn Phe Ser Ser Leu Asp Asp Glu Gly Asp Thr Gln Asn Ile Asp 20 25 30
Ser Trp Phe Glu Glu Lys Ala Asn Leu Glu Asn Lys Leu Gly Lys 35 40 45
Asn Gly Thr Gly Gly Leu Phe Gln Gly Lys Thr Pro Leu Arg Lys Ala 50 55 60
Asn Leu Gln Gln Ala Ile Val Thr Pro Leu Lys Pro Val Asp Asn Thr 65 75 . 80
Tyr Tyr Lys Glu Ala Glu Lys Glu Asn Leu Val Glu Gln Ser Ile Pro 85 90 95
Ser Asn Ala Cys Ser Ser Leu Glu Val Glu Ala Ala Ile Ser Arg Lys 100 105 110
Thr Pro Ala Gln Pro Gln Arg Arg Ser Leu Arg Leu Ser Ala Gln Lys 115 120 125
Asp Leu Glu Gln Lys Glu Lys His His Val Lys Met Lys Ala Lys Arg 130 135 140
Cys Ala Thr Pro Val Ile Ile Asp Glu Ile Leu Pro Ser Lys Lys Met 145 150 155 160

Lys Val Ser Asn Asn Lys Lys Lys Pro Glu Glu Glu Gly Ser Ala His 165 170 175 Gln Asp Thr Ala Glu Lys Asn Ala Ser Ser Pro Glu Lys Ala Lys Gly 180 185 190

Arg His Thr Val Pro Cys Met Pro Pro Ala Lys Gln Lys Phe Leu Lys 195 200 205

Ser Thr Glu Glu Gln Glu Leu Glu Lys Ser Met Lys Met Gln Glu 210 215 220

Val Val Glu Met Arg Lys Lys Asn Glu Glu Phe Lys Lys Leu Ala Leu 225 230 235 240

Ala Gly Ile Gly Gln Pro Val Lys Lys Ser Val Ser Gln Val Thr Lys 245 250 255

Ser Val Asp Phe His Phe Arg Thr Asp Glu Arg Ile Lys Gln His Pro 260 265 270

Lys Asn Gln Glu Glu Tyr Lys Glu Val Asn Phe Thr Ser Glu Leu Arg 275 280 285

Lys His Pro Ser Ser Pro Ala Arg Val Thr Lys Gly Cys Thr Ile Val 290 295 300

Lys Pro Phe Asn Leu Ser Gln Gly Lys Lys Arg Thr Phe Asp Glu Thr 305 310 315 320

Val Ser Thr Tyr Val Pro Leu Ala Gln Gln Val Glu Asp Phe His Lys 325 330 335

Arg Thr Pro Asn Arg Tyr His Leu Arg Ser Lys Lys Asp Asp Ile Asn 340 345 350

Leu Leu Pro Ser Lys Ser Ser Val Thr Lys Ile Cys Arg Asp Pro Gln 355 360 . 365

Thr Pro Val Leu Gln Thr Lys His Arg Ala Arg Ala Val Thr Cys Lys 370 375 380

Ser Thr Ala Glu Leu Glu Ala Glu Glu Leu Glu Lys Leu Gln Gln Tyr 385 390 395 400

Lys Phe Lys Ala Arg Glu Leu Asp Pro Arg Ile Leu Glu Gly Pro 405 410 415

· Ile Leu Pro Lys Lys Pro Pro Val Lys Pro Pro Thr Glu Pro Ile Gly
420 425 430

Phe Asp Leu Glu Ile Glu Lys Arg Ile Gln Glu Arg Glu Ser Lys Lys 435 440 445

Lys Thr Glu Asp Glu His Phe Glu Phe His Ser Arg Pro Cys Pro Thr 450 455 460

Lys Ile Leu Glu Asp Val Val Gly Val Pro Glu Lys Lys Val Leu Pro 465 470 475 480

Ile Thr Val Pro Lys Ser Pro Ala Phe Ala Leu Lys Asn Arg Ile Arg 485 490 495

Met Pro Thr Lys Glu Asp Glu Glu Glu Asp Glu Pro Val Val Ile Lys 500 505 510

Ala Gln Pro Val Pro His Tyr Gly Val Pro Phe Lys Pro Gln Ile Pro 515 520 525

Glu Ala Arg Thr Val Glu Ile Cys Pro Phe Ser Phe Asp Ser Arg Asp 530 535 540

Lys Glu Arg Gln Leu Gln Lys Glu Lys Lys Ile Lys Glu Leu Gln Lys 545 550 555 560

Gly Glu Val Pro Lys Phe Lys Ala Leu Pro Leu Pro His Phe Asp Thr 565 570 575

Ile Asn Leu Pro Glu Lys Lys Val Lys Asn Val Thr Gln Ile Glu Pro 580 585 590

Phe Cys Leu Glu Thr Asp Arg Gly Ala Leu Lys Ala Gln Thr Trp 595 600 605

Lys His Gln Leu Glu Glu Glu Leu Arg Gln Gln Lys Glu Ala Ala Cys 610 615 620

Phe Lys Ala Arg Pro Asn Thr Val Ile Ser Gln Glu Pro Phe Val Pro 625 630 635 640

Lys Lys Glu Lys Lys Ser Val Ala Glu Gly Leu Ser Gly Ser Leu Val 645 650 655

Gln Glu Pro Phe Gln Leu Ala Thr Glu Lys Arg Ala Lys Glu Arg Gln 660 665 670

Glu Leu Glu Lys Arg Met Ala Glu Val Glu Ala Gln Lys Ala Gln Gln 675 680 685

Leu Glu Glu Ala Arg Leu Gln Glu Glu Glu Glu Lys Lys Glu Glu Leu 690 695 700

Ala Arg Leu Arg Arg Glu Leu Val His Lys Ala Asn Pro Ile Arg Lys 705 710 715 720

WO 2004/022778 PCT/AU2003/001166

Tyr Gln Gly Leu Glu Ile Lys Ser Ser Asp Gln Pro Leu Thr Val Pro 725 730 735

Val Ser Pro Lys Phe Ser Thr Arg Phe His Cys  $740 \hspace{1cm} \text{745}$ 

<210> 35

<211> 1246

<212> DNA

<213> NM\_001769 CD9

<220>

<221> CDS

<222> (112)..(795)

<223>

-10	0>	35														
			cago	cgcc	tg c	atct	gtat	c ca	gcgc	cagg	tcc	cgcc	agt	ccca	gctgcg	60
cgc	gccc	ccc	agtc	ccgc	ac c	cgtt:	cggc	c ca	ggct	aagt	tag	ccct	cac (		g ccg t Pro	117
										ctg Leu						165
										ctt Leu						213
										ttc Phe 45						261
										tat Tyr						309
										ggc Gly						357
										ttc Phe						405
										tgg Trp						453
										tac Tyr 125						501
aag	ctg	aaa	acc	aag	gat	gag	ccc	cag	cgg	gaa	acg	ctg	aaa	gcc	atc	549

Lys Leu Lys Thr Lys Asp Glu Pro Gln Arg Glu Thr Leu Lys Ala Ile 135 140 145	
cac tat gcg ttg aac tgc tgt ggt ttg gct ggg ggc gtg gaa cag ttt His Tyr Ala Leu Asn Cys Cys Gly Leu Ala Gly Gly Val Glu Gln Phe 150 160	597
atc tca gac atc tgc ccc aag aag gac gta ctc gaa acc ttc acc gtg Ile Ser Asp Ile Cys Pro Lys Lys Asp Val Leu Glu Thr Phe Thr Val 165 170 175	645
aag too tgt cot gat goo ato aaa gag gto tto gac aat aaa tto cac Lys Ser Cys Pro Asp Ala Ile Lys Glu Val Phe Asp Asn Lys Phe His 180 185 190	693
atc atc ggc gca gtg ggc atc ggc att gcc gtg gtc atg ata ttt ggc Ile Ile Gly Ala Val Gly Ile Gly Ile Ala Val Val Met Ile Phe Gly 195 200 205 210	741
atg atc ttc agt atg atc ttg tgc tgt gct atc cgc agg aac cgc gag Met Ile Phe Ser Met Ile Leu Cys Cys Ala Ile Arg Arg Asn Arg Glu 215 220 225	789
atg gtc tagagtcagc ttacatccct gagcaggaaa gtttacccat gaagattggt Met Val	845
gggatttttt gtttgtttgt tttgttttgt ttgttgtttg ttgtttgttt ttttgccad	t 905
aattttagta ttcattctgc attgctagat aaaágctgaa gttactttat gtttgtctt	t 965
taatgottoa ttoaatattg acatttgtag ttgagogggg ggtttggttt gotttggtt	t 1025
atattttttc agttgtttgt ttttgcttgt tatattaagc agaaatcctg caatgaaag	g 1085
tactatattt gctagactct agacaagata ttgtacataa aagaattttt ttgtcttta	a 1145
atagatacaa atgtetatea aetttaatea agttgtaaet tatattgaag acaatttga	t 1205
acataataaa aaattatgac aatgtcaaaa aaaaaaaaaa	1246
<210> 36	
<211> 228	
2010> DDM	

<213> NM\_001769 CD9

<400> 36

Met Pro Val Lys Gly Gly Thr Lys Cys Ile Lys Tyr Leu Leu Phe Gly 1 5 10 10 15

Phe Asn Phe Ile Phe Trp Leu Ala Gly Ile Ala Val Leu Ala Ile Gly

Leu Trp Leu Arg Phe Asp Ser Gln Thr Lys Ser Ile Phe Glu Gln Glu 35 40 45

Thr Asn Asn Asn Ser Ser Phe Tyr Thr Gly Val Tyr Ile Leu Ile 50 60

Gly Ala Gly Ala Leu Met Met Leu Val Gly Phe Leu Gly Cys Cys Gly 65 70 75 80

Ala Val Gln Glu Ser Gln Cys Met Leu Gly Leu Phe Phe Gly Phe Leu 85 90 95

Leu Val Ile Phe Ala Ile Glu Ile Ala Ala Ala Ile Trp Gly Tyr Ser 100 105 110

His Lys Asp Glu Val Ile Lys Glu Val Gln Glu Phe Tyr Lys Asp Thr 115 120  $\cdot$  125

Tyr Asn Lys Leu Lys Thr Lys Asp Glu Pro Gln Arg Glu Thr Leu Lys 130 135 140

Ala Ile His Tyr Ala Leu Asn Cys Cys Gly Leu Ala Gly Gly Val Glu 145 150 155 160

Gln Phe Ile Ser Asp Ile Cys Pro Lys Lys Asp Val Leu Glu Thr Phe 165 170 175

Thr Val Lys Ser Cys Pro Asp Ala Ile Lys Glu Val Phe Asp Asn Lys 180 185 190

Phe His Ile Ile Gly Ala Val Gly Ile Gly Ile Ala Val Val Met Ile 195 200 205

Phe Gly Met Ile Phe Ser Met Ile Leu Cys Cys Ala Ile Arg Arg Asn 210 215 . 220

Arg Glu Met Val 225

<210> 37

<211> 7557

<212> DNA

<213> NM\_020859 ShrmL, Shroom-related protein

<220>

<221> CDS

<222> (958)..(6942)

<223>

<400> 37

				* .			
ggcagtgctt	actgtgcaga	aagģttgca	a tgcctc	actt tgo	cttaatc	catgttcctc	120
taacttcctt	actctttctc	tcaagagga	g gcaagt	ggct gtg	iacāācċā	cagcagtggc	180
tgatcatcac	tgaaaatacc	aaagaaaag	a actgag	ctgc ctc	cttcata	ttttttccat	240
tgaggattaa	tttaccgtgc	tttttcatt	t tctcta	catc ctg	caaaagt	tttttctct	300
cctaagaaac	aaactatgaa	ctgattgtt	g aaaaaa	agaa gta	aaaagtt	ttagcacage	360
ttctctgtct	cttcgggaca	agttagaaa	a ttctga	agtg ago	cgaagca:	tagtaagtgc	420
tttctttctt	tttaagctac	ttctgggga	g ggagga	ggct att	gtaatgg	taaatttcac	480
tccgagagga	agaaagggtt	gataatcaa	t caaaaa	tgag gta	ttccttt	gagtatttgt	540
gattttctta	ctatattgta	agatgcttt	t aatttt	ctct gta	aaatagg	cagaaatggt	600
tttagtgtgt ·	gtatgtgtga	aataaaagc	t cagaaa	agca atc	ttcagag	cgccactgaa	660
ggaagttttg	acgaacggag	tagagatgt	a taccac	ttgg ggg	cttcagt	gagaacccag	720
aattcctgga	ggaggattta	cattcagaa	a tgttga	agtg aaa	attcctt	ctggttcagc	780
atcttggagt	tcagcttgga	agaacattti	t acgtat	ggaa gaa	tttgctt	ctccaaacct	840
ctcttttggc (	cattgtgtgt	cctgaaggat	t gggaca	actt gtg	ctgtaga	agcactgctt	900
gcctgagttt	gcttcaggca	ttttaaatti	t aacttg	aggg atc	atgtgtt	tggcatg	957
atg agg acc Met Arg Thr 1	act gaa ga Thr Glu As 5	c ttc cac p Phe His	aag cct Lys Pro 10	agt gcc Ser Ala	aca tta Thr Leu	aac tct Asn Ser 15	1005
aac acg gcc Asn Thr Ala	acc aag gg Thr Lys Gl 20	a agg tac y Arg Tyr	att tat Ile Tyr 25	ctg gag Leu Glu	gca ttc Ala Phe 30	ctg gag Leu Glu	1053
gga gga gct Gly Gly Ala 35	ccc tgg gg Pro Trp Gl	t ttt act y Phe Thr 40	cta aag Leu Lys	ggt ggc	ctg gag Leu Glu 45	cac gga His Gly	1101
gaa cca tta Glu Pro Leu 50	atc atc tc Ile Ile Se	t aag gtc r Lys Val 55	gaa gaa Glu Glu	e0 Gly Gly Ggg ggc	aaa gca Lys Ala	gac acc Asp Thr	1149
ctg agc tcc Leu Ser Ser 65	aaa ctg ca Lys Leu Gl 70	g gct ggg n Ala Gly	gat gag Asp Glu	gtt gtg Val Val 75	cac atc His Ile	aat gag Asn Glu 80	1197
gtg act ctg Val Thr Leu	agc agc tco Ser Ser Se: 85	c aga aag r Arg Lys	gag gca Glu Ala 90	gtt tcc Val Ser	ctg gtg Leu Val	aaa gga Lys Gly 95	1245
tcc tac aag Ser Tyr Lys		g Leu Val					1293
cca ggc cat Pro Gly His	gca gat act Ala Asp Thi	ggt gcc Gly Ala 120	tct aac Ser Asn	ttc gtc Phe Val	agc cca Ser Pro 125	gaa cac Glu His	1341
ctc acc tct Leu Thr Ser 130							1389
aaa ctt cgg	ctg aag cac	agg tot a	agt gag	cct gcạ	ggc cga	cct cac	1437

WO 2004/022778 PCT/AU2003/001166

							•									
Lys 145	Leu	Arg	Leu	Lys	His 150	Arg	Ser	Ser	GIu	Pro 155	Ala	Gly	Arg	Pro	His 160	
					Lys	tct Ser										1485
						atg Met										1533
						agt Ser										1581
						cag Gln 215										1629
						tac .Tyr										1677
						cag Gln										1725
						tcc Ser										1773
						cgg Arg										1821
	_	-				ctc Leu 295	-		_		_	_	-			1869
						gac Asp										1917
						ctg Leu										1965
						tat Tyr										2013
						tcc Ser										2061
						ctt Leu 375										2109
						tac Tyr										2157
agc Ser	tcc Ser	tgg Trp	tct Ser	agc Ser 405	ctt Leu	gat Asp	cag Gln	aaa Lys	cgg Arg 410	ctc Leu	tgc Cys	cgg Arg	cct Pro	cag Gln 415	gca Ala	2205
aac	tct	tta	ggc	tcc	ctg	aag	tct	cca	ttc	ata	gag	gag	cag	ctg	cat	2253

	•															
Asr	n Se	r Lei	u Gly 420	y Se:	r Lev	ı Lys	Sei	Pro 425		e Ile	e Gl	u Gli	u Gl 43		u His	
act Thi	gto Val	t Lei 43	ı Glı	g aaq ı Ly:	g agt s Ser	cca Pro	gag Glu 440	ı Asr	ago Sei	Pro	c cc Pr	a gto o Val 44!	l Ly	g cc s Pr	c aag o Lys	2301
His	450	ı Tyı )	t Thi	c Glı	n Lys	455	Gln	Pro	Gl	/ Glr	1 Pro	o Lei 0	ı Le	u Pr	g acc o Thr	2349
Ser 465	: Ile	э Туг	: Ala	ı Val	470	Ser	Leu	ı Glu	Pro	475	Phe	e Ala	a Gl	n Va	g cct l Pro 480	2397
Gln	Pro	Ser	· Val	. Sei 485	Ser	Asn	Gly	Met	Leu 490	Tyr	Pro	o Ala	ı Lei	1 Ala 49	_	2445
GIu	Ser	: Gly	500	: Ile	e Ala	Pro	Gln	Gly 505	Ala	Cys	Ası	Lys	510	: Ala	t acc a Thr	2493
11e	Asp	515	Asn	Gly	Asn	Gln	Asn 520	Gly	Ser	Gly	Arg	9 Pro 525	Gly	, Phe	gcc Ala	2541
Phe	Суs 530	Gln	Pro	Leu	Glu	His 535	Asp	Leu	Leu	Ser	9rc 540	Val	Glu	Lys	g aaa S Lys	2589
545	Glu	Ala	Thr	Ala	Lys 550	Tyr	Val	Pro	Ser	Lys 555	Val	. His	Phe	Суз	560	2637
Val	Pro	Glu	Asn	Glu 565		Asp	Ala	Ser	Leu 570	ГÀЗ	Arg	His	Leu	Thr 575	Pro	2685
Pro	Gln	Gly	Asn 580	Ser	cca Pro	His	Ser	Asn 585	Glu	Arg	<sub>.</sub> Lys	Ser	Thr 590	His	Ser	2733
Asn	Lys	Pro 595	Ser	Ser	cat His	Pro	His 600	Ser	Leu	ГÀЗ	Суз	Pro 605	Gln	Ala	Gln	2781
Ala	Trp 610	Gln	Ala	Gly	gaa Glu	Asp 615	Lys	Arg	Ser	Ser	Arg 620	Leu	Ser	Glu	Pro	2829
625	Glu	Gly	Asp	Phe	cag Gln 630	Glu	Asp	His	Asn	A1a 635	Asn	Leu	Trp	Arg	Arg 640	2877
Leu	Glu	Arg	Glu	Gly 645	cta Leu	Gly	Gln	Ser	Leu 650	Ser	Gly	Asn	Phe	Gly 655	Lys	2925
Thr	Lys	Ser	Ala 660	Phe	tca Ser	Ser :	Leu	Gln 665	Asn	Ile	Pro	Glu	Ser 670	Leu	Arg	2973
aga Arg	His	agc Ser 675	agc Ser	ctg Leu	gag Glu	Leu (	ggc Gly 680	cgg Arg	gga Gly	acc Thr	cag Gln	gag Glu 685	ggt Gly	tac Tyr	ccc Pro	3021
ggg (	ggc	agg	ccc	acc	tgt (	gca (	gtc	aac a	acc	aag	gca	gaa	gac	cct	ggg	3069

Gly	Gly 690	Arg	Pro	Thr	Суз	Ala 695	Val	Asn	Thr	Lys	Ala 700	Glu	Asp	Pro	Gly	
					gac Asp 710											3117
					G1y ggg											3165
					gaa Glu		Pro									3213
	_	_			agg Arg						-	_	_	_		3261
					GJ À âàà											3309
					cga Arg 790											3357
					cat His											3405
					aat Asn										cgt Arg	3453
					gaa Glu											3501
					ttg Leu											3549
					gcg Ala 870											3597
					cgt Arg											3645
					gag Glu											3693
					ccc Pro											3741
					gtc Val											3789
					ccc Pro 950											3837
tcc	tcg	gcc	cac	gtg	ggg	ctg	cgg	agc	ccc	gag	gcg	tcg	gcc	tcc	gcc	3885

-				•		•									
Ser	Ser	Ala	His	Val 965	Gly	Leu A	rg S		ro G 70 .		la Se	r Al	a Se 97	r Ala 5	
							rg H						a Gl	n Glà	3933
	Leu					Pro P					Arg G			cgc cg Arg Ar	
_	-	Thr			-	aag Lys 1015	Lys	-			_	Glu			4026
	atg Met 1025	Asn				atc Ile 1030	Val					Pro	-	ccc Pro	4071
	ggc Gly 1040	Pro				999 Gly 1045	Met					Ser			4116
		Arg				ttc Phe 1060	Glu					Ala			4161
_	ctc Leu 1070	Ser				ccc Pro 1075	Glu					Gln			4206
		Ala				cag Gln 1090	Arg					Arg			4251
	gcc Ala 1100	Ala				ctc Leu 1105	Gln					Leu		gag Glu	4296
-	gcc Ala 1115					ctc Leu 1120	Gln						ctc Leu	-	4341
	tcc Ser 1130			_	_	gcc Ala 1135	Ser	-	-	_		-	cgg		4386
						agg Arg 1150	Glu								4431
						gct Ala 1165									4476
						ggg Gly 1180	Glu								4521
						aca Thr 1195									4566
acc Thr	ccc Pro 1205	agg Arg	gag Glu	cca Pro	tcc Ser	tcc Ser 1210	tgg Trp	GJA aaa	gcc Ala	agg Arg	gcc Ala 1215	Gly Ggg	aag Lys	tcc Ser	4611
atg	tcg	gcc	gag	gac	ctg	ctg	gaa	cgc	tcg	gac	gtc	ctt	gcg	ggc	4656

Met	Ser 1220	Ala	Glu	a Asp	Lev	Leu 1225	Glu	Arg	Sei	Asr	Val 1230		Ala	Gly	
cct Pro	gtc Val 1235	His	gtg Val	ago Arc	, tcc , Ser	agg Arg 1240	Ser	tct	Pro	gcc Ala	acc Thr 1245	Ala	gac Asp	aag Lys	4701
Arg	cag Gln 1250	Asp	gtg Val	ctt Leu	ttg Leu	ggg Gly 1255	Gln	gac Asp	agt Ser	ggc Gly	ttt Phe 1260	Gly	ctt Leu	gtg Val	4746
aag Lys	gat Asp 1265	Pro	tgt Cys	tat Tyr	ttg Leu	gct Ala 1270	Gly	cct Pro	gga Gly	tct Ser	agg Arg 1275	Ser	ctc Leu	agt Ser	4791
tgt Cys	tca Ser 1280	Glu	aga Arg	Gly	caa Gln	gaa Glu 1285	Glu	atg Met	ctg Leu	ctg Leu	ctc Leu 1290	Phe	cac	cat His	4836
	acc Thr 1295	Pro	cgt Arg	tgg Trp	ggt	ggt Gly 1300	Ser	ggc	tgc Cys	aaa Lys	gcc Ala 1305	Ile	ggt Gly	gat Asp	4881
tcc Ser	tcc Ser 1310	Val	cct Pro	agt Ser	gaa Glu	tgt Cys 1315	Pro	gga Gly	acc Thr	ctg Leu	gac Asp 1320	His	cag Gln	agg Arg	4926
caa Gln	gcc Ala 1325	Ser	agg Arg	aca Thr	ccc Pro	tgc Cys 1330	ccc Pro	agg Arg	cca Pro	cca Pro	ctg Leu 1335	Ala	gga Gly	acg Thr	4971
caa Gln	ggg Gly 1340	Leu	gtc Val	aca Thr	gac Asp	acc Thr 1345	agg Arg	gct Ala	gca Ala	ccc Pro	ctg Leu 1350	Thr		att Ile	5016
ggc Gly	acc Thr 1355	cct Pro	ctg Leu	cct Pro	tca Ser	gcc Ala 1360	att Ile	ccc Pro	tct Ser	Gly	tac Tyr 1365	Cys		cag Gln	5061
	ggt Gly 1370	cag Gln	aca Thr	ggg Gly	cga Arg	cag Gln 1375	cct Pro	ctc Leu	ccg Pro	ccc Pro	tac Tyr 1380			gcc Ala	5106
Met	Met 1385	His	Arg	Ser	Asn	Gly 1390	His	Thr	Leu	Thr	1395	Pro	Pro	Gly	5151
cca Pro	aga Arg 1400	ggc Gly	tgt Cys	gag Glu	Gly	gat Asp 1405	ggc Gly	cca Pro	gag Glu	cat His	ggg Gly 1410	gta Val	gaa Glu	gag Glu	5196
gga Gly	acg Thr 1415	agg Arg	aag Lys	agg Arg	gtc Val	tcg Ser 1420	ctg Leu	cct Pro	cag Gln	tgg Trp	cca Pro 1425	cct Pro	cct Pro	tct Ser	5241
cga Arg	gca Ala 1430	aag Lys	tgg Trp	gcc Ala	cac His	gca Ala 1435	gcc Ala	aga Arg	gag Glu	gac Asp	agc Ser 1440	ctt Leu	cct Pro	gag Glu	5286
gaa Glu	tcc Ser 1445	tca Ser	gcc Ala	cct Pro	gat Asp	ttt Phe 1450	gca Ala	aac Asn	ctg Leu	aag Lys	cac His 1455	tat Tyr	caa Gln	aaa Lys	5331
Gln	cag Gln 1460	agt Ser	ctt Leu	cca Pro	Ser	tta Leu 1465	tgc Cys	agc Ser	act Thr	tct Ser	gac Asp 1470	cca Pro	gac Asp	aca Thr	5376
cct	ctt	ggg	gcc	ccg	agc	act	сса	ggg	agg	atc	tcc	ctc	cga	ata	5421

Pro	Leu 1475	Gly	Ala	Pro	Ser	Thr 1480		Gļy	Arg	Ile	Ser 1485		Arg	Ile	
	gag Glu 1490					gac Asp 1495									5466
	gac Asp 1505					agg Arg 1510									5511
_	ccc Pro 1520					ctt Leu 1525	Pro								5556
-	cag Gln 1535	_		-		tat Tyr 1540	Ser	_	_	_					5601
	ccc Pro 1550			-	_	gag Glu 1555	Ala	_	_	_	-	Glu	_		5646
	ggg Gly 1565						.Asn					Val			5691
-	agg Arg 1580					cct Pro 1585									5736
	aca Thr 1595						Gln								5781
_	gag Glu 1610					tcc Ser 1615									5826
-	ggg Gly 1625	tct Ser	ctt Leu	ggt Gly	GJ Ā āāā	cag Gln 1630	cca Pro	gca <sup>,</sup> Ala	ccc Pro	atc Ile	cag Gln 1635	act Thr	caa Gln	agc Ser	5871
	agc Ser 1640					agt Ser 1645									5916
	agt Ser 1655					aag Lys 1660									5961
	ttg Leu 1670														6006
	ttg Leu 1685														6051
	ggt Gly 1700														6096
gta Val	aag Lys 1715	agg Arg	aag Lys	gcc Ala	ata Ile	cag Gln 1720	aga Arg	act Thr	gtc Val	agc Ser	tct Ser 1725	tca Ser	gga Gly	tgt Cys	6141
gaa	ggc	aag	agg	aat	gaa	gac	aag	gaa	gca	gtg	agc	atg	ttg	gtt	6186

WO 2004/022778

Gli	ı Gly 1730	Ly:	s Ar	g Ası	ı Glu	1 Asp 1735		S Glı	a Ala	va:	l Ser 174		Le:	ı Val	
aac Ası	tgc Cys 1745	Pro	t gc	c tac a Tyr	tac Tyr	agt Ser 1750	Va]	j tot L Sei	gct Ala	CC Pro	c aag b Lys 175	Ala	gaç a Glu	g cta 1 Leu	6231
cto Lei	aac Asn 1760	Ly	a ato	c aaa e Lys	gag Glu	atg Met 1765	Pro	a gca Ala	ı gaa	gto Va	g aat l Asn 1770	Glu	g gaa n Glu	ı gag ı Glu	6276
gaa Gli	cag Gln 1775	Ala	a gat a Asp	gto Val	aat Asn	gaa Glu 1780	Lys	aag Lys	gct Ala	gaq Glu	teu 1785	Il∈	gga Gly	agt Ser	6321
cto Lev	acc Thr 1790	His	aaq Lys	ctg Lev	gag Glu	acc Thr 1795	Leu	cag Gln	gag Glu	gcg	g aag Lys 1800	Gly	g ago Ser	ctg Leu	6366
cto Lev	acg Thr 1805	Asp	ato Ile	aag Lys	ctc Leu	aac Asn 1810	Asn	gcc Ala	ctg Leu	Gl <sup>7</sup>	gaa Glu 1815	Glu	gtg Val	gag Glu	6411
gct Ala	ctg Leu 1820	Ile	ago Ser	gag Glu	ctc Leu	tgc Cys 1825	Lys	Pro	aat Asn	gaç	ttt Phe 1830	Asp	aag Lys	tat Tyr	6456
agg Arg	atg Met 1835	Phe	ata Ile	Gly	gat Asp	ttg Leu 1840	Asp	aag Lys	gtg Val	gtc Val	aac Asn 1845	Leu	ctg Leu	ctc Leu	6501
tcc Ser	ctc Leu 1850	Ser	. GJÀ . aàà	cgt Arg	cta Leu	gcc Ala 1855	Arg	gtt Val	gag Glu	aat Asn	gtc Val 1860	Leu	agc Ser	ggc Gly	6546
ctt Leu	ggt Gly 1865	Glu	gat Asp	gcc Ala	agt Ser	aat Asn 1870	Glu	gaa Glu	agg Arg	agc Ser	tct Ser 1875	Leu	tac Tyr	gag Glu	6591
aaa Lys	agg Arg 1880	Lys	atc Ile	ctg Leu	gct Ala	ggt Gly 1885	cag Gln	cat His	gag Glu	gat Asp	gcc Ala 1890	Arg	gag Glu	ctg Leu	6636
aag Lys	gag Glu 1895	Asn	ctg Leu	gat Asp	cgc Arg	agg Arg 1900	gag Glu	cga Arg	gta Val	gtg Val	ctg Leu 1905	ggc	atc Ile	ttg Leu	6681
gcc Ala	aat Asn 1910	tac Tyr	ctt Leu	tca Ser	gag Glu	gag Glu 1915	cag Gln	ctc Leu	cag Gln	gac Asp	tac Tyr 1920	cag Gln	cac His	ttc Phe	6726
gtg Val	aaa Lys 1925	atg Met	aag Lys	tcc Ser	acg Thr	ctc Leu 1930	ctc Leu	att Ile	gag Glu	caa Gln	cgg Arg 1935	aag Lys	ctg Leu	gat Asp	6771
gac Asp	aag Lys 1940	atc Ile	aag Lys	ctg Leu	ggc Gly	cag Gln 1945	gag Glu	cag Gln	gtc Val	aag Lys	tgt Cys 1950	ctg Leu	ctg Leu	gag Glu	6816
agc Ser	ctg Leu 1955	ccc Pro	tca Ser	gat Asp	Phe	att Ile 1960	ccc Pro	aag. Lys	gct Ala	GJ À aaa	gcc Ala 1965	ctg Leu	gct Ala	ctg Leu	6861
ccc Pro	cca Pro 1970	aac Asn	ctc Leu	acg Thr	Ser	gag Glu 1975	ccc Pro	att Ile	cct Pro	gct Ala	ggg Gly 1980	ggc Gly	tgt Cys	act Thr	6906
ttc	agt	ggt	att	ttc	cca	aca	tta	acc	tct	cca	ctt	taac	ctct	tc	6952

Phe Ser Gly Ile Phe Pro Thr Leu Thr Ser Pro Leu 1985 1990 1995
taaaataccc aaccaaaaga tcactgtttc tctcaacact atttaatctg aaaaatgttt
cagtacaaac cactgtttga actatctggg ttattggtgt ttgttcctga tgaaaggaaa
aaaattotot ocaggaggaa goottittoo tiottgooot tootgattga tottotgaga
gctcgaatgc tgctggacac gtaccccttt ctattattac tttgtagtag aaagaaagtt
aatgaaactg agaactgatt ggagggtgtt tgatcattta gtttttaaca ggctgaggca
acatggatca gtgtgtgtcc ccctcaggaa tgtatccaca gtggccttcc ttgctggtgg
gcagtgtatc ctgatggcag ggtacaagta ccattaatga agggtctgca acataaagcc
ttaaaaagac acacactaag aaaactgtaa aaccttgaac attgttattt atatttttta
aaatggaaaa gatcactatg titgttgtgc taaccactta titgattctg tittgtggtg
gacatagatg attacgtttg agctttgtat tttgtgaaaa ccttaatgaa atgaattcca
aagat
<210> 38
<211> 1995
<212> PRT
<213> NM_020859 ShrmL, Shroom-related protein
<400> 38
Met Arg Thr Thr Glu Asp Phe His Lys Pro Ser Ala Thr Leu Asn Ser  1 10 15
Asn Thr Ala Thr Lys Gly Arg Tyr Ile Tyr Leu Glu Ala Phe Leu Glu
Asn Thr Ala Thr Lys Gly Arg Tyr Ile Tyr Leu Glu Ala Phe Leu Glu 20 25 30  Gly Gly Ala Pro Trp Gly Phe Thr Leu Lys Gly Gly Leu Glu His Gly
Asn Thr Ala Thr Lys Gly Arg Tyr Ile Tyr Leu Glu Ala Phe Leu Glu 20 25 30  Gly Gly Ala Pro Trp Gly Phe Thr Leu Lys Gly Gly Leu Glu His Gly 35 40  Glu Pro Leu Ile Ile Ser Lys Val Glu Glu Gly Gly Lys Ala Asp Thr
Asn Thr Ala Thr Lys Gly Arg Tyr Ile Tyr Leu Glu Ala Phe Leu Glu 20  Gly Gly Ala Pro Trp Gly Phe Thr Leu Lys Gly Gly Leu Glu His Gly 35  Glu Pro Leu Ile Ile Ser Lys Val Glu Glu Gly Gly Lys Ala Asp Thr 50  Leu Ser Ser Lys Leu Gln Ala Gly Asp Glu Val Val His Ile Asn Glu
Asn Thr Ala Thr Lys Gly Arg Tyr Ile Tyr Leu Glu Ala Phe Leu Glu 20  Gly Gly Ala Pro Trp Gly Phe Thr Leu Lys Gly Gly Leu Glu His Gly 35  Glu Pro Leu Ile Ile Ser Lys Val Glu Glu Gly Gly Lys Ala Asp Thr 50  Leu Ser Ser Lys Leu Gln Ala Gly Asp Glu Val Val His Ile Asn Glu 75  Val Thr Leu Ser Ser Ser Arg Lys Glu Ala Val Ser Leu Val Lys Gly

Leu Thr Ser Gly Pro Gln His Arg Lys Ala Ala Trp Ser Gly Gly Val 130 135 140

Lys Leu Arg Leu Lys His Arg Ser Ser Glu Pro Ala Gly Arg Pro His 145 150 155 160

Ser Trp His Thr Thr Lys Ser Gly Glu Lys Gln Pro Asp Ala Ser Met 165 170 175

Met Gln Ile Ser Gln Gly Met Ile Gly Pro Pro Trp His Gln Ser Tyr 180 185 190

His Ser Ser Ser Ser Thr Ser Asp Leu Ser Asn Tyr Asp His Ala Tyr 195 200 205

Leu Arg Arg Ser Pro Asp Gln Cys Ser Ser Gln Gly Ser Met Glu Ser 210 215 220

Leu Glu Pro Ser Gly Ala Tyr Pro Pro Cys His Leu Ser Pro Ala Lys 225 230 235 240

Ser Thr Gly Ser Ile Asp Gln Leu Ser His Phe His Asn Lys Arg Asp 245 250 255

Ser Ala Tyr Ser Ser Phe Ser Thr Ser Ser Ser Ile Leu Glu Tyr Pro 260 265 270

His Pro Gly Ile Ser Ala Arg Glu Arg Ser Gly Ser Met Asp Asn Thr 275 280 285

Ser Ala Arg Gly Gly Leu Leu Glu Gly Met Arg Gln Ala Asp Ile Arg 290 295 300

Tyr Val Lys Thr Val Tyr Asp Thr Arg Arg Gly Val Ser Ala Glu Tyr 305 310 315 320

Glu Val Asn Ser Ser Ala Leu Leu Leu Gln Gly Arg Glu Ala Arg Ala 325 330 335

Ser Ala Asn Gly Gln Gly Tyr Asp Lys Trp Ser Asn Ile Pro Arg Gly 340 345 350

Lys Gly Val Pro Pro Pro Ser Trp Ser Gln Gln Cys Pro Ser Ser Leu 355 360 365

Glu Thr Ala Thr Asp Asn Leu Pro Pro Lys Val Gly Ala Pro Leu Pro 370 375 380

Pro Ala Arg Ser Asp Ser Tyr Ala Ala Phe Arg His Arg Glu Arg Pro 385 390 395 400

Ser Ser Trp Ser Ser Leu Asp Gln Lys Arg Leu Cys Arg Pro Gln Ala 405 410 415

Asn Ser Leu Gly Ser Leu Lys Ser Pro Phe Ile Glu Glu Gln Leu His 420 425 430

Thr Val Leu Glu Lys Ser Pro Glu Asn Ser Pro Pro Val Lys Pro Lys 435 440 445

His Asn Tyr Thr Gln Lys Ala Gln Pro Gly Gln Pro Leu Leu Pro Thr 450 455 460

Ser Ile Tyr Ala Val Pro Ser Leu Glu Pro His Phe Ala Gln Val Pro 465 470 475 480

Gln Pro Ser Val Ser Ser Asn Gly Met Leu Tyr Pro Ala Leu Ala Lys 485 490 495

Glu Ser Gly Tyr Ile Ala Pro Gln Gly Ala Cys Asn Lys Met Ala Thr 500 505 510

Ile Asp Glu Asn Gly Asn Gln Asn Gly Ser Gly Arg Pro Gly Phe Ala 515 520 525

Phe Cys Gln Pro Leu Glu His Asp Leu Leu Ser Pro Val Glu Lys Lys . 530 535 540

Pro Glu Ala Thr Ala Lys Tyr Val Pro Ser Lys Val His Phe Cys Ser 545 550 555 560

Val Pro Glu Asn Glu Glu Asp Ala Ser Leu Lys Arg His Leu Thr Pro 565 570 575

Pro Gln Gly Asn Ser Pro His Ser Asn Glu Arg Lys Ser Thr His Ser 580 585 590

Asn Lys Pro Ser Ser His Pro His Ser Leu Lys Cys Pro Gln Ala Gln 595 600 605

Ala Trp Gln Ala Gly Glu Asp Lys Arg Ser Ser Arg Leu Ser Glu Pro 610 620

Trp Glu Gly Asp Phe Gln Glu Asp His Asn Ala Asn Leu Trp Arg Arg 625 630 635 640

Leu Glu Arg Glu Gly Leu Gly Gln Ser Leu Ser Gly Asn Phe Gly Lys 645 650 655

Thr Lys Ser Ala Phe Ser Ser Leu Gln Asn Ile Pro Glu Ser Leu Arg
660 665 670

Arg His Ser Ser Leu Glu Leu Gly Arg Gly Thr Gln Glu Gly Tyr Pro 675 680 685

Gly Gly Arg Pro Thr Cys Ala Val Asn Thr Lys Ala Glu Asp Pro Gly 690 695 700

Arg Lys Ala Ala Pro Asp Leu Gly Ser His Leu Asp Arg Gln Val Ser 705 710 715 720

Thr Asp Pro Ser Pro Glu Glu Pro Pro Ala Pro Ser His Pro His Thr 740 745 750

Ser Ser Leu Gly Arg Arg Gly Pro Gly Pro Gly Ser Ala Ser Ala Leu 755 760 765

Gln Gly Phe Gln Tyr Gly Lys Pro His Cys Ser Val Leu Glu Lys Val 770 775 780

Ser Lys Phe Glu Gln Arg Glu Gln Gly Ser Gln Arg Pro Ser Val Gly 785 790 795 800

Gly Ser Gly Phe Gly His Asn Tyr Arg Pro His Arg Thr Val Ser Thr 805 810 815

Ser Ser Thr Ser Gly Asn Asp Phe Glu Glu Thr Lys Ala His Ile Arg 820 825 830

Phe Ser Glu Ser Ala Glu Pro Leu Gly Asn Gly Glu Gln His Phe Lys 835 840 845

Asn Gly Glu Leu Lys Leu Glu Glu Ala Ser Arg Gln Pro Cys Gly Gln 850 855 860

Gln Leu Ser Gly Gly Ala Ser Asp Ser Gly Arg Gly Pro Gln Arg Pro 865 870 875 880

Asp Ala Arg Leu Leu Arg Ser Gln Ser Thr Phe Gln Leu Ser Ser Glu 885 890 895

Pro Glu Arg Glu Pro Glu Trp Arg Asp Arg Pro Gly Ser Pro Glu Ser 900 905 910

Pro Leu Leu Asp Ala Pro Phe Ser Arg Ala Tyr Arg Asn Ser Ile Lys 915 920 925

Asp Ala Gln Ser Arg Val Leu Gly Ala Thr Ser Phe Arg Arg Arg Asp .930 935 940 Leu Glu Leu Gly Ala Pro Val Ala Ser Arg Ser Trp Arg Pro Arg Pro 945 950 955

Ser Ser Ala His Val Gly Leu Arg Ser Pro Glu Ala Ser Ala Ser Ala 965 970 975

Ser Pro His Thr Pro Arg Glu Arg His Ser Val Thr Pro Ala Glu Gly 980 985 990

Asp Leu Ala Arg Pro Val Pro Pro Ala Ala Arg Arg Gly Ala Arg Arg 995 1000 1005

Arg Leu Thr Pro Glu Gln Lys Lys Arg Ser Tyr Ser Glu Pro Glu 1010 1015 1020

Lys Met Asn Glu Val Gly Ile Val Glu Glu Ala Glu Pro Ala Pro 1025 1030 1035

Leu Gly Pro Gln Arg Asn Gly Met Arg Phe Pro Glu Ser Ser Val 1040 1045 1050

Ala Asp Arg Arg Leu Phe Glu Arg Asp Gly Lys Ala Cys Ser 1055 1060 : 1065

Thr Leu Ser Leu Ser Gly Pro Glu Leu Lys Gln Phe Gln Gln Ser 1070 1075 1080

Ala Leu Ala Asp Tyr Ile Gln Arg Lys Thr Gly Lys Arg Pro Thr 1085 1090 1095

Ser Ala Ala Gly Cys Ser Leu Gln Glu Pro Gly Pro Leu Arg Glu 1100 1105 1110

Arg Ala Gln Ser Ala Tyr Leu Gln Pro Gly Pro Ala Ala Leu Glu 1115 1120 1125

Gly Ser Gly Leu Ala Ser Ala Ser Ser Leu Ser Ser Leu Arg Glu 1130 1135 1140

Pro Ser Leu Gln Pro Arg Arg Glu Ala Thr Leu Leu Pro Ala Thr 1145 1150 1155

Val Ala Glu Thr Gln Gln Ala Pro Arg Asp Arg Ser Ser Ser Phe 1160 1165 1170

Ala Gly Gly Arg Arg Leu Gly Glu Arg Arg Arg Gly Asp Leu Leu 1175 1180 1185

Ser Gly Ala Asn Gly Gly Thr Arg Gly Thr Gln Arg Gly Asp Glu 1190 1195 1200

	Pro 1205	Arg	Glu <sub>.</sub>	Pro	Ser	Ser 1210	Trp	Gly	Ala	Arg	Ala 1215	Gly	Lys	Ser
Met	Ser 1220	Ala	Glu	Asp ·	Leu	Leu 1225	Glu	Arg	Ser	Asp	Val 1230	Leu	Ala	Gly
Pro	Vál 1235		Val	Arg	Ser	Arg 1240	Ser	Ser	Pro	Ala	Thr 1245	Ala	Asp	Lys
Arg	Gln 1250		Val	Leu	Leu	Gly 1255		Asp	Ser	Gly	Phe 1260	Gly	Leu	Val
Lys	Asp 1265		Суѕ	Tyr	Leu	Ala 1270		Pro	Gly	Ser	Arg 1275	Ser	Leu	Ser
	Ser 1280		Arg	Gly	GÌn	Glu 1285		Met	Leu	Leu	Leu 1290	Phe	His	His
Leu	Thr 1295		Arg	Trp	Gly	Gly 1300	Ser	Gly	Суз	Lys	Ala 1305	Ile	Gly	Asp
Ser	Ser 1310		Pro	Ser	Glu	Cys 1315	Pro	Gly	Thr	Leu	Asp 1320	His	Gln	Arg
Gln	Ala 1325		Arg	Thr	Pro	Cys 1330		Arg	Pro	Pro	Leu 1335	Ala	Gly	Thr
Gln	Gly 1340		Val	Thr	Asp	Thr 1345		Ala	Ala	Pro	Leu 1350	Thr	Pro	Ile
Gly	Thr 1355		Leu	Pro	Ser	Ala 1360		Pro	Ser	Gly	Tyr 1365	Cys	Ser	Gln
Asp	Gly 1370		Thr	Gly	Arg	Gln 1375	Pro	Leu	Pro	Pro	Tyr 1380	Thr	Pro	Ala
Met	Met 1385		Arg	Ser	Asn	Gly 1390		Thr	Leu	Thr	Gln 1395	Pro	Pro	Gly
Pro	Arg 1400		Cys	Glu	Gly	Asp 1405		Pro	Glu	His	1410	Val	Glu	Glu
Gly	Thr 1415		Lys	Arg	Val	Ser 1420		Pro	Gln	Trp	Pro 1425	Pro	Pro	Ser
Arg	Ala 1430		Trp	Ala	His	Ala 1435		Arg	Glu	Asp	Ser 1440	Leu	Pro	Glu
Glu	Ser 1445		Ala	Pro	Asp	Phe 1450		Asn	Leu	Lys	His 1455	Tyr	Gln	Lys

- Gln Gln Ser Leu Pro Ser Leu Cys Ser Thr Ser Asp Pro Asp Thr 1460 1465 1470
- Pro Leu Gly Ala Pro Ser Thr Pro Gly Arg Ile Ser Leu Arg Ile 1475 1480 1485
- Ser Glu Ser Val Leu Arg Asp Ser Pro Pro Pro His Glu Asp Tyr 1490 1495 1500
- Glu Asp Glu Val Phe Val Arg Asp Pro His Pro Lys Ala Thr Ser 1505 1510 1515
- Ser Pro Thr Phe Glu Pro Leu Pro Pro Pro Pro Pro Pro Pro Pro 1520 1525 1530
- Ser Gln Glu Thr Pro Val Tyr Ser Met Asp Asp Phe Pro Pro Pro 1535 1540 1545
- Pro Pro His Thr Val Cys Glu Ala Gln Leu Asp Ser Glu Asp Pro 1550 1560
- Glu Gly Pro Arg Pro Ser Phe Asn Lys Leu Ser Lys Val Thr Ile 1565 1570 1575
- Ala Arg Glu Arg His Met Pro Gly Ala Ala His Val Val Gly Ser 1580 1585 1590
- Gln Thr Leu Ala Ser Arg Leu Gln Thr Ser Ile Lys Gly Ser Glu . 1595 1600 1605
- Ala Glu Ser Thr Pro Pro Ser Phe Met Ser Val His Ala Gln Leu 1610 1615 1620
- Ala Gly Ser Leu Gly Gly Gln Pro Ala Pro Ile Gln Thr Gln Ser 1625 1630 1635
- Leu Ser His Asp Pro Val Ser Gly Thr Gln Gly Leu Glu Lys Lys 1640 1650
- Val Ser Pro Asp Pro Gln Lys Ser Ser Glu Asp Ile Arg Thr Glu 1655 1660 1665
- Ala Leu Ala Lys Glu Ile Val His Gln Asp Lys Ser Leu Ala Asp 1670 1680
- Ile Leu Asp Pro Asp Ser Arg Leu Lys Thr Thr Met Asp Leu Met 1685 1690 1695
- Glu Gly Leu Phe Pro Arg Asp Val Asn Leu Leu Lys Glu Asn Ser 1700 1705 1710

- Val Lys Arg Lys Ala Ile Gln Arg Thr Val Ser Ser Ser Gly Cys 1715 1720 1725
- Glu Gly Lys Arg Asn Glu Asp Lys Glu Ala Val Ser Met Leu Val 1730 1735 1740
- Asn Cys Pro Ala Tyr Tyr Ser Val Ser Ala Pro Lys Ala Glu Leu 1745 1750 1755
- Leu Asn Lys Ile Lys Glu Met Pro Ala Glu Val Asn Glu Glu Glu 1760 1765 1770
- Glu Gln Ala Asp Val Asn Glu Lys Lys Ala Glu Leu Ile Gly Ser 1775 1780 1785
- Leu Thr His Lys Leu Glu Thr Leu Gln Glu Ala Lys Gly Ser Leu 1790 1795 1800
- Leu Thr Asp Ile Lys Leu Asn Asn Ala Leu Gly Glu Glu Val Glu 1805 1810 1815
- Ala Leu Ile Ser Glu Leu Cys Lys Pro Asn Glu Phe Asp Lys Tyr 1820 1825 1830
- Arg Met Phe Ile Gly Asp Leu Asp Lys Val Val Asn Leu Leu Leu 1835 1840 1845
- Ser Leu Ser Gly Arg Leu Ala Arg Val Glu Asn Val Leu Ser Gly 1850 1860
- Leu Gly Glu Asp Ala Ser Asn Glu Glu Arg Ser Ser Leu Tyr Glu 1865 1870 1875
- Lys Arg Lys Ile Leu Ala Gly Gln His Glu Asp Ala Arg Glu Leu 1880 1885 1890
- Lys Glu Asn Leu Asp Arg Arg Glu Arg Val Val Leu Gly Ile Leu 1895 1900 1905
- Ala Asn Tyr Leu Ser Glu Glu Gln Leu Gln Asp Tyr Gln His Phe 1910 1915 1920
- Val Lys Met Lys Ser Thr Leu Leu Ile Glu Gln Arg Lys Leu Asp 1925 1930 1935
- Asp Lys Ile Lys Leu Gly Gln Glu Gln Val Lys Cys Leu Leu Glu 1940 1945 1950
- Ser Leu  $\,$  Pro Ser Asp Phe Ile  $\,$  Pro Lys Ala Gly Ala  $\,$  Leu Ala Leu  $\,$  1955  $\,$  1960  $\,$  1965

Pro Pro Asn Leu Thr Ser Glu Pro Ile Pro Ala Gly Gly Cys Thr 1970 1975 Phe Ser Gly Ile Phe Pro Thr Leu Thr Ser Pro Leu 1985 1990 <210> 39 <211> 2529 <212> DNA <213> NM\_004433 ELF3, E74-like factor 3 <220> <221> CDS <222> (203) .. (1315) <223> <400> 39 gagagagaga gagagagaga actagtotog aggtogacgo ggcogcgaat togcggcogc gtcgaccgaa gacgcaggct ctatttagag ccgggtaggg gagcgcacgg ccagatacct cagcgctacc tggcggaact ggatttctct cccgcctgcc ggcctgcctg ccacagccgg actocgocae teeggtagee to atg get gea acc tgt gag att age aac att Met Ala Ala Thr Cys Glu Ile Ser Asn Ile 280 ttt agc aac tac ttc agt geg atg tac agc tcg gag gac tcc acc ctg Phe Ser Asn Tyr Phe Ser Ala Met Tyr Ser Ser Glu Asp Ser Thr Leu 20 gee tet gtt eee eet get gee ace ttt ggg gee gat gae ttg gta etg Ala Ser Val Pro Pro Ala Ala Thr Phe Gly Ala Asp Asp Leu Val Leu acc ctg agc aac ccc cag atg tca ttg gag ggt aca gag aag gcc agc Thr Leu Ser Asn Pro Gln Met Ser Leu Glu Gly Thr Glu Lys Ala Ser tgg ttg ggg gaa cag ccc cag ttc tgg tcg aag acg cag gtt ctg gac Trp Leu Gly Glu Gln Pro Gln Phe Trp Ser Lys Thr Gln Val Leu Asp tgg atc agc tac caa gtg gag aag aac aag tac gac gca agc gcc att 472 Trp Ile Ser Tyr Gln Val Glu Lys Asn Lys Tyr Asp Ala Ser Ala Ile 80 gac ttc tca cga tgt gac atg gat ggc gcc acc ctc tgc aat tgt gcc 520 Asp Phe Ser Arg Cys Asp Met Asp Gly Ala Thr Leu Cys Asn Cys Ala 95 ctt gag gag ctg cgt ctg gtc ttt ggg cct ctg.ggg gac caa ctc cat 568 Leu Glu Glu Leu Arg Leu Val Phe Gly Pro Leu Gly Asp Gln Leu His 115 120 110

					ctc Leu											616
					gag Glu											664
					gac Asp 160											712
					gcc Ala											760
	_				ggc Gly	-		-	Val			-		Thr		808
					cac His											856
					ggc Gly											904
	_	_		_	ccc Pro 240	_			_			-				952
_	_	_	_		gag Glu			-	-				_	_	-	1000
					ggc Gly											1048
			_		ctc Leu					_	-					1096
	_		-		aag Lys		_	_			-		_			1144
					aag Lys 320											1192
					tac Tyr											1240
					tac Tyr											1288
					cag Gln				tgag	ggtt	.gg a	acta	tacc	:c		1335
ggga	ccaa	ac t	cacg	gaco	a ct	cgag	gcct	gca	aacc	ttc	ctgg	ıgagg	ac a	.ggca	ggcca	1395
gato	gccc	ct c	cact	gggg	a at	gctc	ccag	ctg	tgct	gtg	gaga	gaag	ct g	atgt	tttgg	1455

		•				
tgtattgtca	gccatcgtcc	ttggactcgg	agactatggc	ctcgcctccc	caccctcctc	1515
ttggaattac	aagccctggg	gtttgaagct	gactttatag	ctgcaagtgt	atctcctttt	1575
atctggtgcc	tcctcaaacc	cagtctcaga	cactaaatgc	agacaacacc	ttcctcctgc	1635
agacacttgg	actgagccaa	ggaggcttgg	gaggccctag	ggagcaccgt	gatggagagg	1695
acagagcagg	ggctccagca	cttctttctg	gactggcgtt	cacctccctg	ctcagtgctt	1755
gggctccacg	ggcaggggtc	agagcactcc	ctaatttatg	tgctatataa	atatgtcaga	1815
tgtacataga	gatctatttt	ttctaaaaca	ttcccctccc	cactcctctc	ccacagagtg '	1875
ctggactgtt	ccaggccctc	cagtgggctg	atgctgggac	ccttaggatg	gggctcccag	1935
ctcctttctc	ctgtgaatgg	aggcagagac	ctccaataaa	gtgccttctg	ggctttttct	1995
aacctttgtc	ttagctacct	gtgtactgaa	atttgggcct	ttggatcgaa	tatggtcaag	2055
aggttggagg	ggaggaaaat	gaaggtctac	caggatgagg	gtgagggcaa	aggctgacga	2115
agaggggagt	tacagatttc	ctgtagcagg	tgtgggctta	cagacacatg	gactgggctg	2175
ggaggcgagc	aaaggaagca	gctgagactg	ttggagaacg	ttacaagact	tcatgcaaga	2235
aggacatgaa	ctcagaacac	tgaggtcaga	agcatctgct	gtcatgacac	cgctcgagtg	2295
accttgacct	tgaccaagtc	ttgtccttgt	ttaggactga	tttttcctat	taggctaggg	2355
tttggacctg	atgttctcaa	gatgtctaga	attgcatggc	tggccttgtg	gaatagatgg	2415
ttttgcattc	cagccaagtg	tgctgtaaac	tgtatatctg	taatatgaat	cccagctttt	2475
gagtctgaca	aaatcagagt	taggatcttg	taaaggaaaa	aaaaaaaaa	aaaa	2529

<210> 40

<211> 371

<212> PRT

<213> NM\_004433 ELF3, E74-like factor 3

<400> 40

Met Ala Ala Thr Cys Glu Ile Ser Asn Ile Phe Ser Asn Tyr Phe Ser 1  $$10^{\circ}$$ 

Ala Met Tyr Ser Ser Glu Asp Ser Thr Leu Ala Ser Val Pro Pro Ala 20 25 30

Ala Thr Phe Gly Ala Asp Asp Leu Val Leu Thr Leu Ser Asn Pro Gln 35 40 45

Met Ser Leu Glu Gly Thr Glu Lys Ala Ser Trp Leu Gly Glu Gln Pro 50 60

Gln Phe Trp Ser Lys Thr Gln Val Leu Asp Trp Ile Ser Tyr Gln Val 65 70 75 80

Glu Lys Asn Lys Tyr Asp Ala Ser Ala Ile Asp Phe Ser Arg Cys Asp  $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$ 

Met Asp Gly Ala Thr Leu Cys Asn Cys Ala Leu Glu Glu Leu Arg Leu 100 105 110

Val Phe Gly Pro Leu Gly Asp Gln Leu His Ala Gln Leu Arg Asp Leu 115 120 125

Thr Ser Ser Ser Ser Asp Glu Leu Ser Trp Ile Ile Glu Leu Leu Glu 130 135 140

Lys Asp Gly Met Ala Phe Gln Glu Ala Leu Asp Pro Gly Pro Phe Asp 145 150 155 160

Gln Gly Ser Pro Phe Ala Gln Glu Leu Leu Asp Asp Gly Gln Gln Ala 165 170 175

Ser Pro Tyr His Pro Gly Ser Cys Gly Ala Gly Ala Pro Ser Pro Gly 180 185 190

Ser Ser Asp Val Ser Thr Ala Gly Thr Gly Ala Ser Arg Ser Ser His 195 200 205

Ser Ser Asp Ser Gly Gly Ser Asp Val Asp Leu Asp Pro Thr Asp Gly 210 215 220

Lys Leu Phe Pro Ser Asp Gly Phe Arg Asp Cys Lys Lys Gly Asp Pro 225 230 235 240

Lys His Gly Lys Arg Lys Arg Gly Arg Pro Arg Lys Leu Ser Lys Glu 245 250 255

Tyr Trp Asp Cys Leu Glu Gly Lys Lys Ser Lys His Ala Pro Arg Gly 260 265 270

Thr His Leu Trp Glu Phe Ile Arg Asp Ile Leu Ile His Pro Glu Leu 275 280 285

Asn Glu Gly Leu Met Lys Trp Glu Asn Arg His Glu Gly Val Phe Lys 290 295 300

Phe Leu Arg Ser Glu Ala Val Ala Gln Leu Trp Gly Gln Lys Lys 305 310 315 320

Asn Ser Asn Met Thr Tyr Glu Lys Leu Ser Arg Ala Met Arg Tyr Tyr 325 330 335

Tyr Lys Arg Glu Ile Leu Glu Arg Val Asp Gly Arg Arg Leu Val Tyr

113

345 350 340 Lys Phe Gly Lys Asn Ser Ser Gly Trp Lys Glu Glu Glu Val Leu Gln 355 360 Ser Arg Asn 370 <210> 41 <211> 4020 <212> DNA <213> NM\_019027 FLJ20273 <220> <221> CDS <222> (240) .. (1811) <223> <400> 41 gtcaattgcc ctttagtccc aggactaacc ggaagcttct gcaacaggag gacattgaaa 60 ataagatgga acccatccac ataaggattt gcctcaaagg gcactgcaaa aattgaacag 120 aggaatccca aggaagctgc ctgaatttgc ctgtatactc tcgttctgcg acttataaag 180 gaccagacaa atcaaattag tggttttggt ttccgccagc tgtggatgcc tttgacatt 239 atg acc gca gag gat tcc acc gca gcc atg agc agt gac tcg gcc gcc 287 Met Thr Ala Glu Asp Ser Thr Ala Ala Met Ser Ser Asp Ser Ala Ala 10 335 ggg tee teg gee aag gtg eee gag gge gtg geg geg eee aac gag Gly Ser Ser Ala Lys Val Pro Glu Gly Val Ala Gly Ala Pro Asn Glu 383 gca gca ctg ctg gcg ctg atg gag cgc acg ggc tac agc atg gtg caa Ala Ala Leu Leu Ala Leu Met Glu Arg Thr Gly Tyr Ser Met Val Gln 431 gag aac ggg cag cgc aag tac ggc ggc cca ccg ccc ggc tgg gag ggc Glu Asn Gly Gln Arg Lys Tyr Gly Gly Pro Pro Pro Gly Trp Glu Gly 50 55 479 ccg cac ccg cag cgt ggc tgc gag gtc ttc gtg ggc aag atc ccg cgc Pro His Pro Gln Arg Gly Cys Glu Val Phe Val Gly Lys Ile Pro Arg 65 75 527 gac gtg tac gag gac gag ctg gtg ccc gtg ttc gag gcc gtg ggc cgc Asp Val Tyr Glu Asp Glu Leu Val Pro Val Phe Glu Ala Val Gly Arg 90 575 acc tac gag ctg cgc ctc atg atg gac ttt gac ggc aag aac cgc ggc Thr Tyr Glu Leu Arg Leu Met Met Asp Phe Asp Gly Lys Asn Arg Gly

105

110

tac Tyr	gcc Ala	tto Phe 115	val	ato L Met	tao Ty	c tgc c Cys	cac His 120	Lys	g cac s His	gaç Glu	g gco	c aaq a Lys 125	: Ar	g gca	a gtg a Val	623
cgt Arg	gag Glu 130	ı Lev	aad Asr	c aac n Asr	tao Tyi	gag Glu 135	Ile	cgc Arç	Pro	GI7	c cgc Arg 140	j Lei	tei Lei	ggo Gly	e gtg Y Val	671
tgc Cys 145	Cys	ago Ser	gto Val	g gad L Asp	aac Asr 150	ı Cys	cgc Arg	cto Leu	tto Phe	: ato : Ile : 155	: Gl	ggg Gly	ato Ile	e ccc	aag Lys 160	719
atg Met	aag Lys	aag Lys	a cgc	gag Glu 165	Glu	ato Ile	ctg Leu	gaç Glu	gag Glu 170	Ile	gcc Ala	aag Lys	gto Val	acc Thi	gag Glu	767
Gly	gtg Val	ctg Leu	gac Asp 180	Val	ato Ile	gtc Val	tac Tyr	gcc Ala 185	Ser	gcg	gcc Ala	gac Asp	aag Lys 190	Met	, aag : Lys	815
aac Asn	cgc	ggc Gly 195	Phe	gcc Ala	Phe	gtg Val	gag Glu 200	Туг	gag Glu	agc Ser	cac His	cgc Arg 205	Ala	gct Ala	gcc Ala	863
atg Met	gct Ala 210	Arg	cgc	aag Lys	Leu	atg Met 215	cct Pro	Gly	cgc Arg	atc	Gln 220	Leu	tgg Trp	Gly	cac His	911
cag Gln 225	atc Ile	gcc Ala	gtg Val	gac Asp	tgg Trp 230	Ala	gag Gl <sub>u</sub>	cct Pro	gag Glu	atc Ile 235	Asp	gtg Val	gac Asp	gag Glu	gac Asp 240	959
gtg Val	atg Met	gag Glu	acc Thr	gtg Val 245	aag Lys	atc Ile	ctc	tac Tyr	gtg Val 250	cgc Arg	aac Asn	ctc Leu	atg Met	atc Ile 255	Glu	1007
acc Thr	acc Thr	gag Glu	gac Asp 260	acc Thr	atc Ile	aag Lys	aag Lys	agc Ser 265	ttc Phe	ggc Gly	cag Gln	ttc Phe	aac Asn 270	ccc Pro	ggc Gly	1055
tgc Cys	gtg Val	gag Glu 275	cgc Arg	gtc Val	aag Lys	aag Lys	atc Ile 280	cgc Arg	gac Asp	tac Tyr	gcc Ala	ttc Phe 285	gtg Val	cac His	ttc Phe	1103
acc Thr	agc Ser 290	cgc Arg	gag Glu	gat Asp	gcc Ala	gtg Val 295	cat His	gcc Ala	atg Met	aac Asn	aac Asn 300	ctc Leu	aac Asn	GJA	act Thr	1151
gag Glu 305	ctg Leu	gag Glu	ggc Gly	tcg Ser	tgc Cys 310	ctg Leu	gag Glu	gtc Val	acg Thr	ctg Leu 315	gcc Ala	aag Lys	ccc Pro	gtg Val	gac Asp 320	1199
aag Lys	gag Glu	cag Gln	tac Tyr	tcg Ser 325	cgc Arg	tac Tyr	cag Gln	aag Lys	gca Ala 330	gcċ Ala	agg Arg	ggc	ggc Gly	ggc Gly 335	gcg Ala	1247
gct Ala	gag Glu	gca Ala	gcg Ala 340	cag Gln	cag Gln	ccc Pro	agc Ser	tac Tyr 345	gtg Val	tac Tyr	tcc Ser	tgc Cys	gac Asp 350	ccc Pro	tac Tyr	1295
aca Thr	ctg Leu	gcc Ala 355	tac Tyr	tac Tyr	ggc Gly	tac Tyr	ccc Pro 360	tac Tyr	aac Asn	gcg Ala	ctc Leu	att Ile 365	elà aaa	ccc Pro	aac Asn	1343
Arg	gac Asp 370	tac Tyr	ttt Phe	gtg Val	aaa Lys	gta Val 375	gcc Ala	atc Ile	cct Pro	gcc Ala	att Ile 380	'GJÅ aaa	gct Ala	cag Gln	tat Tyr	1391

tcc atg ttt cca gca gct cca gcc cct aaa atg att gaa gat ggc aaa Ser Met Phe Pro Ala Ala Pro Ala Pro Lys Met Ile Glu Asp Gly Lys 385 390 395 400	1439
atc cac aca gtg gag cac atg atc agc ccc att gct gtg cag cca gac Ile His Thr Val Glu His Met Ile Ser Pro Ile Ala Val Gln Pro Asp 405 410 415	1487
cca gcc agt gct gcc gcc gca gcc gcg gcc gca gcc gca gcc Pro Ala Ser Ala	1535
gct gtc att ccc act gtg tcg acg cca cca cct ttc cag ggc cgc cca Ala Val Ile Pro Thr Val Ser Thr Pro Pro Pro Phe Gln Gly Arg Pro 435 440 445	1583
ata act cca gta tac acg gtg gct cca aac gtt cag aga att cct act Ile Thr Pro Val Tyr Thr Val Ala Pro Asn Val Gln Arg Ile Pro Thr 450 460	1631
gcc ggg atc tac ggg gcc agt tac gtg cca ttt gct gct cca gct aca Ala Gly Ile Tyr Gly Ala Ser Tyr Val Pro Phe Ala Ala Pro Ala Thr 465 470 475 480	1679
gcc acg atc gcc aca cta cag aag aac gcg gca gcc gcg gcc gtg Ala Thr Ile Ala Thr Leu Gln Lys Asn Ala Ala Ala Ala Ala Ala Val- 485 490 495	1727
tat gga gga tac gca ggc tac ata cct cag gcc ttc cct gct gcc Tyr Gly Gly Tyr Ala Gly Tyr Ile Pro Gln Ala Phe Pro Ala Ala Ala 500 505 510	1775
att cag gtc ccc atc ccc gac gtc tac cag aca tac tgaggctggt Ile Gln Val Pro Ile Pro Asp Val Tyr Gln Thr Tyr 515 520	1821
gaccagcacg aagacagacc acacaaacac cactgaagga acgcttgact atttatgaag	1881
aaggaacatg ttggattcac acatgcaacc tgaaagtgaa gaatgttagc agatttattt	1941
ctgaattatt ttatatacat gaagttttca ctagttttt aagactattt tcaacttagc	2001
atgcctacgt tcatacattt ccaaaagact tgcaatggtt cgtgccttca ttccatcttt	2061
taaaaatttg tatgctgtac tacatttgta tagaggtttt tgttgttgtt tttttaagga	2121
tatattttca gtatgaaggt tattttctta acttctgcac tccagagatt tctattttgt	2181
agtacettea ataatatate aactatatat taaaaaagea eaettgagga getagggaae	2241
tattttgaaa aatatataca atatttaaag atacaaacag tagtgcttaa aaatactaca	2301
taaagcatta ttttaaaggt tatactggaa agtgcaattt taaaatgagt aaaacctctg	2361
tatttctgct ggcattaagg gttgatggtg ttaccatgta tcatcatggc ggtactattt	2421
tttaaaagaa attaaacact ggatctctcc ttaagccaac attgaaaaga cttgccgcac	2481
ttctgagtcc aaacactgga aagctctcct tgccaccgtt agccggggct cattctccat	2541
gtgeettage ettaaacatg ecceeactee cacatetete accetgteee etceteecca	2601
gatteccaat eccacegeaa tgtttggeaa geetaggaet gataagtage tetgatagag	2661
gagetggtgg ettttataet tetteetggg tttttgttgg ggtttgttgt ttegttgttt	2721
tttgttttt ttttgtttgg ttggggaagt attgtcttct acgtgtgcta ttttcagtag	2781

cagagtaago	acaaggtttt	aatcgagttg	cataagacac	ctttgcatag	ctatttaatt	2841
gcccaatgta	aaactttaat	gccatttcta	atgcttttat	tcatttttga	agtatgagtt	2901
tgtagggaca	aagaatgtat	gttatcgtag	acaagacccc	cagagactct	tttcagcaga	2961
aagttatgct	tctagttgcc	ttaccatgtt	tcttgcaaaa	ctgtccatgg	tcctcaaggg	3021
tgttggaaac	attatgttta	ttaaatgggc	ctctcttcct	ttgctgtgca	cttgatgggt	3081
gaactggatt	ggggtgtgca	catccaggag	gaggaggaga	gacctgtaga	agtttaaaga	3141
tagtttgtaa	atatcttcta	atgcttgttt	ttagtccttt	tatgttggag	aagttcatgg	3201
tatgtagttt	aatgcaaaat	gaaaccattt	tatttcaatg	ttattaaaaa	ggtttgtttt	3261
attaggaagt	taatgtattg	ttgcagtgtt	ttgtgcctgt	ttaaaggctt	ttgtttagca	3321
gagtgaatgt	aaaatacagt	aaaatgttaa	gattgtcatc	tactttttaa	aaaaaaatat	3381
caacttggaa	ttgtttttta	aaggctcaat	caaggaagtg	aggtgtgcaa	taaggtagca	3441
agtaaaacgc	agttgcgttt	ttatgtcatg	ttagagatco	atacaatttt	ccactcacgg	3501
gattittgtt	gatggctgaa	ttcttgtgga	ttcataagag	gatcatgccc	ttagcaagta	3561
cttttgtttt	gttttaaatt	aagagattcc	caaatgcctt	tttccccctc	atcttgaaat	3621
gagatgagtt	tttatgtgta	agcaatattt	atttaactat	tctataaaat	tattgagtgc	3681
ctactgaggc	ctttaagcac	cgctaacatt	cctttccatc	attcttttga	atgacataaa	3741
ataattgtgc	aatgttcctg	atgatgtacc	ccacagctgc	attcaaactc	aaatctgtgg	3801
gaatgagtga	ctcgaccaaa	tgtaattcgg	atcagatcct	catcccctga	ctgtgtgaaa	3861
aagtactct	ccttctagtg	aaggattgtc	acagagtttc	actggatgaa	actatgaccc	3921
gtattctta	ctgtatttta	catgtgcctg	taaattattt	tgccgaaata	agaagaagaa	3981
gaggaagaaa	gaacagtaga	aaaaaaaat	aaaaaaaa			4020

<210> 42

<211> 524

<212> PRT

<213> NM\_019027 FLJ20273

<400> 42

Met Thr Ala Glu Asp Ser Thr Ala Ala Met Ser Ser Asp Ser Ala Ala 1 5 10 15

Gly Ser Ser Ala Lys Val Pro Glu Gly Val Ala Gly Ala Pro Asn Glu 20 25 30

Ala Ala Leu Leu Ala Leu Met Glu Arg Thr Gly Tyr Ser Met Val Gln 35 40 45

Glu Asn Gly Gln Arg Lys Tyr Gly Gly Pro Pro Pro Gly Trp Glu Gly

50 55 60

Pro His Pro Gln Arg Gly Cys Glu Val Phe Val Gly Lys Ile Pro Arg 65 70 75 80

Asp Val Tyr Glu Asp Glu Leu Val Pro Val Phe Glu Ala Val Gly Arg 85 90 95

Thr Tyr Glu Leu Arg Leu Met Met Asp Phe Asp Gly Lys Asn Arg Gly
100 105 110

Tyr Ala Phe Val Met Tyr Cys His Lys His Glu Ala Lys Arg Ala Val 115 120 125

Arg Glu Leu Asn Asn Tyr Glu Ile Arg Pro Gly Arg Leu Leu Gly Val 130 135 140

Cys Cys Ser Val Asp Asn Cys Arg Leu Phe Ile Gly Gly Ile Pro Lys 145 150 155 160

Met Lys Lys Arg Glu Glu Ile Leu Glu Glu Ile Ala Lys Val Thr Glu 165 : 170 175

Gly Val Leu Asp Val Ile Val Tyr Ala Ser Ala Ala Asp Lys Met Lys 180 185 190

Asn Arg Gly Phe Ala Phe Val Glu Tyr Glu Ser His Arg Ala Ala Ala 195 200 205

Met Ala Arg Arg Lys Leu Met Pro Gly Arg Ile Gln Leu Trp Gly His 210 215 220

Gln Ile Ala Val Asp Trp Ala Glu Pro Glu Ile Asp Val Asp Glu Asp 225 230 235 240

Val Met Glu Thr Val Lys Ile Leu Tyr Val Arg Asn Leu Met Ile Glu 245 250 255

Thr Thr Glu Asp Thr Ile Lys Lys Ser Phe Gly Gln Phe Asn Pro Gly 260 265 270

Cys Val Glu Arg Val Lys Lys Ile Arg Asp Tyr Ala Phe Val His Phe 275 280 285

Thr Ser Arg Glu Asp Ala Val His Ala Met Asn Asn Leu Asn Gly Thr 290 295 300

Glu Leu Glu Gly Ser Cys Leu Glu Val Thr Leu Ala Lys Pro Val Asp 305 310 315 320

Lys Glu Gln Tyr Ser Arg Tyr Gln Lys Ala Ala Arg Gly Gly Ala

325 330 335

Ala Glu Ala Gln Gln Pro Ser Tyr Val Tyr Ser Cys Asp Pro Tyr 340 345 350

Thr Leu Ala Tyr Tyr Gly Tyr Pro Tyr Asn Ala Leu Ile Gly Pro Asn 355 360 365

Arg Asp Tyr Phe Val Lys Val Ala Ile Pro Ala Ile Gly Ala Gln Tyr 370 375 380

Ser Met Phe Pro Ala Ala Pro Ala Pro Lys Met Ile Glu Asp Gly Lys 385 390 395 400

Ile His Thr Val Glu His Met Ile Ser Pro Ile Ala Val Gln Pro Asp 405 410 415

Ala Val Ile Pro Thr Val Ser Thr Pro Pro Pro Phe Gln.Gly Arg Pro 435  $\phantom{-}440\phantom{0}$  445

Ile Thr Pro Val Tyr Thr Val Ala Pro Asn Val Gln Arg Ile Pro Thr 450 455 460

Ala Gly Ile Tyr Gly Ala Ser Tyr Val Pro Phe Ala Ala Pro Ala Thr 465 470 475 480

Tyr Gly Gly Tyr Ala Gly Tyr Ile Pro Gln Ala Phe Pro Ala Ala Ala 500 505 510

Ile Gln Val Pro Ile Pro Asp Val Tyr Gln Thr Tyr 515 520

<210> 43

<211> 1451

<212> DNA

<213> NM\_013952 PAX8

<220>

<221> CDS

<222> (161)..(1354)

<223>

<40		43									•					
ttc	agaa	ıgga	ggag	jagac	ac c	gggc	ccaç	gg go	cacco	tcgc	ggg	caac	cgg	acco	aagcag	60
tga	gggc	ctg	cago	cggc	cg ç	gccag	ggca	ag cg	gcag	gege	ggo	ccgg	Jacc	tacg	ggagga	120
ago	cccg	agc	cctc	ggcg	igg d	tgcg	agco	ga ct	cece	ggcg					tcc Ser 5	175
atc Ile	aga Arg	tct Ser	ggc Gly	cat His	gga Gly	, GJA , aaa	cto Lev	g aac 1 Asn	cag Gln 15	ctg Leu	gga Gly	ggg Gly	gco Ala	ttt Phe 20	gtg Val	223
aat Asn	ggc	aga Arg	cct Pro 25	ctg Leu	Pro	gaa Glu	gtg Val	g gtc . Val 30	cgc Arg	cag Gln	Arg	ato	gta Val 35	gac Asp	ctg Leu	271
gcc Ala	cac His	cag Gln 40	ggt	gta Val	agg Arg	ccc Pro	tgc Cys 45	gac Asp	atc Ile	tct Ser	cgc	cag Gln 50	Leu	cgc Arg	gtc Val	319
agc Ser	cat His 55	ggc Gly	tgc Cys	gtc Val	agc Ser	aag Lys 60	ato	ctt Leu	ggc Gly	agg Arg	tac Tyr 65	tac Tyr	gag Glu	act Thr	ggc Gly	367
agc Ser 70	atc Ile	cgg Arg	cct Pro	gga Gly	gtg Val 75	ata Ile	Gly	Gly	tcc Ser	aag Lys 80	ccc Pro	aag Lys	gtg Val	gcc Ala	acc Thr 85	<b>415</b>
ccc Pro	aag Lys	gtg Val	gtg Val	gag Glu 90	aag Lys	att Ile	G1 y ggg	gac Asp	tac Tyr 95	aaa Lys	cgc Arg	cag Gln	aac Asn	cct Pro 100	acc Thr	463
atg Met	ttt Phe	gcc Ala	tgg Trp 105	gag Glu	atc Ile	cga Arg	gac Asp	cgg Arg 110	Leu	ctg Leu	gct Ala	gag Glu	ggc Gly 115	Val	tgt Cys	511
gac Asp	aat Asn	gac Asp 120	act Thr	gtg Val	ccc Pro	agt Ser	gtc Val 125	agc Ser	tcc Ser	att Ile	aat Asn	aga Arg 130	atc Ile	atc Ile	cgg Arg	559
acc Thr	aaa Lys 135	gtg Val	cag Gln	caa Gln	cca Pro	ttc Phe 140	aac Asn	ctc Leu	cct Pro	atg Met	gac Asp 145	agc Ser	tgc Cys	gtg Val	gcc Ala	607
acc Thr 150	aag Lys	tcc Ser	ctg Leu	agt Ser	ccc Pro 155	gga Gly	cac His	acg Thr	ctg Leu	atc Ile 160	ccc Pro	agc Ser	tca Ser	gct Ala	gta Val 165	655
act Thr	ccc Pro	ccg Pro	gag Glu	tca Ser 170	ccc Pro	cag Gln	tcg Ser	gat Asp	tcc Ser 175	ctg Leu	ggc .Gly	tcc Ser	acc Thr	tac Tyr 180	tcc Ser	703
atc Ile	aat Asn	GJ À âââ	ctc Leu 185	ctg Leu	ggc Gly	atc Ile	gct Ala	cag Gln 190	cct Pro	ggc Gly	agc Ser	gac Asp	aag Lys 195	agg Arg	aaa Lys	751
atg Met	gat Asp	gac Asp 200	agt Ser	gat Asp	cag Gln	gat Asp	agc Ser 205	tgc Cys	cga Arg	cta Leu	agc Ser	att Ile 210	gac Asp	tca Ser	cag Gln	799
Ser						cga Arg 220										847

cag Gln 230	cac His	cac His	ctc Leu	gag Glu	ecg Pro 235	Leu	gag Glu	tgc Cys	Pro	ttt Phe 240	gag Glu	cgg Arg	cag Gln	cac His	tac Tyr 245	895
cca Pro	gag Glu	gcc Ala	tat Tyr	gcc Ala 250	tcc Ser	ccc Pro	agc Ser	cac His	acc Thr 255	aaa Lys	ggc	gag Glu	cag Gln	ggc Gly 260	ctc Leu	943
tac Tyr	ccg Pro	ctg Leu	ccc Pro 265	ttg Leu	ctc Leu	aac Asn	agc Ser	acc Thr 270	ctg Leu	gac Asp	gac Asp	ggg	aag Lys 275	gcc Ala	acc Thr	991
ctg Leu	acc Thr	cct Pro 280	tcc Ser	aac Asn	acg Thr	cca Pro	ctg Leu 285	Gly	cgc Arg	aac Asn	ctc Leu	tcg Ser 290	act Thr	cac His	cag Gln	1039
acc Thr	tac Tyr 295	Pro	gtg Val	gtg Val	gca Ala	gct Ala 300	ccg Pro	ccc Pro	ttt Phe	tgg Trp	atc Ile 305	tgc Cys	agc Ser	aag Lys	tcg Ser	1087
gct Ala 310	ccg Pro	Gly ggg	tcc Ser	cgc Arg	cct Pro 315	tca Ser	atg Met	cct Pro	ttc Phe	ccc Pro 320	atg Met	ctg Leu	cct Pro	ccg Pro	tgt Cys 325	1135
acg Thr	ggc Gly	agt Ser	tca Ser	cgg Arg 330	gcc Ala	agg Arg	ccc Pro	tcc Ser	tct Ser 335	cag Gln	ggc Gly	gag Glu	aga Arg	tgg Trp 340	tgg Trp	1183
ggc Gly	cca Pro	cgc Arg	tgc Cys 345	ccg Pro	gat Asp	acc Thr	cac His	ccc Pro 350	aca Thr	tcc Ser	cca Pro	cca Pro	gcg Ala 355	gac Asp	agg Arg	1231
gca Ala	gct Ala	atg Met 360	cct Pro	cct Pro	ctg Leu	cca Pro	tcg Ser 365	cag Gln	gca Ala	tgg Trp	tgg Trp	cag Gln 370	gaa Glu	gtg Val	aat Asn	1279
Thr	ctg Leu 375	gca Ala	atg Met	cct Pro	atg Met	gcc Ala 380	aca Thr	ccc Pro	cct Pro	act Thr	cct Pro 385	cct Pro	aca Thr	gcg Ala	agg Arg	1327
cct ( Pro ( 390	gly ggg	gct Ala	tcc Ser	cca Pro	act Thr 395	cca Pro	gct Ala	tgc Cys	tgag	ttcc	:cc а	tatt	atta	c		1374
agtt	ccac	at c	aagg	ccga	g tg	cacc	gccc	acc	actg	cca	cggc	cttt	ga c	catc	tgtag	1434
ttgc	catg	gg g	acag	tg			•									1451
<210:	> 4	4														
<211	> 3	98				٠										
<212	> P	RT														
<213>	> N	M_01	3952	PAX	8		-									
										•						

<400> 44

Met Pro His Asn Ser Ile Arg Ser Gly His Gly Gly Leu Asn Gln Leu 1 5 10 10 15

Gly Gly Ala Phe Val Asn Gly Arg Pro Leu Pro Glu Val Val Arg Gln 20 25 30

WO 2004/022778 PCT/AU2003/001166

Arg Ile Val Asp Leu Ala His Gln Gly Val Arg Pro Cys Asp Ile Ser
35 40 45

Arg Gln Leu Arg Val Ser His Gly Cys Val Ser Lys Ile Leu Gly Arg
50 55 60

Tyr Tyr Glu Thr Gly Ser Ile Arg Pro Gly Val Ile Gly Gly Ser Lys 65 70 75 80

Pro Lys Val Ala Thr Pro Lys Val Val Glu Lys Ile Gly Asp Tyr Lys 85 90 95

Arg Gln Asn Pro Thr Met Phe Ala Trp Glu Ile Arg Asp Arg Leu Leu 100 105 110

Ala Glu Gly Val Cys Asp Asn Asp Thr Val Pro Ser Val Ser Ser Ile 115 120 125

Asn Arg Ile Ile Arg Thr Lys Val Gln Gln Pro Phe Asn Leu Pro Met 130 135 140

Asp Ser Cys Val Ala Thr Lys Ser Leu Ser Pro Gly His Thr Leu Ile 145 150 155 160

Pro Ser Ser Ala Val Thr Pro Pro Glu Ser Pro Gln Ser Asp Ser Leu 165 170 175

Gly Ser Thr Tyr Ser Ile Asn Gly Leu Leu Gly Ile Ala Gln Pro Gly 180 185 190

Ser Asp Lys Arg Lys Met Asp Asp Ser Asp Gln Asp Ser Cys Arg Leu 195 200 205

Ser Ile Asp Ser Gln Ser Ser Ser Gly Pro Arg Lys His Leu Arg 210 215 220

Thr Asp Ala Phe Ser Gln His His Leu Glu Pro Leu Glu Cys Pro Phe 225 230 235 240

Glu Arg Gln His Tyr Pro Glu Ala Tyr Ala Ser Pro Ser His Thr Lys 245 250 255

Gly Glu Gln Gly Leu Tyr Pro Leu Pro Leu Leu Asn Ser Thr Leu Asp 260 265 270

Asp Gly Lys Ala Thr Leu Thr Pro Ser Asn Thr Pro Leu Gly Arg Asn 275 280 285

Leu Ser Thr His Gln Thr Tyr Pro Val Val Ala Ala Pro Pro Phe Trp 290 295 300

Ile Cys S 305	er Lys Ser Ala Pro G 310	ly Ser Arg Pro 315	Ser Met Pro Phe Pro 320	
Met Leu P	ro Pro Cys Thr Gly Se 325	er Ser Arg Ala 330	Arg Pro Ser Ser Gln 335	
Gly Glu A	rg Trp Trp Gly Pro An 340	rg Cys Pro Asp 345	Thr His Pro Thr Ser	
Pro Pro A	la Asp Arg Ala Ala Me 55 36	et Pro Pro Leu 60	Pro Ser Gln Ala Trp 365	
Trp Gln Gl 370	lu Val Asn Thr Leu Al 375	la Met Pro Met	Ala Thr Pro Pro Thr 380	
Pro Pro Th	or Ala Arg Pro Gly Al 390	a Ser Pro Thr 395		
<210> 45				
<211> 326	5			
<212> DNA	1	•		
<213> AI3	01558 EST			
<400> 45	gttacattca tttgattc	ag tecettataa	accccacacc tcataaacaa	a 60
			ttctagagtc gttctggttt	
			tcacttggca tgtgtgaaag	
attcaggggg	tctgcagctg tttagtgt	tg atgcagttgg	gtcaaaagag tatcatgtta	240
gtcttctgtg	ggttttaggg agggatta	tg gagcctccct	cccaccccac tggctttctt	300
gtgtcacagc	ctttatttct actccg			326
<210> 46				
<211> 153	4			
<212> DNA				
<213> NM_(	018000 FLJ10116			
	•			
<220>				
<221> CDS				
<222> (267	7)(938)			
<223>				

<400 ttt		16 tog (	gggaa	atta	aa ag	gaggg	ggaaa	a aaa	agco	ccga	agaa	aaci	cca d	egec	caaac	60
aaaa	acqca	aag (	gagag	gagg	ad co	gege	geet	t gca	agcc	tog		gogto	ecc o	ggco	gegge	120
												•			gacgct	180
									٠.						gettee	240
		•													acc	293
gge	geeg	grg (	99999	SCCLI	ay c	.0.00									g Thr	233
gtg Val 10	tgc Cys	tgc Cys	tgc Cys	tgc Cys	cgg Arg 15	tgc Cys	gag Glu	tgc Cys	ttg Leu	gag Glu 20	gag Glu	cgc Arg	gcc Ala	ctg Leu	cct Pro 25	341
			ccc Pro													389
			gtg Val 45													437
gat Asp	gtg Val	tcc Ser 60	cac His	aca Thr	gag Glu	gca Ala	gac Asp 65	gac Asp	gac Asp	aga Arg	acc Thr	ctg Leu 70	tac Tyr	aat Asn	ttg Leu	485
			cgt Arg													533
			gat Asp													581
			aag Lys													629
			ttg Leu 125													67:7
aac Asn	aca Thr	agg Arg 140	aaa Lys	gct Ala	cga Arg	gag Glu	atg Met 145	ttg Leu	tta Leu	aaa Lys	ctg Leu	gct Ala 150	gaa Glu	gaa Glu	acc Thr	725
agt Ser	att Ile 155	ttc Phe	cca Pro	aca Thr	agt Ser	tgg Trp 160	gag Glu	ctc Leu	tca Ser	gag Glu	aga Arg 165	tat Tyr	ctc Leu	ttt Phe	gtt Val	773
			ctc Leu													821
gct Ala	cgt Arg	cga Arg	act Thr	tac Tyr 190	ccc Pro	aag Lys	aag Lys	cct Pro	ggg Gly 195	gtt Val	cca Pro	tgc Cys	ctg Leu	gca Ala 200	gat Asp	869
ggc Gly	cag Gln	aaa Lys	gaa Glu 205	ctg Leu	cac His	ctg Leu	tgg Trp	ggg Gly 210	gac Asp	ctc Leu	tca Ser	tgc Cys	aga Arg 215	ctt Leu	gca Ala	917
cat	atg	cag	gga	gta	ttg	cac	tgaa	agato	ett t	gato	gaco	ct to	ttci	tatta	3	968

His Met Gln Gly Val Leu His 220

agaagataat	tttcaaaagg	gagcaatgct	gtgaatgcag	cttgcttctc	tctacagatt	102
gagaagtcca	gcttcaaaag	ttacttgcca	cttaagcaag	gaacttgtca	agagatcatg	108
gttcatgtta	ctgaaaagac	tttaaggatt	tgtaaggtta	atccatagat	tgctgagaac	1148
aatggaaata	tttttatttt	tacagatttt	gcacttctga	attcaggtta	aaaactaact	1208
tgtatttagt	ctgcttagag	gactgtgact	tgaaaatttt	tatataccaa	tgagcttttt	1268
ggtagcgtcc	acaatgttta	aaatatttca	taggcgagat	ccgtgttctc	catttattaa	1328
tgcattgtag	accaatttaa	ctgctgtgtt	tcaggaaaat	tcttcctagt	ttaataagca	1388
agctaaaagt	tttattttt	atatttagtg	cttaatcttt	gcctcatgtt	atgtaaaatt	1448
agcctgcaga	tattttctct	caattctgta	gactctcgca	agataaacat	tcaaacagtg	1508
aacaaacaa	taaaataaat	aaacct				1534

<210> 47

<211> 224

<212> PRT

<213> NM\_018000 FLJ10116

<400> 47

Met Gly Leu Arg Asp Trp Leu Arg Thr Val Cys Cys Cys Cys Arg Cys 1 5 10 15

Glu Cys Leu Glu Glu Arg Ala Leu Pro Glu Lys Glu Pro Leu Val Ser 20 25 30

Asp Asn Asn Pro Tyr Ser Ser Phe Gly Ala Thr Leu Val Arg Asp Asp 35 40 45

Glu Lys Asn Leu Trp Ser Met Pro His Asp Val Ser His Thr Glu Ala 50 55 60

Asp Asp Asp Arg Thr Leu Tyr Asn Leu Ile Val Ile Arg Asn Gln Gln 65 70 75 80

Ala Lys Asp Ser Glu Glu Trp Gln Lys Leu Asn Tyr Asp Ile His Thr 85 90 95

Leu Arg Gln Val Arg Arg Glu Val Arg Asn Arg Trp Lys Cys Ile Leu 100 105 110

Glu Asp Leu Gly Phe Gln Lys Glu Ala Asp Ser Leu Leu Ser Val Thr 115 120 125 WO 2004/022778 PCT/AU2003/001166

125

						•										
Lys 1	Leu 130	Ser	Thr	Ile	Ser	Asp 135		Lys	Asn	Thr	Arg 140		Ala	Arg	Glu	
Met 1 145	Leu	Leu	Lys	Leu	Ala 150		Glu	Thr	Ser	Ile 155		Pro	Thr	Ser	Trp 160	
Glu 1	Leu	Ser	Glu	Arg 165		Leu	Phe	Val	Val 170		Arg	Leu	Ile	Ala 175		
Asp A	Ala	Ala	Glu 180	Glu	Phe	Phe	Lys	Leu 185		Arg	Arg	Thr	Tyr 190		Lys	
Lys I	Pro	Gly 195	Val	Pro	Суз	Leu	Ala 200		Gly	Gln	Lys	Glu 205	Leu	His	Leu	
Trp 6	Sly 210	Asp	Leu	Ser	Cys	Arg 215	Leu	Ala	His	Met	Gln 220	Gly	Val	Leu	His	
<210>	- 4	8														
<211>	. 2	385														
<212>	D	NA														
<213>	· N	M_1	4472	4 FL	J305:	32			•							
<220>									,		÷					
<221>	С	DS														
<222>	. (	71).	. (14	141)					٠							
<223>									•	•					-	
<400> gatac	-		caga	igtco	et ge	gcag	regte	g ego	gete	ttc	ctgg	cggc	etq (	cqcad	gatat	q 60
aaaat		aa a	itg t	ca a	at ç		rga a	iga t	cc a	igg a	at c sn A	gg g	rac a	agg o	ege	109
tac garage Tyr A	sp (	gag Glu	gtc Val	cca Pro	agc Ser	gac Asp 20	ctg Leu	ccc Pro	tat Tyr	caa Gln	gat Asp 25	acc Thr	acc Thr	ata Ile	aga Arg	157
acc ca Thr H: 30	ac d is l	cca Pro	att Ile	Leu	cat His 35	gac Asp	agt Ser	gag Glu	cgg Arg	gca Ala 40	gtg Val	agc Ser	gct Ala	gat Asp	ccc Pro 45	205
ttg co Leu Pi	ca d ro I	cca Pro	Pro	cct Pro 50	ctc Leu	cca Pro	tta Leu	Gln	cca Pro 55	cca Pro	ttc Phe	ggc Gly	cca Pro	gac Asp 60	ttc Phe	253
tac to Tyr Se	ca a er S	er.	gac Asp 65	aca Thr	gaa Glu	gaa Glu	cca Pro	gct Ala 70	ata Ile	gcg Ala	cca Pro	Asp	ctc Leu 75	aaa Lys	cca Pro	301
gta aç	gg c	gc	ttt	gtc	cct	gac	tcc	tgg .	aag	aac	ttt	ttc	aga	ggg	aag	349

Val	. Arc	80 80	J Ph∈	e Val	. Pro	Asp	Ser 85	Trp	Lys	Asr	Phe	Phe 90	Arg	g Gly	Lys	
aaa Lys	aag Lys 95	gac Asp	Pro	gaa Glu	tgg Trp	gat Asp 100	Lys	ccg Pro	gtg Val	ser	gat Asp 105	Ile	agg Arg	tac Tyr	atc Ile	397
tec Ser 110	Asp	gga Gly	gtg Val	gag Glu	tgt Cys 115	Ser	Pro	cca Pro	gcc Ala	Ser 120	Pro	gca	aga Arg	cca Pro	aac Asn 125	445
cac His	cgt Arg	tcg Ser	Pro	cto Leu 130	Asn	tcc Ser	tgc Cys	aaa Lys	gat Asp 135	Pro	tac Tyr	gga Gly	ggg	tca Ser 140		493
				Ser				gct Ala 150	Asp					Arg		541
ccc Pro	tat Tyr	gga Gly 160	Ser	cta Leu	gac Asp	cga Arg	cac His 165	aca Thr	caa Gln	aca Thr	gtt Val	cga Arg 170	Thr	tac Tyr	agt Ser	589
		Val						aga Arg								637
gca Ala 190	Gly	ctg Leu	ctg Leu	aga Arg	ata Ile 195	ctg Leu	ggt Gly	gtg Val	gtg Val	gag Glu 200	Leu	ctt Leu	ttg Leu	GJ A GGG	gcc Ala 205	685
ggt Gly	gtc Val	ttt Phe	gct Ala	tgt Cys 210	gtc Val	aca Thr	gct Ala	tac Tyr	att Ile 215	cac His	aag Lys	gac Asp	agt Ser	gag Glu 220	tgg Trp	733
tac Tyr	aac Asn	ttg Leu	ttt Phe 225	Gly	tat Tyr	tca Ser	caa Gln	ccg Pro 230	tat Tyr	GJ A GGC	atg Met	gga Gly	ggc Gly 235	gtt Val	ggt Gly	781
						Gly		tat Tyr								829
cct Pro	ttt Phe 255	gta Val	ctc Leu	gtg Val	gtt Val	gct Ala 260	gga Gly	tta Leu	gct Ala	tgg Trp	atc Ile 265	·acc Thr	acc Thr	att Ile	att Ile	877
att Ile 270	ctg Leu	gtt Val	ctt Leu	ggc Gly	atg Met 275	tcc Ser	atg Met	tat Tyr	tac Tyr	cgg Arg 280	Thr	att Ile	ctt Leu	ctg Leu	gac Asp 285	925
tct Ser	aat Asn	tgg Trp	tgg Trp	ccc Pro 290	cta Leu	act Thr	gaa Glu	ttt Phe	gga Gly 295	att Ile	aac Asn	gtt Val	gcc Ala	ttg Leu 300	ttt Phe	973
att Ile	ttg Leu	tat Tyr	atg Met 305	gcc Ala	gca Ala	gcc Ala	ata Ile	gtc Val 310	tat Tyr	gtg Val	aat Asn	gat Asp	acc Thr 315	aac Asn	cga Arg	1021
ggt Gly	ggc Gly	ctc Leu 320	tgc Cys	tac Tyr	tat Tyr	Pro	tta Leu 325	ttt Phe	aat Asn	aca Thr	cca Pro	gtg Val 330	aat Asn	gca Ala	gtg Val	1069
ttc Phe	tgc Cys 335	cgg Arg	gta Val	gaa Glu	Gly	gga Gly 340	cag Gln	ata Ile	gct Ala	gca Ala	atg Met 345	atc Ile	ttc Phe	ctg Leu	ttt Phe	1117
gtc	acc	atg	ata	gtt	tat	ctc	att	agt	gcť	ttg	gtt	tgc	cta	aag	tta	1165

Val Thr Met Ile Val Tyr Leu Ile Ser Ala Leu Val Cys Leu Lys Leu 350 365	
tgg agg cat gag gca gct cgg aga cat aga gaa tat atg gaa caa cag Trp Arg His Glu Ala Ala Arg Arg His Arg Glu Tyr Met Glu Gln Gln 370 375 380	1213
gag ata aat gag cca tca ttg tca tcg aaa agg aaa atg tgt gaa atg Glu Ile Asn Glu Pro Ser Leu Ser Ser Lys Arg Lys Met Cys Glu Met 385 390 395	1261
gcc acc agt ggt gac aga caa aga gac tca gaa gtt aat ttc aag gaa Ala Thr Ser Gly Asp Arg Gln Arg Asp Ser Glu Val Asn Phe Lys Glu 400 405 410	1309
ctg aga aca gca aaa atg aaa cct gaa cta ctg agt gga cac atc ccc Leu Arg Thr Ala Lys Met Lys Pro Glu Leu Leu Ser Gly His Ile Pro 415 420 425	1357
cca cgc cca gct aat ttt ttt gta ttt tta gta gag atg ggg ttt cac Pro Arg Pro Ala Asn Phe Phe Val Phe Leu Val Glu Met Gly Phe His 435 440 445	. 1405
cgt gtt agc cag gat gat ctc gat ctc ctg acc tca tgatccaccc Arg Val Ser Gln Asp Asp Leu Asp Leu Thr Ser 450 455	1451
gcctcagcct cccaaagtgt tgggattaca ggcgtgagtc accgcgccca gctggtattg	1511
cttttctatt ccctttggac atacatgcta cagtcccaca atgtagcatt tccttggaaa	1571
ctcccttttt ttttttttt gagatggagt ttcgctcttg ttgcccaggc tggagtacag	1631
tggtatgatc ttggctcact gcagcctctg cctcctgggt tcaagcgatt ctcctgcctc	1691
tgcctcccaa gtagctggga ttacaggcac ccaccaccat gcccagctaa ttttttgtat	1751
ttttagtaga gacaggattt cactatgttg gccaggttgg tctcaagctc ctgacctcag	1811
atgatctacc agccteggee ttetgaagtg etgggattea ggtgtgagee aetgtgeeea	1871
gcagggatgc ttcatctttc taagaattat cttggctttg gactttattc ataaatgttt	1931
tatttctgtt agtatgaaca atagactgcc ttaacaaagt ttttttttaa acaaaatcgt	1991
tottgttgga ttttattcag cagcatctat catgtagata aattcccagg tgtagcatta	2051
cagettetga etaatatage tgecatteag acaattaatg tteaaagagt tttetaaagt	2111
gataaaacca aagaaaagca tgtggaaaag cagaagctta gaaagttgtg gtcactgaat	2171
gcactccctg gtttttattt gtcagtgaaa tctttatgca ttcattgtta atattttaat	2231
tccatggctt tgtaggctgt gctgtgtctg aaggggtaac acctagggaa acatgaggcc	2291
ccttatggga cccccaaat ggaacaactt cactttctct tttatgtatt gagccctgtg	2351
ttaacatttc acttaagaag agcaccagtg cttt	2385

<210> 49

<211> 457

<212> PRT

<213> NM\_144724 FLJ30532

<400> 49

Val Pro Ser Asp Leu Pro Tyr Gln Asp Thr Thr Ile Arg Thr His Pro 20 25 30

Ile Leu His Asp Ser Glu Arg Ala Val Ser Ala Asp Pro Leu Pro Pro 35 40 45

Pro Pro Leu Pro Leu Gln Pro Pro Phe Gly Pro Asp Phe Tyr Ser Ser 50 55 60

Asp Thr Glu Glu Pro Ala Ile Ala Pro Asp Leu Lys Pro Val Arg Arg 65 70 . 75 80

Phe Val Pro Asp Ser Trp Lys Asn Phe Phe Arg Gly Lys Lys Asp 85 90 95

Pro Glu Trp Asp Lys Pro Val Ser Asp Ile Arg Tyr Ile Ser Asp Gly
100 105 110

Val Glu Cys Ser Pro Pro Ala Ser Pro Ala Arg Pro Asn His Arg Ser 115 120 125

Pro Leu Asn Ser Cys Lys Asp Pro Tyr Gly Gly Ser Glu Gly Thr Phe 130 135 140

Ser Ser Arg Lys Glu Ala Asp Ala Val Phe Pro Arg Asp Pro Tyr Gly 145 150 150 165

Ser Leu Asp Arg His Thr Gln Thr Val Arg Thr Tyr Ser Glu Lys Val 165 170 175

Glu Glu Tyr Asn Leu Arg Tyr Ser Tyr Met Lys Ser Trp Ala Gly Leu 180 185 190

Leu Arg Ile Leu Gly Val Val Glu Leu Leu Gly Ala Gly Val Phe 195 200 205

Ala Cys Val Thr Ala Tyr Ile His Lys Asp Ser Glu Trp Tyr Asn Leu 210 215 220

Phe Gly Tyr Ser Gln Pro Tyr Gly Met Gly Gly Val Gly Gly Leu Gly 225 230 235 240

Ser Met Tyr Gly Gly Tyr Tyr Tyr Thr Gly Pro Lys Thr Pro Phe Val 245 250 255

Leu Val Val Ala Gly Leu Ala Trp Ile Thr Thr Ile Ile Ile Leu Val 260 265 270

Leu Gly Met Ser Met Tyr Tyr Arg Thr Ile Leu Leu Asp Ser Asn Trp 275 280 285

Trp Pro Leu Thr Glu Phe Gly Ile Asn Val Ala Leu Phe Ile Leu Tyr 290 295 300

Met Ala Ala Ala Ile Val Tyr Val Asn Asp Thr Asn Arg Gly Gly Leu 305 310 315 320

Cys Tyr Tyr Pro Leu Phe Asn Thr Pro Val Asn Ala Val Phe Cys Arg 325 330 335

Val Glu Gly Gly Gln Ile Ala Ala Met Ile Phe Leu Phe Val Thr Met 340 345 350

Ile Val Tyr Leu Ile Ser Ala Leu Val Cys Leu Lys Leu Trp Arg His 355 360 365

Glu Ala Ala Arg Arg His Arg Glu Tyr Met Glu Gln Gln Glu Ile Asn 370 375 380

Glu Pro Ser Leu Ser Ser Lys Arg Lys Met Cys Glu Met Ala Thr Ser 385 390 395 400

Gly Asp Arg Gln Arg Asp Ser Glu Val Asn Phe Lys Glu Leu Arg Thr 405 410 415

Ala Lys Met Lys Pro Glu Leu Leu Ser Gly His Ile Pro Pro Arg Pro 420 425 430

Ala Asn Phe Phe Val Phe Leu Val Glu Met Gly Phe His Arg Val Ser 435 440 445

Gln Asp Asp Leu Asp Leu Leu Thr Ser 450 455

<210> 50

<211> 2280

<212> DNA

<213> NM\_006424 SLC34A2

<220>

<221> CDS

130

<222> (64)..(2133)

<223>

<4(	00>	50								•						
cgg	geca	aggt	ttco	caggo	etc (	gccg	ccgo	c to	cato	ccaç	g cad	ctg	egga	ggga	igeget	g 60
aco	Met	g gct : Ala	ccc Pro	tgg Trp	p cct Pro 5	gaa Glu	tto Lei	gga Gly	gat Asp	gcc Ala 10	caç Glr	cco Pro	aac Asr	ccc Pro	gat Asp 15	108
aag Lys	tac Tyr	cto Lev	gaa Glu	ggg Gly 20	gco Ala	gca Ala	ggt Gly	cag Gln	cag Gln 25	Pro	act Thr	gco Ala	cct Pro	gat Asp 30	aaa Lys	156
ago Ser	aaa Lys	gag Glu	acc Thr 35	aac Asn	aaa Lys	aca Thr	gat Asp	aac Asn 40	act Thr	gag Glu	gca Ala	cct Pro	gta Val 45	acc Thr	Lys	204
att Ile	gaa Glu	ctt Lev 50	ctg Leu	r ccg	tcc Ser	tac Tyr	Ser 55	acg Thr	gct Ala	aça Thr	. ctg	ata Ile 60	gat Asp	gag Glu	ccc Pro	252
act Thr	gag Glu 65	gtg Val	gat Asp	gac Asp	Pro	tgg Trp 70	aac Asn	cta Leu	ccc Pro	act Thr	ctt Leu 75	cag Gln	gac Asp	tcg Ser	GJ Å GGG	300
ato Ile 80	aag Lys	tgg Trp	tca Ser	gag Glu	aga Arg 85	gac Asp	acc Thr	aaa Lys	Gly	aag Lys .90	att Ile	ctc	tgt Cys	ttc Phe	ttc Phe 95	348
caa Gln	ggg	att Ile	G1 y	aga Arg 100	ttg Leu	att Ile	tta Leu	ctt Leu	ctc Leu 105	gga Gly	ttt Phe	ctc	tac Tyr	ttt Phe 110	ttc Phe	396
gtg Val	tgc Cys	tcc Ser	ctg Leu 115	Asp	att Ile	ctt Leu	agt Ser	agc Ser 120	gcc Ala	ttc Phe	cag Gln	ctg Leu	gtt Val 125	Gly	gga Gly	444
aaa Lys	atg Met	gca Ala 130	gga Gly	cag Gln	ttc Phe	ttc Phe	agc Ser 135	aac Asn	agc Ser	tct Ser	att Ile	atg Met 140	tcc Ser	aac Asn	cct Pro	492
ttg Leu	ttg Leu 145	ggg Gly	ctg Leu	gtg Val	atc Ile	ggg Gly 150	gtg Val	ctg Leu	gtg Val	acc Thr	gtc Val 155	ttg Leu	gtg Val	cag Gln	agc Ser	540
tcc Ser 160	agc Ser	acc Thr	tca Ser	acg Thr	tcc Ser 165	atc Ile	gtt Val	gtc Val	agc Ser	atg Met 170	gtg Val	tcc Ser	tct Ser	tca Ser	ttg Leu 175	588
ctc Leu	act Thr	gtt Val	cgg Arg	gct Ala 180	gcc Ala	atc Ile	ccc Pro	att Ile	atc Ile 185	atg Met	GJA aaa	gcc Ala	aac Asn	att Ile 190	gga Gly	636
acg Thr	tca Ser	atc Ile	acc Thr 195	aac Asn	act Thr	att Ile	gtt Val	gcg Ala 200	ctc Leu	atg Met	cag Gln	gtg Val	gga Gly 205	gat Asp	cgg Arg	684
agt Ser	gag Glu	ttc Phe 210	aga Arg	aga Arg	gct Ala	ttt Phe	gca Ala 215	gga Gly	gcc Ala	act Thr	gtc Val	cat His 220	gac Asp	ttc Phe	ttc Phe	732
aac Asn	tgg Trp 225	ctg Leu	tcc Ser	gtg Val	ttg Leu	gtg Val 230	ctc Leu	ttg Leu	ccc Pro	gtg Val	gag Glu 235	gtg Val	gcc Ala	acc Thr	cat His	78,0

											•					
tac Tyr 240	ctc Leu	gag Glu	atc Ile	ata Ile	acc Thr 245	cag Gln	ctt Leu	ata Ile	gtg Val	gag Glu 250	Ser	ttc Phe	cac His	ttc Phe	aag Lys 255	828
aat Asn	gga Gly	gaa Glu	gat Asp	gcc Ala 260	cca Pro	gat Asp	ctt Leu	ctg Leu	aaa Lys 265	gtc Val	atc Ile	act Thr	aag Lys	ccc Pro 270	ttc Phe	876
aca Thr	aag Lys	ctc Leu	att Ile 275	gtc Val	cag Gln	ctg Leu	gat Asp	aaa Lys 280	aaa Lys	gtt Val	atc Ile	agc Ser	caa Gln 285	att Ile	gca Ala	924
atg Met	aac Asn	gat Asp 290	gaa Glu	aaa Lys	gcg Ala	aaa Lys	aac Asn 295	aag Lys	agt Ser	ctt Leu	gtc Val	aag Lys 300	att Ile	tgg Trp	tgc Cys	972
aaa Lys	act Thr 305	ttt Phe	acc Thr	aac Asn	aag Lys	acc Thr 310	cag Gln	att Ile	aac Asn	gtc Val	act Thr 315	gtt Val	ccc Pro	tcg Ser	act Thr	1020
gct Ala 320	aac Asn	tgc Cys	acc Thr	tcc Ser	cct Pro 325	tcc Ser	ctc Leu	tgt Cys	Trp	acg Thr 330	gat Asp	ggc Gly	atc Ile	caa Gln	aac Asn 335	1068
tgg Trp	acc Thr	atg Met	aag Lys	aat Asn 340	gtg Val	acc Thr	tac Tyr	aag Lys	gag Glu 345	aac Asn	atc Ile	gcc Ala	aaa Lys	tgc Cys 350	cag Gln	1116
cat His	atc Ile	ttt Phe	gtg Val 355	aat Asn	ttc Phe	cac His	ctc Leu	ccg Pro 360	gat Asp	ctt Leu	gct Ala	gtg Val	ggc Gly 365	acc Thr	atc Ile	1164
ttg Leu	ctc Leu	ata Ile 370	ctc Leu	tcc Ser	ctg Leu	ctg Leu	gtc Val 375	ctc Leu	tgt Cys	ggt Gly	tgc Cys	ctg Leu 380	atc Ile	atg Met	att Ile	1212
gtc Val	aag Lys 385	atc Ile	ctg Leu	ggc	tct Ser	gtg Val 390	ctc Leu	aag Lys	Gly	cag Gln	gtc Val 395	gcc Ala	act Thr	gtc Val	atc Ile	1260
aag Lys 400	aag Lys	acc Thr	atc Ile	aac Asn	act Thr 405	gat Asp	ttc Phe	ccc Pro	ttt ·Phe	ccc Pro 410	ttt Phe	gca Ala	tgg Trp	ttg Leu	act Thr 415	1308
ggc	tac Tyr	ctg Leu	gcc Ala	atc Ile 420	ctc Leu	gtc Val	Gly ggg	gca Ala	ggc Gly 425	atg Met	acc Thr	ttc Phe	atc Ile	gta Val 430	cag Gln	1356
agc Ser	agc Ser	tct Ser	gtg Val 435	ttc Phe	acg Thr	tcg Ser	gcc Ala	ttg Leu 440	acc Thr	ccc Pro	ctg Leu	att Ile	gga Gly 445	Ile	ggc Gly	1404
gtg Val	ata Ile	acc Thr 450	Ile	gag Glu	agg Arg	gct Ala	tat Tyr 455	Pro	ctc	acg Thr	ctg Leu	ggc Gly 460	Ser	aac Asn	atc Ile	1452
Gly	acc Thr 465	Thr	acc	acc Thr	gcc Ala	atc Ile 470	Leu	gcc Ala	gcc Ala	tta Leu	gcc Ala 475	·Ser	cct Pro	ggc	aat Asn	1500
gca Ala 480	Leu	agg Arg	agt Ser	tca Ser	ctc Leu 485	Gln	atc Ile	gcc Ala	ctg Leu	tgc Cys 490	His	ttt Phe	ttc Phe	ttc Phe	aac Asn 495	1548
atc Ile	tcc Ser	ggc	atc Ile	ttg Leu 500	Leu	tgg Trp	tac Tyr	ccg Pro	atc Ile 505	Pro	ttc Phe	act Thr	cgc	ctg Leu 510	ccc Pro	1596

atc Ile	cgc Arg	atg Met	gcc Ala 515	Lys	Gly ggg	ctg Leu	Gly	aac Asn 520	atc Ile	tct Ser	gcc Ala	aag Lys	tat Tyr 525	cgc Arg	tgg Trp	1644
												atc Ile 540				1692
												gtt Val				1740
											Cys	ctc Leu				1788
												cag Gln				1836
												gat Asp				1884
tcc Ser	aag Lys	ttc Phe 610	acc Thr	ggc Gly	tgc Cys	ttc Phe	cag Gln 615	atg Met	cgc Arg	tgc Cys	tgc Cys	tac Tyr 620	tgc Cys	tgc Cys	cgc Arg	1932
gtg Val	tgc Cys 625	tgc Cys	cgc Arg	gcg Ala	tgc Cys	tgc Cys 630	ttg Leu	ctg Leu	tgt Cys	ggc Gly	tgc Cys 635	ccc Pro	aag Lys	tgc. Cys	tgc Cys	1980
cgc Arg 640	tgc Cys	agc Ser	aag Lys	tgc Cys	tgc Cys 645	gag Glu	gac Asp	ttg Leu	gag Glu	gag Glu 650	gcg Ala	cag Gln	gag Glu	GJ A GGG	cag Gln 655	2028
gat Asp	gtc Val	cct Pro	gtc Val	aag Lys 660	gct Ala	cct Pro	gag Glu	acc Thr	ttt Phe 665	gat Asp	aac Asn	ata Ile	acc Thr	att Ile 670	agc Ser	2076
aga Arg	gag Glu	gct Ala	cag Gln 675	ggt Gly	gag Glu	gtc Val	cct Pro	gcc Ala 680	tcg Ser	gac Asp	tca Ser	aag Lys	acc Thr 685	gaa Glu	tgc Cys	2124
acg Thr	gcc Ala	ttg Leu 690	tagg	ggac	gc c	ccag	rattg	ıt ca	ıggga	tggg	ggg	atgg	tcc			2173
ttga	gttt	tg c	atgo	tctc	c to	cctc	ccac	ttc	tgca	ccc	tttc	acca	cc t	cgag	gagat	2233
ttgc	tccc	ca t	tago	gaat	g aa	attg	atgo	agt	ccta	aaa	aaaa	aaa				2280
<210	> 5	1								•						
<211	> 6	90														
<212	> P	RT														
<213	> N	м_00	6424	SLC	34A2											

<400> 51

Met Ala Pro Trp Pro Glu Leu Gly Asp Ala Gln Pro Asn Pro Asp Lys 1 10 15

Tyr Leu Glu Gly Ala Ala Gly Gln Gln Pro Thr Ala Pro Asp Lys Ser 20 25 30

Lys Glu Thr Asn Lys Thr Asp Asn Thr Glu Ala Pro Val Thr Lys Ile 35 40 45

Glu Leu Leu Pro Ser Tyr Ser Thr Ala Thr Leu Ile Asp Glu Pro Thr 50 55 60

Glu Val Asp Asp Pro Trp Asn Leu Pro Thr Leu Gln Asp Ser Gly Ile 65 70 75 80

Lys Trp Ser Glu Arg Asp Thr Lys Gly Lys Ile Leu Cys Phe Phe Gln 85 90 95

Gly Ile Gly Arg Leu Ile Leu Leu Leu Gly Phe Leu Tyr Phe Phe Val

Cys Ser Leu Asp Ile Leu Ser Ser Ala Phe Gln Leu Val Gly Gly Lys 115 120 125

Met Ala Gly Gln Phe Phe Ser Asn Ser Ser Ile Met Ser Asn Pro Leu 130 135 140

Leu Gly Leu Val Ile Gly Val Leu Val Thr Val Leu Val Gln Ser Ser 145 150 155 160

Ser Thr Ser Thr Ser Ile Val Val Ser Met Val Ser Ser Ser Leu Leu
. 165 170 175

Thr Val Arg Ala Ala Ile Pro Ile Ile Met Gly Ala Asn Ile Gly Thr 180 185 190

Ser Ile Thr Asn Thr Ile Val Ala Leu Met Gln Val Gly Asp Arg Ser 195 200 205

Glu Phe Arg Arg Ala Phe Ala Gly Ala Thr Val His Asp Phe Phe Asn 210 215 220

Trp Leu Ser Val Leu Val Leu Leu Pro Val Glu Val Ala Thr His Tyr 225 230 235 240

Leu Glu Ile Ile Thr Gln Leu Ile Val Glu Ser Phe His Phe Lys Asn

Gly Glu Asp Ala Pro Asp Leu Leu Lys Val Ile Thr Lys Pro Phe Thr 260 265 270

Lys Leu Ile Val Gln Leu Asp Lys Lys Val Ile Ser Gln Ile Ala Met

280 275 285 Asn Asp Glu Lys Ala Lys Asn Lys Ser Leu Val Lys Ile Trp Cys Lys 295 Thr Phe Thr Asn Lys Thr Gln Ile Asn Val Thr Val Pro Ser Thr Ala Asn Cys Thr Ser Pro Ser Leu Cys Trp Thr Asp Gly Ile Gln Asn Trp 330 Thr Met Lys Asn Val Thr Tyr Lys Glu Asn Ile Ala Lys Cys Gln His 345 Ile Phe Val Asn Phe His Leu Pro Asp Leu Ala Val Gly Thr Ile Leu 360 Leu Ile Leu Ser Leu Leu Val Leu Cys Gly Cys Leu Ile Met Ile Val 375 380 Lys Ile Leu Gly Ser Val Leu Lys Gly Gln Val Ala Thr Val Ile Lys 390 395 Lys Thr Ile Asn Thr Asp Phe Pro Phe Pro Phe Ala Trp Leu Thr Gly 405 410 Tyr Leu Ala Ile Leu Val Gly Ala Gly Met Thr Phe Ile Val Gln Ser 425 Ser Ser Val Phe Thr Ser Ala Leu Thr Pro Leu Ile Gly Ile Gly Val 440 . . 445 Ile Thr Ile Glu Arg Ala Tyr Pro Leu Thr Leu Gly Ser Asm Ile Gly 455 Thr Thr Thr Ala Ile Leu Ala Ala Leu Ala Ser Pro Gly Asn Ala 470 Leu Arg Ser Ser Leu Gln Ile Ala Leu Cys His Phe Phe Phe Asn Ile 490 Ser Gly Ile Leu Leu Trp Tyr Pro Ile Pro Phe Thr Arg Leu Pro Ile 505 Arg Met Ala Lys Gly Leu Gly Asn Ile Ser Ala Lys Tyr Arg Trp Phe Ala Val Phe Tyr Leu Ile Ile Phe Phe Phe Leu Ile Pro Leu Thr Val 530

Phe Gly Leu Ser Leu Ala Gly Trp Arg Val Leu Val Gly Val Gly Val

545					550					555					560	
Pro	Val	Val	Phe	Ile 565	Ile	Ile	Leu	Val	Leu 570	Cys	Leu	Arg	Leu	Leu 575	Gln	4
Ser	Arg	Cys	Pro 580	Arg	Val	Leu	Pro	Lys 585	Lys	Leu	Gln	Asn	Trp 590	Asn	Phe	
Leu	Pro	Leu 595	Trp	Met	Arg	Ser	Leu 600	Lys	Pro	Trp	Asp	Ala 605	Val	Val	Ser	
Lys	Phe 610	Thr	Gly	Суз	Phe	Gln 615	Met	Arg	Суз	Cys	Tyr 620	Cys	Cys	Arg	Val	
Cys 625	Суз	Arg	Ala	Cys	Cys 630	Leu	Leu	Cys	Gly	Cys 635	Pro	Lys	Суз	Cys	Arg 640	
Cys	Ser	ьуз	Сув	Cys 645	Glu	Asp	Leu	Glu	Glu 650	Ala	Gln	Glu	Gly	Gln 655	Asp	
Val	Pro	Val	Lys 660	Ala	Pro	Glu	Thr	Phe 665	Asp	Asn	Ile	Thr	Ile 670	Ser	Arg	•
Glu	Ala	Gln 675	Gly	Glu	Val	Pro	Ala 680	Ser	Asp	Ser	Lys	Thr 685	Glu	Cys	Thr	
Ala	Leu 690										i					
<210	)> 5	52														
<211	.> 5	529														
<212	!> [	NA														
<213	i> P	W959	311,	DKF	'Zp43	4 <b>J</b> 03	7, E	ST .			-					
<220	>															
<221	> n	isc_	feat	ure												
<222	> (	393)	(3	93)												
<223	> t	he r	esid	lue a	t po	siti	on 3	93 i	s A,	TC	or	G				
<400 ctgt		2 .tc a	atgg	ttct	c tt	ccct	tttc	cat	cctc	caa	acct	ggcc	tg a	.gcct	cctga	60
agtt	gctg	ct g	tgaa	tctg	a aa	gact	tgaa	aag	cctc	cac	ctgc	tgtg	tg g	actt	catct	120
caag	gggc	cc a	gcct	cctc	t gg	actc	cacc	ttg	gacc	tca	gtga	ctca	ga a	cttc	tgcct	180
ctaa	acta	ct c	taaa	atcc	а да	ctat	aga t	ata	ttet	cta	aacc	ttca	aa a	ctct	agaat	240

gtccatattt attttatgt tcttggcttt gtgttttagg aaaagtgaat cttgctgt	tt 300
tcaataatgt gaatgctatg ttctgggaaa atccactatg acatctaagt tttgtgta	ca 360
gagagatatt titgcaacta titccacctt cinccacaac cccccacact ccactcca	ca 420
ctcttgagtc tctttaccta atggtctcta cctaatggac cctcgtggcc aaaaagtc	ca 480
ttaaaccaga aaggtgattg gaaaaaaaaa aaaaaaaact cgagggggg	529
<210> 53	
<211> 2100	
<212> DNA	
<213> AF111713 JAM1, junctional adhesion molecule 1	
<220>	
<221> CDS	
<222> (287)(1183)	
<223>	
\2237	
<400> 53	
<400> 53 cgaggccgag agggcgggcc agggccgcac tecggagact cgcggttgct acgcgcacc	a 60
tggctggaga cggagtttcg ctcttgatgc ccagcaggct ggagtgcgat ggcgcgatt	t 120
cggctcactg caacctcctc ctcccagagg tacttctcag ccctctagct ccaactgag	ja 180
acccagccag tcaggaagtc gctacttcgg gaacaccaac caatcagggg gccgtcacc	t 240
gctgaaggag tccttcggcg gctgttgtgt cgggagcctg atcgcg atg ggg aca	295
Met Gly Thr 1	
aag gcg caa gtc gag agg aaa ctg ttg tgc ctc ttc ata ttg gcg atc Lys Ala Gln Val Glu Arg Lys Leu Leu Cys Leu Phe Ile Leu Ala Ile	343
5 10 15	
ctg ttg tgc tcc ctg gca ttg ggc agt gtt aca gtg cac tct tct gaa Leu Leu Cys Ser Leu Ala Leu Gly Ser Val Thr Val His Ser Ser Glu	391
20 25 30 35	
cct gaa gtc aga att cct gag aat aat cct gtg aag ttg tcc tgt gcc Pro Glu Val Arg Ile Pro Glu Asn Asn Pro Val Lys Leu Ser Cys Ala	439
40 45 50	
tac tcg ggc ttt tct tct ccc cgt gtg gag tgg aag ttt gac caa gga Tyr Ser Gly Phe Ser Ser Pro Arg Val Glu Trp Lys Phe Asp Gln Gly	487
55 60 65	
gac acc acc aga ctc gtt tgc tat aat aac aag atc aca gct tcc tat	535
Asp Thr Thr Arg Leu Val Cys Tyr Asn Asn Lys Ile Thr Ala Ser Tyr 70 75 80	
gag gac egg gtg acc ttc ttg cca act ggt atc acc ttc aag tcc gtg	583
Glu Asp Arg Val Thr Phe Leu Pro Thr Gly Ile Thr Phe Lys Ser Val 85 90 95	
·	

aca Thr 100	cgg Arg	gaa Glu	gac Asp	act Thr	ggg Gly 105	aca Thr	tac Tyr	act Thr	tgt Cys	atg Met 110	gtc Val	tct Ser	gag Glu	gaa Glu	ggc Gly 115	631
ggc Gly	aac Asn	agc Ser	tat Tyr	ggg Gly 120	gag Glu	gtc Val	aag Lys	gtc Val	aag Lys 125	ctc Leu	atc Ile	gtg Val	ctt Leu	gtg Val 130	cct Pro	679
cca Pro	tcc Ser	aag Lys	cct Pro 135	aca Thr	gtt Val	aac Asn	atc Ile	ccc Pro 140	tcc Ser	tct Ser	gcc Ala	acc Thr	att Ile 145	GJ À âââ	aac Asn	727
cgg Arg	gca Ala	gtg Val 150	ctg Leu	aca Thr	tgc Cys	tca Ser	gaa Glu 155	caa Gln	gat Asp	ggt Gly	tcc Ser	cca Pro 160	ccț Pro	tct Ser	gaa Glu	775
											acg Thr 175					823
											aat Asn					871
	_	_		_		_		-		_	act Thr		-			919
-		-								_	act Thr					967
											atc Ile					1015
											ttt Phe 255	Gly				1063
_		_	_				_	-		_	aaa Lys			_	_	1111
_	_				-	_				_	agt Ser	_	_	-		1159
	cag Gln							tgaç	jecto	gt d	egget	caco	eg co	ctato	eatct	1213
gcat	ttgc	ect t	acto	aggt	g ct	acto	gact	ctg	gccc	ctg	atgt	ctgt	ag t	ttca	acagga	1273
tgcc	ttat	tt g	tctt	ctad	a co	ccac	aggg	ccc	ccta	ctt	ctto	ggat	gt (	gttt	taata	1333
atgt	cago	ta t	gtgo	ccca	it co	tcct	tcat	gcc	ctcc	ctc	cctt	tcct	ac o	cacto	gctgag	1393
tggc	ctgg	jaa c	ttgt	ttaa	ıa gt	gttt	atto	ccc	attt	ctt	tgaç	ggat	ca ç	ggaag	gaatc	1453
ctgo	gtat	.gc c	atto	jactt	c cc	ttct	aagt	aga	cago	aaa	aatç	gegg	gg g	gtcgc	aggaa	1513
tctg	cact	ca a	ctgo	ccac	c to	gctg	gcag	r gga	tctt	tga:	ataç	gtat	ct t	gago	ttggt	1573
tctg	ggct	ct t	tcct	tgtg	it ac	tgac	gaco	agg	Igcca	gct	gtto	taga	igt g	ggaa	ttaga	1633
ggct	agag	icā ā	ctga	aatç	g tt	gttt	ggtg	atg	acac	tgg	ggto	ctto	ca t	ctct	ggggc	1693

			•			
ccactctctt	ctgtcttccc	atgggaagtg	ccactgggat	ccctctgccc	tgtcctcctg	1753
aatacaagct	gactgacatt	gactgtgtct	gtgġaaaatg	ggagctcttg	ttgtggagag	1813
catagtaaat	tttcagagaa	cttgaagcga	aaaggattta	aaaccgctgc	tctaaagaaa	1873
agaaaactgg	aggetgggeg	cagtggctca	cgcctgtaat	cccagaggct	gaggcaggcg	1933
gatcacctga	ggtcgggagt	tcgggatcag	cctgaccaac	atggagaaac	cctgctggaa	1993
atacagagtt	agccaggcat	ggtggtgcat	gcctgtagtc	ccagctgctc	aggagcctgg	2053
caacaagagc	aaaactccag	ctcaaaaaaa	aaaaaaaaa	aaaaaaa	•	2100
				-	•	
<210> 54						

<211> 299

<212> PRT

<213> AF111713 JAM1, junctional adhesion molecule 1

<400> 54

Met Gly Thr Lys Ala Gln Val Glu Arg Lys Leu Leu Cys Leu Phe Ile 1  $\phantom{\bigg|}$  5  $\phantom{\bigg|}$  10  $\phantom{\bigg|}$  15

Leu Ala Ile Leu Leu Cys Ser Leu Ala Leu Gly Ser Val Thr Val His  $20 \hspace{1cm} 25 \hspace{1cm} 30$ 

Ser Ser Glu Pro Glu Val Arg Ile Pro Glu Asn Asn Pro Val Lys Leu 35 40 45

Ser Cys Ala Tyr Ser Gly Phe Ser Ser Pro Arg Val Glu Trp Lys Phe 50 55 60

Asp Gln Gly Asp Thr Thr Arg Leu Val Cys Tyr Asn Asn Lys Ile Thr 65 70 .75 80

Ala Ser Tyr Glu Asp Arg Val Thr Phe Leu Pro Thr Gly Ile Thr Phe 85 90 95

Lys Ser Val Thr Arg Glu Asp Thr Gly Thr Tyr Thr Cys Met Val Ser

Glu Glu Gly Gly Asn Ser Tyr Gly Glu Val Lys Val Lys Leu Ile Val

Leu Val Pro Pro Ser Lys Pro Thr Val Asn Ile Pro Ser Ser Ala Thr 130 135 140

Ile Gly Asn Arg Ala Val Leu Thr Cys Ser Glu Gln Asp Gly Ser Pro

·											00		-			
Pro	Ser	Glu	Tyr	Thr 165	Trp	Phe	Lys	Asp	Gly 170		Val	Met	Pro	Thr 175	Asn	
Pro	Lys	Ser	Thr	Arg	Ala	Phe	Ser	Asn	Ser	Ser	Tyr	Val	Leu	Asn	Pro	
			180					185					190			
Thr	Thr	Gly 195	Glu	Leu	Val	Phe	Asp 200	Pro	Leu	Ser	Ala	.ser 205	Asp	Thr	Gly	
Glu	Туг 210	Ser	Суз	Glu	Ala	Arg 215	Asn	Gly	Tyr	Gly	Thr 220	Pro	Met	Thr	Ser	
Asn 225	Ala	Val	Arg	Met	Glu 230	Ala	Val	Glu	Arg	Asn 235	Val	Gly	Val	Ile	Val 240	
Ala	Ala	Val	Leu	Val 245	Thr	Leu	Ile	Leu	Leu 250	Gly	Ile	Leu	Val	Phe 255	Gly	
Ile	Trp	Phe	Ala 260	Tyr	Ser	Arg	Gly	His 265		Asp	Arg	Thr	Lys 270	Lys	Gly	
Thr	Ser	Ser 275	Lys	Lys	Val	Ile	Tyr 280	Ser	Gln	Pro	Ser	Ala 285	Arg	Ser	Glu	
Gly	Glu 290	Phe	Lys	Gln	Thr	Ser 295	Ser	Phe	Leu	Val						
<210	)> !	55						;		,						
<211	.> 2	2154														
<212	!> I	ONA									•					
<213	> 1	M_00	6636	MTH	FD2,	met	hyle	ne t	etra	hydr	ofol	ate				
<220	<b>&gt;</b>															
<221		DS						•								
<222		77).	. (11	08)												
<223	>															
<400 atat	_	5 gc g	tggc	ccgc	g cg	cgcg:	cttc	cct	cccg	gcg (	cagt	cacc	gg c	gcgg	tctat	60
ggct	gcga	ct t	ctcta	a at Me	g tc t Se	t gc r Ala	t tte	g gc u Al	t gc a Al	c cg	g ct	g ct u Le	g ca u Gl: 10	g cc n Pr	c gcg o Ala	112
cac His	Ser					Leu A					Leu :					160

								•								
aat Asn	gaa Glu 30	gct Ala	gtt Val	gtc Val	att Ile	tct Ser 35	gga Gly	agg Arg	aaa Lys	ctg Leu	gcc Ala 40	cag Gln	cag Gln	atc Ile	aag Lys	208
cag Gln 45	gaa Glu	gtg Val	cgg Arg	cag Gln	gag Glu 50	gta Val	gaa Glu	gag Glu	tgg Trp	gtg Val 55	gcc Ala	tca Ser	ggc Gly	aac Asn	aaa Lys 60	256
cgg Arg	cca Pro	cac His	ctg Leu	agt Ser 65	gtg Val	atc Ile	ctg Leu	gtt Val	ggc Gly 70	gag Glu	aat Asn	cct Pro	gca Ala	agt Ser 75	cac His	304
tcc Ser	tat Tyr	gtc Val	ctc Leu 80	aac Asn	aaa Lys	acc Thr	agg Arg	gca Ala 85	gct Ala	gca Ala	gtt Val	gtg Val	gga Gly 90	atc Ile	aac Asn	352
agt Ser	gag Glu	aca Thr 95	att Ile	atg Met	aaa Lys	cca Pro	gct Ala 100	tca Ser	att Ile	tca Ser	gag Glu	gaa Glu 105	gaa Glu	ttg Leu	ttg Leu	400
aat Asn	tta Leu 110	atc Ile	aat Asn	aaa Lys	ctg Leu	aat Asn 115	aat Asn	gat Asp	gat Asp	aat Asn	gta Val 120	gat Asp	ggc	ctc Leu	ctt Leu	448
gtt Val 125	cag Gln	ttg Leu	cct Pro	ctt Leu	cca Pro 130	gag Glu	cat His	att Ile	gat Asp	gag Glu 135	aga Arg	agg Arg	atc Ile	tgc Cys	aat Asn 140	496
gct Ala	gtt Val	tct Ser	cca Pro	gac Asp 145	aag Lys	gat Asp	gtt Val	gat Asp	ggc Gly 150	ttt Phe	cat His	gta Val	att Ile	aat Asn 155	gta Val	544
gga Gly	cga Arg	atg Met	tgt Cys 160	ttg Leu	gat Asp	cag Gln	tat Tyr	tcc Ser 165	Met	tta Leu	ccg	gct	act Thr 170	cca Pro	tgg Trp	592
ggt Gly	gtg Val	tgg Trp 175	gaa Glu	ata Ile	atc Ile	aag Lys	cga Arg 180	act Thr	ggc Gly	att Ile	cca Pro	acc Thr 185	cta Leu	GJ À âââ	aag Lys	640
aat Asn	gtg Val 190	gtt Val	gtg Val	gct Ala	gga Gly	agg Arg 195	tca Ser	aaa Lys	aac Asn	gtt Val	gga Gly 200	atg Met	ccc Pro	att Ile	gca Ala	688
atg Met 205	tta Leu	ctg Leu	cac His	aca Thr	gat Asp 210	GJA aaa	gcg Ala	cat His	gaa Glu	cgt Arg 215	ccc Pro	gga Gly	ggt Gly	gat Asp	gcc Ala 220	736
act Thr	gtt Val	aca Thr	ata Ile	tct Ser 225	cat His	cga Arg	tat Tyr	act Thr	ccc Pro 230	Lys	gag Glu	cag Gln	ttg Leu	aag Lys 235	Lys	784
cat His	aca Thr	att Ile	ctt Leu 240	gca Ala	gat Asp	att Ile	gta Val	ata Ile 245	tct Ser	gct Ala	gca Ala	ggt Gly	att Ile 250	cca Pro	aat Asn	832
ctg Leu	atc Ile	aca Thr 255	Ala	gat Asp	atg Met	atc Ile	aag Lys 260	Glu	gga Gly	gca Ala	gca Ala	gtc Val 265	att Ile	gat Asp	gtg Val	880
gga Gly	ata Ile 270	Asn	aga Arg	gtt Val	cac His	gat Asp 275	cct Pro	gta Val	act Thr	gcc Ala	aaa Lys 280	Pro	aag Lys	ttg Leu	gtt Val	928
gga Gly 285	gat Asp	gtg Val	gat Asp	ttt Phe	gaa Glu 290	gga Gly	gtc Val	aga Arg	caa Gln	aaa Lys 295	Ala	Gly	tat Tyr	atc Ile	act Thr 300	976

cca gtt cct gga ggt gtt ggc ccc atg aca gtg gca atg cta atg aag Pro Val Pro Gly Gly Val Gly Pro Met Thr Val Ala Met Leu Met Lys 305 310 315	1024
aat acc att att gct gca aaa aag gtg ctg agg ctt gaa gag cga gaa Asn Thr Ile Ile Ala Ala Lys Lys Val Leu Arg Leu Glu Glu Arg Glu 320 325 330	1072
gtg ctg aag tct aaa gag ctt ggg gta gcc act aat taactactgt Val Leu Lys Ser Lys Glu Leu Gly Val Ala Thr Asn 335 340	1118
gtcttctgtg tcacaaacag cactccaggc cagctcaaga agcaaagcag gccaatagaa	1178
atgcaatatt tttaatttat tctactgaaa tggtttaaaa tgatgccttg tatttattga	1238
aagettaaat gggtgggtgt ttetgeacat acetetgeag taceteacea gggageatte	1298
cagtatcatg cagggtcctg tgatctagcc aggagcagcc attaacctag tgattaatat	1358
gggagacatt accatatgga ggatggatgc ttcactttgt caagcacctc agttacacat	1418
tegeetttte taggattgea ttteecaagt getattgeaa taacagttga taeteatttt	1478
aggtaccaga ccttttgagt tcaactgate aaaccaaagg aaaagtgttg ctagagaaaa	1538
ttggggaaaa ggtgaaaaag aaaaaatggt agtaattgag cagaaaaaaa ttaatttata	1598
tatgtattga ttggcaacca gatttatcta agtagaactg aattggctag gaaaaaagaa	1658
amactgcatg ttaatcattt tectaagetg teettttgag gettagteag tttattggga	1718
aaatgtttag gattatteet tgetattagt acteatttta tgtatgttae cetteagtaa	1778
gttctcccca ttttagtttt ctaggactga aaggattctt ttctacatta tacatgtgtg	1838
ttgtcatatt tggcttttgc tatatacttt aacttcattg ttaaattttt gtattgtata	1898
gtttctttgg tgtatcttaa aacctatttt tgaaaaacaa acttggcttg ataatcattt	1958
gggcagettg ggtaagtacg caacttactt ttccaccaaa gaactgtcag cagetgeetg	2018
cttttctgtg atgtatgtat cctgttgact tttccagaaa ttttttaaga gtttgagtta	2078
ctattgaatt taatcagact ttctgattaa agggttttct ttctttttta ataaaacaca	2138
tctgtctggt atggta	2154

<210> 56

<211> 344

<212> PRT

<213> NM\_006636 MTHFD2, methylene tetrahydrofolate

<400> 56

Leu Arg Leu Arg Pro Phe His Leu Ala Ala Val Arg Asn Glu Ala Val 20 25 30

- Val Ile Ser Gly Arg Lys Leu Ala Gln Gln Ile Lys Gln Glu Val Arg 35 40 45
- Gln Glu Val Glu Glu Trp Val Ala Ser Gly Asn Lys Arg Pro His Leu 50 60
- Ser Val Ile Leu Val Gly Glu Asn Pro Ala Ser His Ser Tyr Val Leu 65 70 75 80
- Asn Lys Thr Arg Ala Ala Ala Val Val Gly Ile Asn Ser Glu Thr Ile 85 90 95
- Met Lys Pro Ala Ser Ile Ser Glu Glu Glu Leu Leu Asn Leu Ile Asn 100 105 110
- Lys Leu Asn Asn Asp Asp Asn Val Asp Gly Leu Leu Val Gln Leu Pro 115 120 125
- Leu Pro Glu His Ile Asp Glu Arg Arg Ile Cys Asn Ala Val Ser Pro 130 135 140
- Asp Lys Asp Val Asp Gly Phe His Val Ile Asn Val Gly Arg Met Cys 145 150 155 160
- Leu Asp Gln Tyr Ser Met Leu Pro Ala Thr Pro Trp Gly Val Trp Glu
  165 170 175
- Ile Ile Lys Arg Thr Gly Ile Pro Thr Leu Gly Lys Asn Val Val 180 185 190
- Ala Gly Arg Ser Lys Asn Val Gly Met Pro Ile Ala Met Leu Leu His 195 200 205
- Thr Asp Gly Ala His Glu Arg Pro Gly Gly Asp Ala Thr Val Thr Ile 210 215 220
- Ser His Arg Tyr Thr Pro Lys Glu Gln Leu Lys Lys His Thr Ile Leu 225 230 235 240
- Ala Asp Ile Val Ile Ser Ala Ala Gly Ile Pro Asn Leu Ile Thr Ala 245 250 255
- Asp Met Ile Lys Glu Gly Ala Ala Val Ile Asp Val Gly Ile Asn Arg 260 265 270
- Val His Asp Pro Val Thr Ala Lys Pro Lys Leu Val Gly Asp Val Asp 275 280 285
- Phe Glu Gly Val Arg Gln Lys Ala Gly Tyr Ile Thr Pro Val Pro Gly 290 295 300

Gly Val Gly Pro Met Thr Val Ala Met Leu Met Lys Asn Thr Ile Ile Ala Ala Lys Lys Val Leu Arg Leu Glu Glu Arg Glu Val Leu Lys Ser 330 Lys Glu Leu Gly Val Ala Thr Asn 340 <210> 57 <211> 1117 <212> DNA <213> NM\_006149 galectin 4, LGALS4 <220> <221> CDS <222> (57)..(1025) <223> 59 atctcccact cetgeagete ttetcacagg accageeact agegeageet egageg atg Met 107 gcc tat gtc ccc gca ccg ggc tac cag ccc acc tac aac ccg acg ctg Ala Tyr Val Pro Ala Pro Gly Tyr Gln Pro Thr Tyr Asn Pro Thr Leu 10 155 cct tac tac cag ccc atc ccg ggc ggg ctc aac gtg gga atg tct gtt Pro Tyr Tyr Gln Pro Ile Pro Gly Gly Leu Asn Val Gly Met Ser Val 203 tac atc caa gga gtg gcc agc gag cac atg aag cgg ttc ttc gtg aac Tyr Ile Gln Gly Val Ala Ser Glu His Met Lys Arg Phe Phe Val Asn 40 251 ttt gtg gtt ggg cag gat ccg ggc tca gac gtc gcc ttc cac ttc aat Phe Val Val Gly Gln Asp Pro Gly Ser Asp Val Ala Phe His Phe Asn ccg cgg ttt gac ggc tgg gac aag gtg gtc ttc aac acg ttg cag ggc 299 Pro Arg Phe Asp Gly Trp Asp Lys Val Val Phe Asn Thr Leu Gln Gly 70 347 ggg aag tgg ggc agc gag gag agg aag agg agc atg ccc ttc aaa aag Gly Lys Trp Gly Ser Glu Glu Arg Lys Arg Ser Met Pro Phe Lys Lys 85 90 395 ggt gec gec ttt gag etg gte tte ata gte.etg get gag eac tae aag Gly Ala Ala Phe Glu Leu Val Phe Ile Val Leu Ala Glu His Tyr Lys 105

gtg gtg gta aat gga aat ccc tte tat gag tac ggg cac cgg ctt ccc Val Val Val Asn Gly Asn Pro Phe Tyr Glu Tyr Gly His Arg Leu Pro 443

144

115	120	125	
cta cag atg gtc acc cac Leu Gln Met Val Thr His 130	ctg caa gtg gat ggg Leu Gln Val Asp Gly 140	Asp Leu Gln Leu Gln	491
tca atc aac ttc atc gga	ggc cag ccc ctc cgc	g eee cag gga eee eeg	539
Ser Ile Asn Phe Ile Gly	Gly Gln Pro Leu Arg	g Pro Gln Gly Pro Pro	
150	155	160	
atg atg cca cct tac cct	ggt ccc gga cat tgc	e cat caa cag ctg aac	587
Met Met Pro Pro Tyr Pro	Gly Pro Gly His Cys	s His Gln Gln Leu Asn	
165	170	175	
age ctg ccc acc atg gas	gga ccc cca acc tto	e aac eeg eet gtg eea	635
Ser Leu Pro Thr Met Glu	Gly Pro Pro Thr Phe	e Asn Pro Pro Val Pro	
180	185	190	
tat ttc ggg agg ctg caa	gga ggg ctc aca gct	c cga aga acc atc atc	683
Tyr Phe Gly Arg Leu Glr	Gly Gly Leu Thr Ala	a Arg Arg Thr Ile Ile	
195	200	205	
atc aag ggc tat gtg cct lle Lys Gly Tyr Val Pro 210 215	Pro Thr Gly Lys Ser	r Phe Ala Ile Asn Phe	731
aag gtg ggc tcc tca ggg	gac ata gct ctg cac	c att aat ccc cgc atg	779
Lys Val Gly Ser Ser Gly	Asp Ile Ala Leu His	s Ile Asn Pro Arg Met	
230	235	240	
ggc aac ggt acc gtg gtc	cgg aac agc ctt ctc	g aat ggc tcg tgg gga	827
Gly Asn Gly Thr Val Val	Arg Asn Ser Leu Leu	u Asn Gly Ser Trp Gly	
245	250	255	
tcc gag gag aag aag atc	acc cac aac cca ttt	t ggt ccc gga cag ttc	875
Ser Glu Glu Lys Lys Ile	Thr His Asn Pro Phe	e Gly Pro Gly Gln Phe	
260	265	.270	
ttt gat ctg tcc att cgc	tgt ggc ttg gat cgc	c ttc aag gtt tac gcc	923
Phe Asp Leu Ser Ile Arc	Cys Gly Leu Asp Arc	g Phe Lys Val Tyr Ala	
275	280	285	
aat ggc cag cac ctc ttc Asn Gly Gln His Leu Pho 290 295	Asp Phe Ala His Arg	g Leu Ser Ala Phe Gln	971
agg gtg gac aca ttg gaa	atc cag ggt gat gto	c acc ttg tcc tat gtc	1019
Arg Val Asp Thr Leu Glu	The Gln Gly Asp Va	l Thr Leu Ser Tyr Val	
310	315	320	
cag atc taatctattc ctgo	ggccat aactcatggg aa	aaacagaat tatcccctag	1075
gactcctttc taagccccta a	taaaatgtc tgagggtgtc	e te	1117
<210> 58			
<211> 323 <212> PRT			
<213> NM_006149 galect	in 4, LGALS4		

Met Ala Tyr Val Pro Ala Pro Gly Tyr Gln Pro Thr Tyr Asn Pro Thr 1 5 10 15

Leu Pro Tyr Tyr Gln Pro Ile Pro Gly Gly Leu Asn Val Gly Met Ser 20 25 30

Val Tyr Ile Gln Gly Val Ala Ser Glu His Met Lys Arg Phe Phe Val 35 40 45

Asn Phe Val Val Gly Gln Asp Pro Gly Ser Asp Val Ala Phe His Phe 50 55 60

Asn Pro Arg Phe Asp Gly Trp Asp Lys Val Val Phe Asn Thr Leu Gln 65 70 75 80

Gly Gly Lys Trp Gly Ser Glu Glu Arg Lys Arg Ser Met Pro Phe Lys 85 90 95

Lys Gly Ala Ala Phe Glu Leu Val Phe Ile Val Leu Ala Glu His Tyr 100 105 110

Lys Val Val Val Asn Gly Asn Pro Phe Tyr Glu Tyr Gly His Arg Leu 115 120 125

Pro Leu Gln Met Val Thr His Leu Gln Val Asp Gly Asp Leu Gln Leu 130 135 140

Gln Ser Ile Asn Phe Ile Gly Gly Gln Pro Leu Arg Pro Gln Gly Pro 145 150 155 160

Pro Met Met Pro Pro Tyr Pro Gly Pro Gly His Cys His Gln Gln Leu 165 170 175

Asn Ser Leu Pro Thr Met Glu Gly Pro Pro Thr Phe Asn Pro Pro Val

Pro Tyr Phe Gly Arg Leu Gln Gly Gly Leu Thr Ala Arg Arg Thr Ile 195 200 205

. Ile Ile Lys Gly Tyr Val Pro Pro Thr Gly Lys Ser Phe Ala Ile Asn 210 215 220

Phe Lys Val Gly Ser Ser Gly Asp Ile Ala Leu His Ile Asn Pro Arg 225 230 235 240

Met Gly Asn Gly Thr Val Val Arg Asn Ser Leu Leu Asn Gly Ser Trp 245 250 255

Gly Ser Glu Glu Lys Lys Ile Thr His Asn Pro Phe Gly Pro Gly Gln 260 265 270

456

									•							•
Phe	Phe	Asp 275	Leu	Ser	Ile	Arg	Cys 280	Gly	Leu	Asp	Arg	Phe 285	Lys	Val	Tyr	
Ala	Asn 290	Gly	Gln	His	Leu	Phe 295	Asp	Phe	Ala	His	Arg 300	Leu	Ser	Ala	Phe	
Gln 305	Arg	Val	Asp	Thr	Leu 310		Ile	Gln	Gly	Asp 315	Val	Thr	Leu	Ser	Tyr 320	
Val	Gln	Ile		•												
<21	0>	59													•	
<21	1>	3697						-								
<21	2>	DNA														
<21	3> :	0_MM	0406	3 ca	dher	in 1	7, C	DH17								
<22	0>										-					
<22	1>	CDS														
<22	2>	(121	) (:	2616	)											
<22	3>															
<40 agg	_	59 att (	ccca	agga	aorat	tacto	ecaqi	t cai	tage	aaga	atc	cqa	cca (	ctga	atggaa	60
															aaact	120
_	ata								• -		_					168
	Ile															
	gca Ala															216
	atg Met															264
	cag Gln 50															312
	aca Thr								Arg							360
	aga Arg															408

90

gca gcc ctg gac gct aat gga att ata gtg gag ggt cca gtc cct atc Ala Ala Leu Asp Ala Asn Gly Ile Ile Val Glu Gly Pro Val Pro Ile

			100	)				105	5				110	)		
Thr	ata Ile	gaa Glu	ı Val	aaq Lys	g gad	ato Ile	aac Asn 120	Asp	aat Asr	cga Arg	Pro	acq Thi	Phe	t cto	c cag ı Gln	504
tca Ser	аа <u>с</u> Буз 130	Туг	gaa Glu	ggo Gl	tca Sei	gta Val 135	. Arg	g Cag	aac Asr	tct Ser	cgc Arg	, Pro	gga Gly	a aaq y Lys	g ccc s Pro	552
ttc Phe 145	Leu	tat Tyr	gto Val	aat Asr	gco Ala 150	Thr	gac Asp	cto Lev	gat Asp	gat Asp 155	Pro	geo Ala	act Thr	ccc Pro	aat Asn 160	600
ggc	cag Gln	ctt Leu	tat Tyr	tac Tyr 165	Glr	att Ile	gtc Val	ato Ile	cag Gln 170	Lev	cco Pro	ato Met	ato : Ile	aac Asr 175	aat Asn	648
gtc Val	atg Met	tac Tyr	Phe 180	Gln	r ato	aac Asn	aac Asn	aaa Lys 185	Thr	gga	gco Ala	ato Ile	tct Ser 190	Let	acc Thr	696
cga Arg	gag Glu	gga Gly 195	Ser	cag Gln	gaa Glu	ttg Leu	aat Asn 200	Pro	gct Ala	aag Lys	aat Asn	Pro 205	Ser	tat Tyr	aat Asn	744
ctg Leu	gtg Val 210	Ile	tca Ser	gtg Val	aag Lys	gac Asp 215	atg Met	gga Gly	Gly	cag Gln	agt Ser 220	Glu	aat	tco Ser	ttc Phe	792
agt Ser 225	gat Asp	acc Thr	aca Thr	tct Ser	gtg Val 230	Asp	atc Ile	ata Ile	gtg Val	aca Thr 235	Glu	aat Asn	att	tgg Trp	aaa Lys 240	840
gca Ala	cca Pro	aaa Lys	cct Pro	gtg Val 245	Glu	atg Met	gtg Val	gaa Glu	aac Asn 250	tca Ser	act Thr	gat Asp	cct Pro	Cac His 255	Pro	888
atc Ile	aaa Lys	atc Ile	act Thr 260	cag Gln	gtg Val	cgg	tgg Trp	aat Asn 265	gat Asp	ccc Pro	ggt Gly	gca Ala	caa Gln 270	Tyr	tcc Ser	936
tta Leu	gtt Val	gac Asp 275	aaa Lys	gag Glu	aag Lys	ctg Leu	cca Pro 280	aga Arg	ttc Phe	cca Pro	ttt Phe	tca Ser 285	att Ile	gac Asp	cag Gln	984
gaa Glu	gga Gly 290	gat Asp	att Ile	tac Tyr	gtg Val	act Thr 295	cag Gln	ccc Pro	ttg Leu	gac Asp	cga Arg 300	gaa Glu	gaa Glu	aag Lys	gat Asp	1032
gca Ala 305	tat Tyr	gtt Val	ttt Phe	tat Tyr	gca Ala 310	gtt Val	gca Ala	aag Lys	gat Asp	gag Glu 315	tac Tyr	gga Gly	aaa Lys	cca Pro	ctt Leu 320	1080
tca Ser	tat Tyr	ccg Pro	ctg Leu	gaa Glu 325	att Ile	cat His	gta Val	aaa Lys	gtt Val 330	aaa Lys	gat Asp	att Ile	aat Asn	gat Asp 335	aat Asn	1128
cca Pro	cct Pro	aca Thr	tgt Cys 340	ccg Pro	tca Ser	cca Pro	gta Val	acc Thr 345	gta Val	ttt Phe	gag Glu	gtc Val	cag Gln 350	gag Glu	aat Asn	1176
gaa Glu	cga Arg	ctg Leu 355	ggt Gly	aac Asn	agt Ser	atç Ile	360 GJ Å āāā	acc Thr	ctt Leu	act Thr	gca Ala	cat His 365	gac Asp	agg Arg	gat Asp	1224
gaa Glu	gaa Glu	aat Asn	act Thr	gcc Ala	aac Asn	agt Ser	ttt Phe	cta Leu	aac Asn	tac Tyr	agg Arg	att Ile	gtg Val	gag Glu	caa Gln	1272

	370	)				375		,			380	)				
act Thr 385	Pro	aaa Lys	ctt Lev	Pro	atg Met 390	Asp	gga Gly	cto Lev	tto Phe	cta Lev 395	·Ile	caa e Glr	aco Thi	tat Tyi	gct Ala 400	1320
gga Gly	atg Met	tta Lev	cag Gln	tta Lev 405	Ala	aaa Lys	caģ Gln	tcc Ser	ttg Leu 410	Lys	aaq Lys	g caa s Glr	gat Asp	act Thi 415	cct Pro	1368
				Thr					Āsp					Thr	ctt Leu	1416
tgt Cys	ttt Phe	gtg Val 435	Gln	atc Ile	aac Asn	gtt Val	att Ile 440	Asp	atc Ile	aat Asn	gat Asp	cag Glr 445	Ile	ccc Pro	atc Ile	1464
ttt Phe	gaa Glu 450	Lys	tca Ser	gat Asp	tat Tyr	gga Gly 455	Asn	Leu	act Thr	ctt Leu	gct Ala 460	Glu	gac Asp	aca Thr	aac Asn	1512
att Ile 465	Gly	tcc Ser	acc	atc Ile	tta Leu 470	Thr	atc Ile	cag Gln	gcc Ala	act Thr 475	Asp	gct Ala	gat Asp	gag Glu	r cca Pro 480	1560
ttt Phe	act Thr	GJ Å āāā	agt Ser	tct Ser 485	Lys	att Ile	ctg Leu	tat Tyr	cat His 490	atc Ile	ata Ile	aag Lys	gga Gly	gac Asp 495	agt Ser	1608
gag Glu	gga Gly	cgc Arg	ctg Leu 500	GJ A Gaa	gtt Val	gac Asp	aca Thr	gat Asp 505	ccc Pro	cat His	acc Thr	aac Asn	acc Thr 510	G1y	tat Tyr	1656
gtc Val	ata Ile	att Ile 515	aaa Lys	aag Lys	cct Pro	ctt Leu	gat Asp 520	ttt Phe	gaa Glu	aca Thr	gca Ala	gct Ala 525	gtt Val	tcc Ser	aac Asn	1704
att Ile	gtg Val 530	ttc Phe	aaa Lys	gca Ala	gaa Glu	aat Asn 535	cct Pro	gag Glu	cct Pro	cta Leu	gtg Val 540	ttt Phe	ggt Gly	gtg Val	aag Lys	1752
tac Tyr 545	aat Asn	gca Ala	agt Ser	tct Ser	ttt Phe 550	gcc Ala	aag Lys	ttc Phe	acg Thr	ctt Leu 555	att Ile	gtg Val	aca Thr	gat Asp	gtg Val 560	1800
aat Asn	gaa Glu	gca Ala	cct Pro	caa Gln 565	ttt Phe	tcc Ser	caa Gln	cac His	gta Val 570	ttc Phe	caa Gln	gcg Ala	aaa Lys	gtc Val 575	agt Ser	1848
gag Glu	gat Asp	gta Val	gct Ala 580	ata Ile	ggc Gly	act Thr	aaa Lys	gtg Val 585	Gly ggc	aat Asn	gtg Val	act Thr	gcc Ala 590	aag Lys	gat Asp	1896
cca Pro	gaa Glu	ggt Gly 595	ctg Leu	gac Asp	ata Ile	agc Ser	tat Tyr 600	tca Ser	ctg Leu	agg. Arg	gga Gly	gac Asp 605	aca Thr	aga Arg	ggt Gly	1944
tgg Trp	ctt Leu 610	aaa Lys	att Ile	gac Asp	cac His	gtg Val 615	act Thr	ggt Gly	gag Glu	atc Ile	ttt Phe 620	agt Ser	gtg Val	gct Ala	cca Pro	1992
ttg Leu 625	gac Asp	aga Arg	gaa Glu	gcc Ala	gga Gly 630	agt Ser	cca Pro	tat Tyr	cgg Arg	gta Val 635	caa Gln	gtg Val	gtg Val	gcc Ala	aca Thr 640	2040
gaa Glu	gta Val	GJ Å āāā	G1 y ggg	tct Ser	tcc Ser	ttg Leu	agc Ser	tct Ser	gtg Val	tca Ser	gag Glu	ttc Phe	cac His	ctg Leu	atc Ile	2088

				•	
	645	650	· .	<b>655</b> .	
ctt atg gat gtg Leu Met Asp Val 660	aat gac aac o Asn Asp Asn	cct ecc agg Pro Pro Arg 665	cta gcc aag Leu Ala Lys	gac tac acg 2 Asp Tyr Thr 670	136
ggc ttg ttc ttc Gly Leu Phe Phe 675	Cys His Pro	ctc agt gca Leu Ser Ala 680	cct gga agt Pro Gly Ser 685	ctc att ttc 2 Leu Ile Phe	184
gag gct act gat Glu Ala Thr Asp 690					232
ttt tcc ctc ggc Phe Ser Leu Gly 705				-	280
atc aat ggt act Ile Asn Gly Thr			Arg His Thr		328
gag agg gag tat Glu Arg Glu Tyr 740					2376
ccc ttg gaa ggc Pro Leu Glu Gly 755	Ile Val Ser			, , , ,	2424
gaa gga agt tgt Glu Gly Ser Cys 770					472
gtg ggc atg gca Val Gly Met Ala 785					:520
ata att tta gca Ile Ile Leu Ala		-	Lys Lys Asp	J J	:568
gat aat gtt gaa Asp Asn Val Glu 820					616
tgaatttgaa aagga	aatgtt tgaatt	tata tagcaa	gtgc tatttcaç	gca acaaccatct 2	676
catcctatta cttt	tcatct aacgtg	catt ataatt	tttt aaacagat	at tecetettgt 2	736
cctttaatat ttgc	taaata tttctt	tttt gaggtg	gagt cttgctct	gt cgcccaggct 2	796
ggagtacagt ggtg	tgatec cagetea	actg caacct	cege etectgge	gtt cacatgattc 2	856
tectgeetea gette	cctaag tagctg	ggtt tacagg	cacc caccacca	itg cccagctaat 2	916
ttttgtattt ttaa					976
acgtcaagtg atct				,	036
acttagatat ttcat					096
tcctctctgc aaatq		•			156
taaatattot gtaca				,	216
gcctggattc catt	atgit tittete	yatt ccatco	tyty teeeette	at coetgactee 3	210

						•	•										
tttgg	gtat	tt c	cacto	aatt	t ca	aaca	tttc		.gaga	aga	aaaa	cgto	Jag 9	gacto	i.k agga	a 33	336
aaata	aaat	.aa a	taaa	agaa	c ag	goott	ttcc	ctt	agta	atta	acaç	gaaat	gt i	ttct	gtgtc	a 33	396
ttaac	ccat	ct t	taat	caat	g to	gacat	gtto	g cto	ettte	gct	gaaa	ttct	tc a	aactt	ggaa	a 34	156
tgaca	acag	ac c	caca	ıgaaç	g to	jttca	aaca	a caa	accta	actc	tgca	aacc	ett (	ggtaa	agga	a 35	516
ccagt	cag	rct g	gcca	ıgatt	t co	ctcac	ctaco	tgo	cato	gcat	acat	gctg	gcg (	catgt	tttc	t 35	576
tcatt	cgt	at c	gttac	gtaaa	g tt	ttg	gttat	tat	tatat	tta	acat	gtgg	gaa (	gaaaa	acaag	a 36	636
catga	aaaa	ga g	gtggt	gaca	a at	caag	gaata	a aac	cacto	gtt	gtag	gtcag	gtt 1	ttgti	tgtt	a 36	696
a																. 30	597
<210>	> 6	0															
<211>	> 8	32															
<212>	> F	RT															
<213>	> N	м_ос	14063	3 cac	iheri	in 17	7, CI	)H17									
<400>	> 6	50					•		٠٠ -								
Met 1 1	lle	Leu	Gln	Ala 5	His	Leu	His	Ser	Leu 10	Суз	Leu	Leu	Met	Leu 15	Tyr		
Leu A	Ala	Thr	Gly 20	Tyr	Gly	Gln	Glu	Gly 25	Lys	Phe	Ser	Gly	Pro 30	Leu	Lys		
Pro N	Met	Thr 35	Phe	Ser	Ile	Tyr	Glu 40	Gly	Gln	Glu	Pro	Ser 45	Gln	Ile	Ile		
Phe (	51n 50	Phe	Lys	Ala	Asn	Pro 55	Pro	Ala	Val	Thr	Phe 60	Glu	Leu	Thr	Gly		
Glu 1 65	ľhr	Asp	Asn	Ile	Phe 70	Val	Ile	Glu	Arg	Glu 75	Gly	Leu	Leu	Tyr	Tyr 80		
Asn A	Arg	Ala	Leu	Asp 85	Arg	Glu	Thr	Arg	Ser 90	Thr	His	Asn	Leu	G1n .95	Val		
Ala <i>F</i>	Ala	Leu	Asp 100	Ala	Asn	Gly	Ile	Ile 105	Val	Glu	Gly	Pro	Val 110	Pro	Ile		
Thr 1	Ile	Glu 115		Lys	Asp	Ile	Asn 120	Asp	Asn	Arg	Pro	Thr 125	Phe	Leu	Gľn		•
Ser I	Lys L30	Tyr	Glu	Gly	Ser	Val 135	Arg	Gln	Asn	Ser	Arg 140	Pro	Gly	Lys	Pro		

Phe Leu Tyr Val Asn Ala Thr Asp Leu Asp Asp Pro Ala Thr Pro Asn 145 150 155 160

- Gly Gln Leu Tyr Tyr Gln Ile Val Ile Gln Leu Pro Met Ile Asn Asn 165 170 175
- Val Met Tyr Phe Gln Ile Asn Asn Lys Thr Gly Ala Ile Ser Leu Thr 180 185 190
- Arg Glu Gly Ser Gln Glu Leu Asn Pro Ala Lys Asn Pro Ser Tyr Asn 195 200 205
- Leu Val Ile Ser Val Lys Asp Met Gly Gly Gln Ser Glu Asn Ser Phe 210 215 220
- Ser Asp Thr Thr Ser Val Asp Ile Ile Val Thr Glu Asn Ile Trp Lys 225 230 235 240
- Ala Pro Lys Pro Val Glu Met Val Glu Asn Ser Thr Asp Pro His Pro 245 250 255
- Ile Lys Ile Thr Gln Val Arg Trp Asn Asp Pro Gly Ala Gln Tyr Ser 260 265 270
- Leu Val Asp Lys Glu Lys Leu Pro Arg Phe Pro Phe Ser Ile Asp Gln 275 280 285
- Glu Gly Asp Ile Tyr Val Thr Gln Pro Leu Asp Arg Glu Glu Lys Asp 290 295 300
- Ala Tyr Val Phe Tyr Ala Val Ala Lys Asp Glu Tyr Gly Lys Pro Leu 305 310 315 320
- Ser Tyr Pro Leu Glu Ile His Val Lys Val Lys Asp Ile Asn Asp Asn 325 330 335
- Pro Pro Thr Cys Pro Ser Pro Val Thr Val Phe Glu Val Gln Glu Asn 340 345 350
- Glu Arg Leu Gly Asn Ser Ile Gly Thr Leu Thr Ala His Asp Arg Asp 355 360 365
- Glu Glu Asn Thr Ala Asn Ser Phe Leu Asn Tyr Arg Ile Val Glu Gln 370 380
- Thr Pro Lys Leu Pro Met Asp Gly Leu Phe Leu Ile Gln Thr Tyr Ala 385 390 395 400
- Gly Met Leu Gln Leu Ala Lys Gln Ser Leu Lys Lys Gln Asp Thr Pro
  405 410 415
- Gln Tyr Asn Leu Thr Ile Glu Val Ser Asp Lys Asp Phe Lys Thr Leu 420 425 430

- Cys Phe Val Gln Ile Asn Val Ile Asp Ile Asn Asp Gln Ile Pro Ile 435 440 445
- Phe Glu Lys Ser Asp Tyr Gly Asn Leu Thr Leu Ala Glu Asp Thr Asn 450 455 460
- Ile Gly Ser Thr Ile Leu Thr Ile Gln Ala Thr Asp Ala Asp Glu Pro 465 470 475 480
- Phe Thr Gly Ser Ser Lys Ile Leu Tyr His Ile Ile Lys Gly Asp Ser 485 490 495
- Glu Gly Arg Leu Gly Val Asp Thr Asp Pro His Thr Asn Thr Gly Tyr 500 505 510
- Val Ile Ile Lys Lys Pro Leu Asp Phe Glu Thr Ala Ala Val Ser Asn 515 520 525
- Ile Val Phe Lys Ala Glu Asn Pro Glu Pro Leu Val Phe Gly Val Lys 530 535 540
- Asn Glu Ala Pro Gln Phe Ser Gln His Val Phe Gln Ala Lys Val Ser 565 570 575
- Glu Asp Val Ala Ile Gly Thr Lys Val Gly Asn Val Thr Ala Lys Asp 580 585 590
- Pro Glu Gly Leu Asp Ile Ser Tyr Ser Leu Arg Gly Asp Thr Arg Gly 595 600 605
- Trp Leu Lys Ile Asp His Val Thr Gly Glu Ile Phe Ser Val Ala Pro 610 615 620
- Leu Asp Arg Glu Ala Gly Ser Pro Tyr Arg Val Gln Val Val Ala Thr 625 630 635 640
- Glu Val Gly Gly Ser Ser Leu Ser Ser Val Ser Glu Phe His Leu Ile 645 650 655
- Leu Met Asp Val Asn Asp Asn Pro Pro Arg Leu Ala Lys Asp Tyr Thr 660 665 670
- Gly Leu Phe Phe Cys His Pro Leu Ser Ala Pro Gly Ser Leu Ile Phe 675 680 685
- Glu Ala Thr Asp Asp Asp Gln His Leu Phe Arg Gly Pro His Phe Thr 690 695 700

Phe 705	Ser	Leu	Gly	Ser	Gly 710	Ser	Leu	Gln	Asn	Asp 715	Trp	Glu	Val	Ser	Lys 720	
Ile	Asn	Gly	Thr	His 725	Ala	Arg	Leu	Ser	Thr 730	Arg	His	Thr	Glu	Phe 735	Glu	
Glu	Arg	Glu	Tyr 740	Val	Val	Leu	Ile	Arg 745		Asn	Asp	Gly	Gly 750	Arg	Pro	
Pro	Leu	Glu 755	Gly	Ile	Val	Ser	Leu 760	Pro	Val	Thr	Phe	Cys 765	Ser	Cys	Val	
Glu	Gly 770	Ser	Суз	Phe	Arg	Pro 775	Ala	Gly	His	Gln	Thr 780	Gly	Ile	Pro	Thr	
Val 785	Gly	Met	Ala	Val	Gly 790	Ile	Leu	Leu	Thr	Thr 795		Leu	Val	Ile	Gly 800	
Ile	Ile	Leu	Ala	Val 805	Val	Phe	Ile	Arg	Ile 810	Lys	Lys	Ąsp	Lys	Gly 815	Lys	
Asp	Asn	Val	Glu 820	Ser	Ala	Gln	Ala	Ser 825	Glu	Val	Lys	Pro	Leu 830	Arg	Ser	
<210	)> (	51				٠										
<211	.> 2	2920														
<212	?> !	ONA	•							,						
<213	3> 1	M_00	5588	; me	prin	A,	alph	a							•	
									•							
<220		יחמ														
<222		DS (10)	. (22	471												
<223		. 10).	. (22	3/)												
<400	> 6	51														
cttg	cago											u Ph			c ttg r Leu	51
				Ile	gca Ala 20				Ile					Glu		99
					gat Asp											147

aat tta gct gca ggc ttg gac ctc ttt caa ggg gac atc ctc ttg cag Asn Leu Ala Ala Gly Leu Asp Leu Phe Gln Gly Asp Ile Leu Leu Gln  $\,$ 

			50					55	•		·		60			
								Pro				tgg Trp 75				243
att Ile	cct Pro 80	tac Tyr	atc Ile	ttg Leu	gct Ala	gat Asp 85	aat Asn	ttg Leu	GJ Å aaa	ctg Leu	aat Asn 90	gct Ala	aaa Lys	gga Gly	gcc Ala	291
att Ile 95	ctg Leu	tat Tyr	gcc Ala	ttt Phe	gag Glu 100	atg Met	ttc Phe	cgt Arg	ctc	aag Lys 105	tcc Ser	tgt Cys	gtg Val	gat Asp	ttc Phe 110	339
aag Lys	ccc Pro	tat Tyr	gaa Glu	gga Gly 115	gag Glu	agc Ser	tca Ser	tat Tyr	atc Ile 120	ata Ile	ttt Phe	caa Gln	cag Gln	ttt Phe 125	gat Asp	387
ggg G1y	tgc Cys	tgg Trp	tct Ser 130	gag Glu	gtt Val	ggt Gly	gac Asp	caa Gln 135	cat His	gtg Val	gga Gly	cag Gln	aac Asn 140	att Ile	tcc Ser	435
						Tyr						cac His 155				483
cat His	gct Ala 160	ttg Leu	gga Gly	ttt Phe	tac Tyr	cac His 165	gag Glu	cag Gln	tca Ser	agg Arg	acg Thr 170	gac Asp	cgg Arg	gat Asp	gat Asp	531
tat Tyr 175	gtg Val	aac Asn	atc Ile	tgg Trp	tgg Trp 180	gac Asp	caa Gln	att Ile	ctt Leu	tca Ser 185	ggt Gly	tac Tyr	cag Gln	cac His	aac Asn 190	579
												aat Asn				627
												ttt Phe				675
												ttt Phe 235				723
												gag Glu				771
cga Arg 255	atg Met	tac Tyr	aat Asn	tgc Cys	acc Thr 260	aca Thr	act Thr	cac His	act Thr	ctt Leu 265	ttg Leu	gac Asp	cac His	tgt Cys	act Thr 270	819
												acc Thr				867
												gaa Glu				915
						Thr						atg Met 315				963
acc Thr	agc Ser	tcg Ser	G] À ààà	tcc Ser	gcg Ala	gaa Glu	gag Glu	gca Ala	ġcc Ala	cta Leu	ctg Leu	gag Glu	tct Ser	cgg Arg	att Ile	1011

	320					325					330	)				
	Tyr					Gln			ctg Leu		Phe				atg Met 350	1059
					Asp				gtc Val 360	Trp					Asp	1107
				Val					_					Gln	gga Gly	1155
gat Asp	gat Asp	gac Asp 385	His	aat Asn	tgg Trp	aaa Lys	att Ile 390	Ala	cat His	gtg Val	gtg Val	ctc Leu 395	Lys	gag Glu	gaa Glu	1203
cag Gln	aag Lys 400	Phe	cgc Arg	tac Tyr	ctt Leu	ttc Phe 405	Gln	ggc Gly	aca Thr	aaa Lys	ggc Gly 410	Asp	cct Pro	cag Gln	aac Asn	1251
	Thr								atc Ile		Leu					1299
tgc Cys	ccc Pro	aca Thr	ggg ggg	gtc Val 435	Trp	aca Thr	gtc Val	cgg Arg	aat Asn 440	ttc Phe	tcc Ser	caa Gln	gtc Val	ctt Leu 445	gag Glu	1347
aac Asn	acc Thr	agc Ser	aaa Lys 450	ggg Gly	gac Asp	aag Lys	ctt Leu	cag Gln 455	agc Ser	cct Pro	cga Arg	ttc Phe	tac Tyr 460	aat Asn	tcg Ser	1395
gag Glu	gga Gly	tat Tyr 465	ggt Gly	ttt Phe	ggg	gta Val	act Thr 470	tta Leu	tac Tyr	cca Pro	aat Asn	agc Ser 475	aga Arg	gaa Glu	agc Ser	1443
tct Ser	ggt Gly 480	tac Tyr	t.tg Leu	aga Arg	ctt Leu	gct Ala 485	ttt Phe	çat His	gtg Val	tgc Cys	agt Ser 490	G1 y ggg	gag Glu	aac Asn	gat Asp	1491
									aga Arg							1539
*	•	~ `	~ ~	-	_		_	_	agg Arg 520		_					1587
ttc Phe	act Thr	acc Thr	tcg Ser 530	aag Lys	tcg Ser	cac His	aca Thr	tct Ser 535	cca Pro	gcg Ala	ata Ile	aat Asn	gac Asp 540	act Thr	gtc Val	1635
									acc Thr							1683
tgt Cys	ttt Phe 560	aga Arg	agc Ser	atc Ile	gac Asp	ttg Leu 565	ggc Gly	tgg Trp	agt Ser	ggt Gly	ttc Phe 570	att Ile	tcc Ser	cac His	caa Gln	1731
atg Met 575	ctg Leu	aaa Lys	agg Arg	agg Arg	agt Ser 580	ttc Phe	ctg Leu	aaa Lys	aat .Asn	gat Asp 585	gac Asp	ctc Leu	atc Ile	ata Ile	ttt Phe 590	1779
gtg Val	gac Asp	ttt Phe	gaa Glu	gat Asp	atc Ile	acc Thr	cac His	ctc Leu	agc Ser	cag Gln	act Thr	gaa Glu	gtt Val	ccc Pro	tct Ser	1827

	• •	·	
. 595		600	605
aaa ggc aaa aga ctg a Lys Gly Lys Arg Leu S 610	agc ccc caa ggc Ser Pro Gln Gly 615	ctc att ctc caa Leu Ile Leu Gln	ggc cag gag 1875 Gly Gln Glu 620
cag cag gtc tcc gaa g Gln Gln Val Ser Glu G 625	gaa ggt tcg gga Glu Gly Ser Gly 630	aag gcc atg tta Lys Ala Met Leu 635	gag gaa gcc 1923 Glu Glu Ala
cta cct gtc agc ctg a Leu Pro Val Ser Leu S 640	agc cag ggg cag Ser Gln Gly Gln 645	ccc agc cga cag Pro Ser Arg Gln 650	aag cgg tcg 1971 Lys Arg Ser
gtg gag aac aca ggc ( Val Glu Asn Thr Gly 1 655	ccc ctg gag gac Pro Leu Glu Asp 660	cat aac tgg cca His Asn Trp Pro 665	cag tac ttc 2019 Gln Tyr Phe 670
aga gac cca tgt gac o Arg Asp Pro Cys Asp 1 675	cca aac cct tgc Pro Asn Pro Cys	caa aat gac ggc Gln Asn Asp Gly 680	atc tgt gtg · 2067 Ile Cys Val 685
aac gtg aag ggg atg g Asn Val Lys Gly Met 2 690	gcg agc tgc agg Ala Ser Cys Arg 695	tgc atc tct gga Cys Ile Ser Gly	cat gct ttc 2115 His Ala Phe 700
ttc tac acg ggg gag c Phe Tyr Thr Gly Glu 7 705			
ctg ggc atg gtg atc g Leu Gly Met Val Ile G 720			
tcc atc atc gcc atc of Ser Ile Ile Ala Ile 1735			edgeet 2257
gctggcattg gccagaccad	c agcagcacct cc	tccatgca ggcctta	act ttcccatgtt 2317
caatgcagtt tggggcagct	t tttttatcag cc	ttgctttg gatagga	ect ccaaggacta 2377
agcetecage eccatgtgtg	g accettgtca te	tctctgcc ccacata:	att atgttacttt 2437
gctatgtgct cctaatgtat	t ctagtgtgtc ct	gtgacaac actcatca	aca cttcattgta 2497
aatcacttgt tttattgact	t gtctttccta ta	gactgtaa gctccat	gag ggcaggcaca 2557
tgttgttctc attgaccgtg	g ctggccccag tg	cctagatg catggct	ggc acattgttgg 2617
cactcaacaa tggttgaatg	g aataaaacaa ta	aatgaatg aataact	ag atatagaaac 2677
tctcatttat attgcagatt	t gaatatatat ga	tgaaattc ttatgtt	gaa tatgttagaa 2737
tcaaatactc atttttcatt	t agatacagta gt	gtcatcac tctttta	nga tottgttaaa 2797
gatttcaaat aaaggtactt	t ctggcgagcc ag	gctgcaca gcatttg	ett teetetgaga 2857
ttctaagaga aggcctttaa	a taaatttaat aa	atattgag ttagcaa	aaa aaaaaaaaaa 2917
aaa		•	2920

<210> 62

<211> 746

<212> PRT

<213> NM\_005588; meprin A, alpha

<400> 62

Met Ala Trp Ile Arg Ser Thr Cys Ile Leu Phe Phe Thr Leu Leu Phe 1 5 10 15

Ala His Ile Ala Ala Val Pro Ile Lys His Leu Pro Glu Glu Asn Val 20 25 30

His Asp Ala Asp Phe Gly Glu Gln Lys Asp Ile Ser Glu Ile Asn Leu 35 40 45

Ala Ala Gly Leu Asp Leu Phe Gln Gly Asp Ile Leu Leu Gln Lys Ser 50 55 60

Arg Asn Gly Leu Arg Asp Pro Asn Thr Arg Trp Thr Phe Pro Ile Pro 65 70 75 80

Tyr Ile Leu Ala Asp Asn Leu Gly Leu Asn Ala Lys Gly Ala Ile Leu 85 90 95

Tyr Ala Phe Glu Met Phe Arg Leu Lys Ser Cys Val Asp Phe Lys Pro 100 105 110

Tyr Glu Gly Glu Ser Ser Tyr Ile Ile Phe Gln Gln Phe Asp Gly Cys 115 120 125

Trp Ser Glu Val Gly Asp Gln His Val Gly Gln Asn Ile Ser Ile Gly 130 135 140

Gln Gly Cys Ala Tyr Lys Ala Ile Ile Glu His Glu Ile Leu His Ala 145 150 155 160

Leu Gly Phe Tyr His Glu Gln Ser Arg Thr Asp Arg Asp Asp Tyr Val 165 170 175

Asn Ile Trp Trp Asp Gln Ile Leu Ser Gly Tyr Gln His Asn Phe Asp 180 185 190

Thr Tyr Asp Asp Ser Leu Ile Thr Asp Leu Asn Thr Pro Tyr Asp Tyr 195 200 205

Glu Ser Leu Met His Tyr Gln Pro Phe Ser Phe Asn Lys Asn Ala Ser 210 215 220

Val Pro Thr Ile Thr Ala Lys Ile Pro Glu Phe Asn Ser Ile Ile Gly 225 230 235 240 Gln Arg Leu Asp Phe Ser Ala Ile Asp Leu Glu Arg Leu Asn Arg Met 245 250 255

Tyr Asn Cys Thr Thr Thr His Thr Leu Leu Asp His Cys Thr Phe Glu 260 265 270

Lys Ala Asn Ile Cys Gly Met Ile Gln Gly Thr Arg Asp Asp Thr Asp 275 280 285

Trp Ala His Gln Asp Ser Ala Gln Ala Gly Glu Val Asp His Thr Leu 290 295 300

Leu Gly Gln Cys Thr Gly Ala Gly Tyr Phe Met Gln Phe Ser Thr Ser 305 310 315 320

Ser Gly Ser Ala Glu Glu Ala Ala Leu Leu Glu Ser Arg Ile Leu Tyr 325 330 335

Pro Lys Arg Lys Gln Gln Cys Leu Gln Phe Phe Tyr Lys Met Thr Gly 340 345 350

Ser Pro Ser Asp Arg Leu Val Val Trp Val Arg Arg Asp Ser Thr 355 360 365

Gly Asn Val Arg Lys Leu Val Lys Val Gln Thr Phe Gln Gly Asp Asp 370 375 380

Asp His Asn Trp Lys Ile Ala His Val Val Leu Lys Glu Glu Gln Lys 385 390 395 400

Phe Arg Tyr Leu Phe Gln Gly Thr Lys Gly Asp Pro Gln Asn Ser Thr 405 410 415

Gly Gly Ile Tyr Leu Asp Asp Ile Thr Leu Thr Glu Thr Pro Cys Pro 420 425 430

Thr Gly Val Trp Thr Val Arg Asn Phe Ser Gln Val Leu Glu Asn Thr 435 440 445

Ser Lys Gly Asp Lys Leu Gln Ser Pro Arg Phe Tyr Asn Ser Glu Gly 450 455

Tyr Gly Phe Gly Val Thr Leu Tyr Pro Asn Ser Arg Glu Ser Ser Gly 465 470 475 480

Tyr Leu Arg Leu Ala Phe His Val Cys Ser Gly Glu Asn Asp Ala Ile 485 490 495

Leu Glu Trp Pro Val Glu Asn Arg Gln Val Ile Ile Thr Ile Leu Asp 500 505 510

Gln Glu Pro Asp Val Arg Asn Arg Met Ser Ser Ser Met Val Phe Thr 515 520 525

Thr Ser Lys Ser His Thr Ser Pro Ala Ile Asn Asp Thr Val Ile Trp 530 535 540

Asp Arg Pro Ser Arg Val Gly Thr Tyr His Thr Asp Cys Asn Cys Phe 545 550 560

Arg Ser Ile Asp Leu Gly Trp Ser Gly Phe Ile Ser His Gln Met Leu 565 570 575

Lys Arg Arg Ser Phe Leu Lys Asn Asp Asp Leu Ile Ile Phe Val Asp 580 585 590

Phe Glu Asp Ile Thr His Leu Ser Gln Thr Glu Val Pro Ser Lys Gly 595 600 605

Lys Arg Leu Ser Pro Gln Gly Leu Ile Leu Gln Gly Gln Gln Gln 610 615 620

Val Ser Glu Glu Gly Ser Gly Lys Ala Met Leu Glu Glu Ala Leu Pro 625 630 635 640

Val Ser Leu Ser Gln Gly Gln Pro Ser Arg Gln Lys Arg Ser Val Glu 645 650 655

Asn Thr Gly Pro Leu Glu Asp His Asn Trp Pro Gln Tyr Phe Arg Asp 660 665 670

Pro Cys Asp Pro Asn Pro Cys Gln Asn Asp Gly Ile Cys Val Asn Val 675 680 . 685

Lys Gly Met Ala Ser Cys Arg Cys Ile Ser Gly His Ala Phe Phe Tyr 690 695 700

Thr Gly Glu Arg Cys Gln Ser Ala Glu Val His Gly Ser Val Leu Gly 705 710 715 720

Met Val Ile Gly Gly Thr Ala Gly Val Ile Phe Leu Thr Phe Ser Ile 725 730 735

Ile Ala Ile Leu Ser Gln Arg Pro Arg Lys
740 745

<210> 63

<211> 8838

<212> DNA

<213> NM\_015902 EDD, hyperplastic discs protein, DD5

<220> <221> CDS <222> (34) .. (8430) <223> <400> 63 egecetegag tggaggaega gaaggaaage ace atg acg tee ate cat tte gtg 54 . Met Thr Ser Ile His Phe Val 1 gtt cac ccg ctg ccg ggc acc gag gac cag ctc aat gac agg tta cga 102 Val His Pro Leu Pro Gly Thr Glu Asp Gln Leu Asn Asp Arg Leu Arg 10 15 gaa gtt tct gag aag ctg aac aaa tat aat tta aac agc cac ccc cct Glu Val Ser Glu Lys Leu Asn Lys Tyr Asn Leu Asn Ser His Pro Pro 30 ttg aat gta ttg gaa cag gct act att aaa cag tgt gtg gtg gga cca 198 Leu Asn Val Leu Glu Gln Ala Thr Ile Lys Gln Cys Val Val Gly Pro 45 . 50 aat cat gct gcc ttt ctt ctt gag gat ggt aga gtt tgc agg att ggt 246 Asn His Ala Ala Phe Leu Leu Glu Asp Gly Arg Val Cys Arg Ile Gly ttt tca gta cag cca gac aga ttg gaa ttg ggt aaa cct gat aat aat Phe Ser Val Gln Pro Asp Arg Leu Glu Leu Gly Lys Pro Asp Asn Asn gat ggg tca aag ttg aac agc aac tcg ggg gca ggg agg acg tca agg Asp Gly Ser Lys Leu Asn Ser Asn Ser Gly Ala Gly Arg Thr Ser Arg cct ggt agg aca agc gac tct cca tgg ttt ctc tca ggt tct gag act Pro Gly Arg Thr Ser Asp Ser Pro Trp Phe Leu Ser Gly Ser Glu Thr 110 cta ggc agg ctg gca ggc aac acc tta gga agc cgc tgg agt tct gga Leu Gly Arg Leu Ala Gly Asn Thr Leu Gly Ser Arg Trp Ser Ser Gly 125 ' 130 gtg ggt gga agt ggt gga gga tcc tct ggt agg tca tca gct gga gct Val Gly Gly Ser Gly Gly Gly Ser Ser Gly Arg Ser Ser Ala Gly Ala 534 cga gat tcc cgc cgg cag act cga gtt att cgg aca gga cgg gat cga Arg Asp Ser Arg Arg Gln Thr Arg Val Ile Arg Thr Gly Arg Asp Arg 155 160 ggg tot ggg ott ttg ggc agt cag coc cag coa gtt att coa gca tot 582 Gly Ser Gly Leu Leu Gly Ser Gln Pro Gln Pro Val Ile Pro Ala Ser gtc att cca gag gag ctg att tca cag gcc caa gtt gtt tta caa ggc 630 Val Ile Pro Glu Glu Leu Ile Ser Gln Ala Gln Val Val Leu Gln Gly 190 195

aaa too aga agt gto att att oga gaa ott oag aga aca aat ott gat Lys Ser Arg Ser Val Ile Ile Arg Glu Leu Gln Arg Thr Asn Leu Asp

210

205

200

	•	•												-		
	aac Asn															726
	gat Asp															774
	atg Met															822
	att Ile 265	_	-	_	-	_			_	-		_				87.0
	cct Pro															918
	ctc Leu															966
_	tcc Ser	-		_			-	_	Arg			_				1014
	gat Asp															1062
	aag Lys 345															1110
	cag Gln															1158
-	ctg Leu			-		_	_	_	-	_						1206
	tgg Trp															1254
	tta Leu															1302
ata Ile	gtc Val 425	ctc Leu	ctg Leu	tct Ser	gca Ala	aat Asn 430	agc Ser	ata Ile	aga Arg	gca Ala	act Thr 435	gta Val	gct Ala	aca Thr	gaa Glu	1350
aat Asn 440	aac Asn	aag Lys	gtt Val	gct Ala	aca Thr 445	tgg Trp	gtg Val	gat Asp	gaa Glu	act Thr 450	tta Leu	agt Ser	tct Ser	gtg Val	gct Ala 455	1398
	aaa Lys															1446
cgg Arg	ata Ile	gtt Val	tct Ser 475	tta Leu	cat His	tgc Cys	tgt Cys	gcc Ala 480	ctt Leu	tac Tyr	acc Thr	tgc Cys	gct Ala 485	cag Gln	ctg Leu	1494

			Leu			tgg Trp		Val								1542
aaa Lys	atg Met 505	Leu	gag Glu	aaa Lys	gct Ala	aga Arg 510	gca Ala	aaa Lys	aat Asn	aaa Lys	aag Lys 515	Pro	aaa Lys	tcc Ser	agt Ser	1590
gct Ala 520	Gly	att Ile	tct Ser	tca Ser	atg Met 525	ccg Pro	aac Asn	ato	act Thr	gtt Val 530	Gly	acc Thr	cag Gln	gta Val	tgc Cys 535	1638
					Leu	tat Tyr				Ala					Ile	1686
agt Ser	gct Ala	Gly	att Ile 555	cct Pro	aaa Lys	gtt Val	ggt Gly	gtc Val 560	Leu	atg Met	gag Glu	tca Ser	gtt Val 565	tgg Trp	aat Asn	1734
			_			ttt Phe			_			-	_	-		1782
						aaa Lys 590										1830
						gaa Glu										1878
acg Thr	tgt Cys	agt Ser	gat Asp	gca Ala 620	tcc Ser	tca Ser	att Ile	gcc Ala	agc Ser 625	agt Ser	gca Ala	tca Ser	atg Met	cca Pro 630	tac Tyr	1926
aaa Lys	cga Arg	cga Arg	cgg Arg 635	tca Ser	acc Thr	cct Pro	gca Ala	cca Pro 640	aaa Lys	gaa Glu	gag Glu	gaa Glu	aag Lys 645	gtg Val	aat Asn	1974
gaa Glu	gag Glu	cag Gln 650	tgg Trp	tct Ser	ctt Leu	cgg Arg	gaa Glu 655	gtg Val	gtt Val	ttt Phe	gtg Val	gaa Glu 660	gat Asp	gtc Val	aag Lys	2022
aat Asn	gtt Val 665	cct Pro	gtt Val	Gly Gly	aag Lys	gtg Val 670	cta Leu	aaa Lys	gta Val	gat Asp	ggt Gly 675	gcc Ala	tat Tyr	gtt Val	gct Ala	2070
gta Val 680	aaa Lys	ttt Phe	cca Pro	gga Gly	acc Thr 685	tcc Ser	agt Ser	aat Asn	act	aac Asn 690	tgt Cys	cag Gln	aac Asn	agc Ser	tct Ser 695	2118
ggt Gly	cca Pro	gat Asp	gct Ala	gac Asp 700	cct Pro	tct Ser	tct Ser	ctc Leu	ctg Leu 705	cag Gln	gat Asp	tgt Cys	agg Arg	tta Leu 710	ctt Leu	2166
						gtt Val										2214
						act Thr										2262
						aat Asn 750										2310

ctg Leu 760	Lys	act Thr	gga Gly	aat Asn	Trp 765	Val	cga Arg	tac Tyr	tgt Cys	atc Ile 770	Phe	gat Asp	ctt	gct Ala	aca Thr 775	2358
gga Gly	aaa Lys	gca Ala	gaa Glu	cag Gln 780	gaa Glu	aat Asn	aat Asn	ttt Phe	cct Pro 785	Thr	agc Ser	agc Ser	att Ile	gct Ala 790	Phe	2406
ctt Leu	ggt	cag Gln	aat Asn 795	Glu	agg Arg	aat Asn	gta Val	gcc Ala 800	Ile	ttc Phe	açt Thr	gct Ala	gga Gly 805	Gln	gaa Glu	2454
tct Ser	Pro	Ile 810	Ile	ctt Leu	cga	gat Asp	gga Gly 815	Asn	ggt	acc Thr	atc Ile	tac Tyr 820	cca Pro	atg Met	gcc Ala	2502
aaa Lys	gat Asp 825	Суз	atg Met	gga Gly	gga Gly	ata Ile 830	agg Arg	gat Asp	ccc Pro	gat Asp	tgg Trp 835	ctg Leu	gat Asp	ctt Leu	cca Pro	2550
Pro 840	att Ile	agt Ser	agt Ser	ctt Leu	gga Gly 845	atg Met	ggt Gly	gtg Val	cat His	tct Ser 850	tta Leu	ata Ile	aat Asn	ctt Leu	cct Pro 855	2598
gcc Ala	aat Asn	tca Ser	aca Thr	atc Ile 860	aaa Lys	aag Lys	aaa Lys	gct Ala	gct Ala 865	gtt Val	atc Ile	atc Ile	atg Met	gct Ala 870	gta Val	2646
gag Glu	aaa Lys	caa Gln	acc Thr 875	tta Leu	atg Met	caa Gln	cac His	att Ile 880	ctg Leu	cgc Arg	tgt Cys	gac Asp	tat Tyr 885	gag Glu	gcc Ala	2694
tgt Cys	cga Arg	caa Gln 890	tat Tyr	cta Leu	atg Met	aat Asn	ctt Leu 895	gag Glu	caa Gln	gcg Ala	gtt Val	gtt Val 900	tta Leu	gag Glu	cag Gln	2742
aat Asn	cta Leu 905	cag Gln	atg Met	ctg Leu	cag Gln	aca Thr 910	ttc Phe	atc Ile	agc Ser	cac His	aga Arg 915	tgt Cys	gat Asp	gga Gly	aat Asn	2790
cga Arg 920	aat Asn	att Ile	ttg Leu	cat His	gct Ala 925	tgt Cys	gta Val	tca Ser	gtt Val	tgc Cys 930	ttt Phe	cca Pro	acc Thr	agc Ser	aat Asn 935	2838
aaa Lys	gaa Glu	act Thr	aaa Lys	gaa Glu 940	gaa Glu	gag Glu	gaa Glu	gcg Ala	gag Glu 945	cgt Arg	tct Ser	gaa Glu	aga Arg	aat Asn 950	aca Thr	2886
Phe	Ala	Glu	Arg 955	Leu	Ser	Ala	Val	Glu 960	Ala	Ile	Ala		Ala 965	Ile	Ser	2934
gtt Val	gtt Val	tca Ser 970	agt Ser	aat Asn	Gly ggc	Pro	ggt Gly 975	aat Asn	cgg Arg	gct Ala	Gly	tca Ser 980	tca Ser	agt Ser	agc Ser	2982
cga Arg	agt Ser 985	ttg Leu	aga Arg	tta Leu	Arg	gaa Glu i 990	atg Met	atg Met	aga Arg	Arg	tcg Ser 995	ttg Leu .	aga Arg	gca Ala	gct Ala	3030
ggt Gly 1000	ttg Leu	ggt Gly	aga Arg	cat His	gaa Glu 100	gc Al	t gg a Gl	a gc y Al	t tc a Se	a to r Se: 10:	r S	gt g er A			_	3075
gat Asp 1015	cca Pro	gtt Val	tca Ser	ccc Pro	Pro			t cc			r T	gg gt rp Va		-		3120

•	cct Pro 1030	Pro	gcg Ala	g ato a Met	g gat : Asp	cct Pro 103	Asp	ggt Gly	gac Asp	att Ile	gat Asp 1040	Phe	t ato	c cto	g gcc ı Ala		3165
	ccc Pro 1045	Ala	gtg Val	r er?	tct Ser	ctt Leu 1050	Thr	aca Thr	gca Ala	gca	acc Thr 1055	Gly	act Thi	ggt Gl	caa / Gln		3210
	gga Gly 1060	Pro	ago Ser	acc Thr	tcc Ser	act Thr 1065	Ile	cca Pro	ggt Gly	Pro	tcc Ser 1070	Thi	gaq Gli	g cca pro	tct Ser		3255
	gta Val 1075	Val	gaa Glu	tco Ser	aag Lys	gat Asp 1080	Arg	aag Lys	gcg Ala	aat Asn	gct Ala 1085	His	ttt Phe	ata 11e	ttg Leu		3300
	aaa Lys 1090	Leu	tta Leu	tgt Cys	gac Asp	agt Ser 1095	Val	gtt Val	ctc Leu	cag Gln	ccc Pro 1100	Туг	cta Lev	cga Arg	gaa Glu		3345
	ctt Leu 1105	Leu	tct Ser	gcc Ala	aag Lys	gat Asp 1110	Ala	aga Arg	Gly	atg Met	acc Thr 1115	Pro	ttt Phe	ato Met	tca Ser		3390
	gct Ala 1120	gta Val	agt Ser	Gly	cga Arg	gct Ala 1125	Tyr	cct Pro	gct Ala	gca Ala	att Ile 1130	Thr	ato Ile	tta Leu	gaa Glu		3435
	act Thr 1135	gct Ala	cag Gln	aaa Lys	att Ile	gca Ala 1140	Lys	gct Ala	gaa Glu	ata Ile	tcc Ser 1145	Ser	agt Ser	gaa Glu	aaa Lys		3480
	gag Glu 1150	gaa Glu	gat Asp	gta Val	ttc Phe	atg Met 1155	Gly	atg Met	gtt Val	tgc Cys	cca Pro 1160	Ser	ggt	acc	aac Asn		3525
	cct Pro 1165	gat Asp	gac Asp	-tct Ser	cct Pro	tta Leu 1170	Tyr	gtt Val	tta Leu	tgt Cys	tgt Cys 1175	Asn	gac Așp	act Thr	tgc Cys		3570
	agt Ser 1180	ttt Phe	aca Thr	tgg Trp	act Thr	gga Gly 1185	Ala	gag Glu	cac His	att Ile	aac Asn 1190	Gln	gat Asp	att Ile	ttt Phe		3615
	gag Glu 1195	tgt Cys	cga Arg	act Thr	tgt Cys	ggc Gly 1200	Leu	ctg Leu	gag Glu	tca Ser	ctg Leu 1205	Cys	tgt Cys	tgt Cys	acg Thr	·	3660
	gaa Glu 1210	tgt Cys	gca Ala	agg Arg	gtt Val	tgt Cys 1215	cat His	aaa Lys	ggt Gly	cat His	gat Asp 1220	tgc Cys	aaa Lys	ctc Leu	aaa Lys		3705
	cgg Arg 1225	aca Thr	tca Ser	cca Pro	aca Thr	gcc Ala 1230	tac Tyr	tgt Cys	gat Asp	tgt Cys	tgg Trp 1235	gag Glu	aaa Lys	tgt Cys	aaa Lys		3750
	tgt Cys 1240	aaa Lys	act Thr	ctt Leu	att Ile	gct Ala 1245	gga Gly	cag Gln	Lys	tct Ser	gct Ala 1250	cgt Arg	ctt Leu	-			3795
						act Thr 1260							ctg Leu				3840
:						ctc Leu 1275							aca Thr				3885

agg Arg 1285	Glr	ace Th:	g gt r Va	g gad 1 Gli	g cat u His 129	Cy:	t caa	a tad	c ag	g cca g Pro 129	Pr	t cg o Ar	a at g Il	c agg e Arg	3930
gaa Glu 1300	Asp	cg Arg	t aa g As:	c cga n Ara	a aaa g Lys 130	Thi	a gco	a Se	t cc r Pr	t gaa o Glu 131	As	t tc p Se	a ga r As	t atg p Met	3975
cca Pro 1315	Asp	cat His	t gat s Asj	t tta p Le:	a gag ı Glu 1320	Pro	cca Pro	a aga o Arg	a tti g Phe	t gcc e Ala 132	Gl	g ct	t gc u Al	a ttg a Leu	4020
gag Glu 1330	Arg	gtt Val	cta L Lei	a cad	gac Asp 133!	Tr	aat Asr	gco Ala	Let	g aaa u Lys 134	Se	t ate	g at	t atg e Met	4065
ttt Phe 1345	GLY	tcg Ser	g cad	g gag n Glu	aat Asn 1350	Lys	gac Asp	cct Pro	ctt Lei	agt Ser 135	Ala	age a Se	c ag c Se	t aga r Arg	4110
ata Ile 1360	Gly	cat His	ctt Lei	ttg Lev	cca Pro 1365	Glu	gag Glu	caa Glr	gta Val	tac Tyr 137	Let	aat 1 Asr	caq n Gli	g caa n Gln	4155
agt Ser 1375	Gly	aca Thr	att : Ile	cgg Arg	ctg Leu 1380	Asp	tgt Cys	ttc Phe	act Thr	cat His 138	Cys	ctt Lei	ata 1116	a gtt e Val	4200
aag Lys 1390	tgt Cys	aca Thr	gca Ala	gat Asp	att Ile 1395	Leu	ctt Leu	tta Leu	gat Asp	act Thr 1400	Leu	cta Lev	ggt Gly	aca Thr	4245
cta Leu 1405	gtg Val	aaa Lys	gaa Glu	ctc Leu	caa Gln 1410	Asn	aaa Lys	tat Tyr	aca Thr	cct Pro 1415	Gly	cgt Arg	aga Arg	gaa Glu	4290
gaa Glu 1420	gct Ala	att Ile	gct Ala	gtg Val	aca Thr 1425	Met	agg Arg	ttt Phe	cta Leu	cgt Arg 1430	Ser	gtg Val	gca Ala	aga Arg	4335
gtt Val 1435	ttt Phe	gtt Val	att Ile	ctg Leu	agt Ser 1440	Val	gaa Glu	atg Met	gct Ala	tca Ser 1445	Ser	aaa Lys	aag Lys	aaa Lys	4380
aac Asn 1450	aac Asn	ttt Phe	att Ile	cca Pro	cag Gln 1455	Pro	att Ile	gga Gly	aaa Lys	tgc Cys 1460	Lys			ttc Phe	4425
caa Gln 1465	gca Ala	ttg Leu	cta Leu	cct Pro	tac Tyr 1470	Ala	gtg Val	gaa Glu	gaa Glu	ttg Leu 1475	Cys	aac Asn	gta Val	gca Ala	4470
gag Glu 1480	tca Ser	ctg Leu	att Ile	gtt Val	cct Pro 1485	gtc Val	aga Arg	atg Met	GJ À ààà	att Ile 1490	gct Ala	cgt Arg	cca Pro	act Thr	4515
gca d Ala 1 1495	cca Pro	ttt Phe	acc Thr	ctg Leu	gct Ala 1500	agt Ser	act Thr	agc Ser	ata Ile	gat Asp 1505	gcc Ala	atg Met	cag Gln	ggc Gly	4560
agt of Ser (	gaa ( Glu (	gaa Glu	tta Leu	ttt Phe	tca Ser 1515	gtg Val	gaa Glu	cca Pro	cta Leu	cca Pro 1520	cca Pro				4605
tct of Ser A 1525	gat d Asp (	cag Sln	tct Ser	Ser	agc Ser 1530	tcc Ser	agt Ser	cag Gln	Ser	cag Gln 1535	tca Ser	tcc Ser	tac Tyr	atc Ile	4650

															•
atc Ile 154	Ar	g aa g As	t cc n Pr	a ca o Gl	g cag n Gln 154	Arg	g cg g Ar	c ato	c ag	c cag r Gln 155	Se	a ca r Gl	g cc n Pr	c gtt o Val	4695 L
cgg Arg 155	G1	c ag y Ar	a ga g As	t ga p Gl	a gaa u Glu 156	Glı	g gat n Asj	t gat o Asp	at o Il	t gtt e Val 156	Se:	a gc r Al	a ga a As	t gto p Val	4740
gaa Glu 157	GT.	g gt u Va	t ga l Gl	g gt u Va	g gtg l Val 157	Glu	g ggt	t gtg y Val	g gci	t gga a Gly 158	Gl	a ga u Gl	g ga u As	t cat p His	4785
cat His 158	As	t gaa	a ca u Gl	g ga n Gli	a gaa u Glu 1590	His	Gly	g gaa 7 Glu	ı gaa	a aat 1 Asn 159	Ala	t gad a Gli	g gc u Al	a gag a Glu	4830
gga Gly 1600	GL	a cai	t ga s As <sub>l</sub>	t gaq p Gli	g cat 1 His 160	Asp	gaa Glu	a gad Asp	617 636	g agt Ser 161	Ası	ato Mei	g gad	g ctg u Leu	4875
gac Asp 1615	Let	g tta 1 Lei	a gca a Ala	a gca a Ala	a gct a Ala 1620	Glu	aca	gaa Glu	agt Ser	gat Asp 162	Sei	gaa Glu	a agi	t aac r Asn	4920
cac His 1630	Sei	aac Asr	caa Glr	a gat n Asp	aat Asn 1635	Ala	agt	. ggg	cgc	aga Arg 1640	Ser	gti Val	gto Val	e act L Thr	4965
gca Ala 1645	Ala	act Thr	gct Ala	ggt Gly	tca Ser 1650	Glu	gca Ala	gga Gly	gca Ala	agc Ser 1655	Ser	gtt Val	cct Pro	gcc Ala	5010
ttc Phe 1660	Phe	tct Ser	gaa Glu	gat Asp	gat Asp 1665	Ser	caa Gln	tcg Ser	aat Asn	gac Asp 1670	Ser	agt Ser	gat Asp	tct Ser	5055
gat Asp 1675	Ser	agt Ser	agt Ser	agt Ser	cag Gln 1680	Ser	gac Asp	gac Asp	ata Ile	gaa Glu 1685	Gln			ttt Phe	5100
atg Met 1690	Leu	gat Asp	gag Glu	cca Pro	tta Leu 1695	Glu	aga Arg	acc Thr	aca Thr	aat Asn 1700	Ser			gcc Ala	5145
aat Asn 1705	GTÄ	gct Ala	gcc Ala	caa Gln	gct Ala 1710	Pro	cgt Arg	tca Ser	atg Met	cag Gln 1715	$\mathtt{Trp}$			cgc Arg	5190
aac Asn 1720	acc Thr	cag Gln	cat His	cag Gln	cga Arg 1725	gca Ala	gcc Ala	agt Ser	aca Thr	gcc Ala 1730	Pro	tcc Ser	agt Ser	aca Thr	5235
tct Ser 1735	aca Thr	cca Pro	gca Ala	gca Ala	agt Ser 1740	tca Ser	gcg <sup>.</sup> Ala	ggt Gly	ttg Leu	att Ile 1745	Tyr	att Ile	gat Asp	cct Pro	5280
tca Ser 1750	aac Asn	tta Leu	cgc Arg	cgg Arg	agt Ser 1755	ggt Gly	acc Thr	atc Ile	agt Ser	aca Thr 1760	agt Ser	gct Ala	gca Ala	gct Ala	5325
gca Ala 1765	gca Ala	gct Ala	gct Ala	ttg Leu	gaa Glu 1770	gct Ala	agc Ser	aac Asn	gcc Ala	agc Ser 1775	agt Ser	tac Tyr	cta Leu	aca Thr	5370
tct Ser 1780	gca Ala	agc Ser	agt Ser	tta Leu	gcc Ala 1785	agg (	gct Ala	tac Tyr	Ser	att Ile 1790	gtc Val	att Ile	aga Arg	caa Gln	5415

atc Ile 1795	Se	g ga r As	c tt p Le	g at u Me	g ggc t Gly 180	Lev	att Ile	cc Pr	t aa o Ly	g tat s Tyr 180	As	t ca n Hi	c ct s Le	a gta u Val	5460
tac Tyr 1810	Sei	t ca r Gl	g at n Il	t cc e Pr	a gca o Ala 181	Ala	gtg Val	g aa: L Ly:	a tte	g act u Thr 182	Ty.	c car r Gl	a ga n As	t gca p Ala	5505
gta Val 1825	Asr	e tta	a cae u Gl	g aa n Ası	c tat n Tyr 1830	Val	gaa Glu	gaa Gli	a aag 1 Lys	g ctt s Leu 183	Il	t cc	c ac	t tgg r Trp	5550
aac Asn 1840	Trp	g ato Mei	g gte t Vai	c agi	t att r Ile 1849	Met	gat Asp	tci Sei	act	gaa Glu 185	Ala	t caa a Gli	a tta	a cgt u Arg	5595
tat Tyr 1855	Gly	tct Sei	t gca r Ala	a tta a Lei	a gca 1 Ala 1860	Ser	gct Ala	ggt	gat Asp	cct Pro 186	Gly	a cat	cca Pro	a aat o Asn	5640
cat His 1870	Pro	ctt Leu	cac His	gct Ala	tct Ser 1875	Gln	aat Asn	tca Ser	gcg Ala	jaga Arg 1880	Arc	ı gaç g Glı	g ago	g atg g Met	5685
act Thr 1885	Ala	cga Arc	n gaa g Glu	a gaa a Glu	gct Ala 1890	Ser	tta Leu	cga Arg	aca Thr	ctt Leu 1895	Glu	r GJ? r gg¢	aga Arq	a cga g Arg	5730
cgt Arg 1900	Ala	Thr	ttg Lev	j ctt Lev	agc Ser 1905	Ala	cgt Arg	caa Gln	gga Gly	atg Met 1910	Met	tct Ser	gca	a cga a Arg	5775
gga Gly 1915	Asp	ttc Phe	cta Leu	aat Asn	tat Tyr 1920	Ala	ctg Leu	tct	cta Leu	atg Met 1925	Arg	tct Ser	cat	: aat : Asn	5820
gat Asp 1930	gag Glu	cat His	tct Ser	gat Asp	gtt Val 1935	Leu	cca Pro	gtt Val	ttg Leu	gat Asp 1940	Val	tgc Cys	tca Ser	ttg Leu	5865
aag Lys 1945	cat His	gtg Val	gca Ala	tat Tyr	gtt Val 1950	ttt Phe	caa Gln	gca Ala	ctt Leu	ata Ile 1955	Tyr	tgg Trp	att Ile	aag Lys	5910
gca Ala 1960	atg Met	aat Asn	cag Gln	cag Gln	aca Thr 1965	aca Thr	ttg Leu	gat Asp	aca Thr	cct Pro 1970	Gln	cta Leu	gaa Glu	cgc Arg	5955
aaa Lys 1975	agg Arg	acg Thr	cga Arg	gaa Glu	ctc Leu 1980	ttg Leu	gaa Glu	ctg Leu	Gly	att Ile 1985	Asp	aat Asn	gaa Glu	gat Asp	6000
tca Ser 1990	gaa Glu	cat His	gaa Glu	aat Asn	gat Asp 1995	gat Asp	gac Asp	acc Thr	aat Asn	caa Gln 2000	agt Ser	gct Ala		ttg Leu	6045
aat Asn 2005	gat Asp	aag Lys	gat Asp	gat Asp	gac Asp 2010	tct Ser	ctt Leu	cct Pro	gca Ala	gaa Glu 2015		ggc Gly			6090
cat His 2020	cca Pro	ttt Phe	ttc Phe	cga Arg	cgt Arg 2025	tca ( Ser i	gac Asp	tcc Ser	atg Met	aca Thr 2030		ctt Leu.			6135
ata Ile : 2035	ccc Pro	cca Pro	aat Asn	Pro	ttt Phe 2040	gaa q Glu N	gtg /al :	cct Pro	Leu	gct Ala 2045	gaa Glu				6180

		Ala				cat His 2055	Leu					Ala				6225
		Leu				cca Pro 2070	Ser					Ser				6270
, •	agt Ser 2080	Ser				tta Leu 2085	Met					Asp				6315
	cta Leu 2095	Glu				aca Thr 2100	Lys					Āla				6360
	aat Asn 2110	Val				caa Gln 2115	Asn					Glu				6405
	cag Gln 2125	Pro	gtg Val	ctg Leu	cca Pro	gaa Glu 2130	gaa Glu	act Thr	gag Glu	agt Ser	tca Ser 2135	Lys	cca Pro	Gly	cca Pro	6450
	tct Ser 2140					gct Ala 2145						Ser				6495
	gaa Glu 2155					gaa Glu 2160					cct Pro 2165	Leu				6540
	agg Arg 2170					ttt Phe 2175						Ser		gat Asp	_	6585
	ctg Leu 2185					cgc . Arg 2190								agg Arg	-	6630
	ttc Phe 2200	atg Met	gaa Glu	gat Asp	gtt Val	gga Gly 2205	gca Ala	gaa Glu	cct Pro	gga Gly	tca Ser 2210	atc Ile	cta Leu	act Thr	gaa Glu	6675
	ttg Leu 2215	ggt Gly	ggt Gly	ttt Phe	gag Glu	gta Val 2220	aaa Lys	gaa Glu	tca Ser	aaa Lys	ttc Phe 2225	cgc Arg	aga Arg	gaa Glu	atg Met	6720
•	gaa Glu 2230					cag Gln 2235				Asp						6765
į						ctt Leu 2250										6810
i						cga Arg 2265										6855
1	cac His 2275	aga Arg	gta Val	aaa Lys	gtc Val	aca Thr 2280	ttt Phe	aag Lys	gat Asp	gag Glu	cca Pro 2285	gga Gly	gag Glu	ggc Gly	agt Ser	6900
(					Ser	ttt Phe 2295										6945

									•						
tca Ser 2305	Asn					Asn				atc Ile 2315	Gln				6990
aaa Lys 2320	Gly	acc Thr	cac His	aca Thr	agt Ser 2325	Leu	atg Met	cag Gln	aga Arg	tta Leu 2330	Arg	aac Asn	cga Arg	gga Gly	7035
gag Glu 2335	Arg					Glu				gaa Glu 2345	Met				7080
agt Ser 2350	Gly					Ser				cgg Arg 2360	Asp				7125
aga Arg 2365	Arg					Asp				ttt Phe 2375	Arg				7170
gaa Glu 2380	Gly									ttg Leu 2390					7215
cag Gln 2395	Ala									gta Val 2405					7260
cca Pro 2410	Āla					Ile				ttg Leu 2420		-	tta Leu		7305
cca Pro 2425	Ala	cag Gln	ctg Leu	ctt Leu	ctc Leu 2430	Leu	cta Leu	gca Ala	agt Ser	gag Glu 2435	gat Asp		ctg Leu		7350
gca Ala 2440	Arg									att Ile 2450					7395
gaa Glu 2455	Asn									gga Gly 2465					7440
tca Ser 2470										cgc Arg 2480					7485
cga Arg 2485										aca Thr 2495					7530
gac Asp 2500										aaa Lys 2510					7575
act Thr 2515										agg Arg 2525					7620
aga Arg 2530										ctg Leu 2540					7665
ctc Leu 2545										att Ile 2555					7710

							•								
ggt Gly 2560	Arç	aaa J Lys	gto Val	aat Asr	tgg Trp 2565	His	gat Asp	ttt Phe	gct	ttt Phe 2570	Phe	gat Asp	cct Pro	gta Val	7755
atg Met 2575	Tyr				cgg Arg 2580	Gln			Lev		Ser				7800
gat Asp 2590	Ala	gat Asp	gct Ala	gtt Val	ttc Phe 2595	Ser	gca Ala	atg Met	gat Asp	ttg Leu 2600	Ala	ttt Phe	gca Ala	att	7845
gac Asp 2605	Leu	tgt Cys	aaa Lys	gaa Glu	gaa Glu 2610	Gly	gga Gly	gga Gly	cag Gln	gtt Val 2615	Glu	ctc Leu	att Ile	cct Pro	7890
aat Asn 2620	Gly	gta Val	aat Asn	ata Ile	cca Pro 2625	Val	act Thr	cca Pro	cag Gln	aat Asn 2630	Val	tat Tyr	gag Glu	tat Tyr	7935
gtg Val 2635	Arg	aaa Lys	tac Tyr	gca Ala	gaa Glu 2640	His	aga Arg	atg Met	ttg Leu	gta Val 2645	Val	gca Ala	gaa Glu	cag Gln	7980
ccc Pro 2650	Leu	cat His	gca Ala	atg Met	agg Arg 2655	Lys	ggt Gly	cta Leu	cta Leu	gat Asp 2660	Val	ctt Leu	cca Pro	aaa Lys	8025
aat Asn 2665	Ser	tta Leu	gaa Glu	gat Asp	tta Leu 2670	Thr	gca Ala	gaa Glu	gat Asp	ttt Phe 2675	Arg	ctt Leu	ttg Leu	gta Val	8070
aat Asn 2680	Gly	tgc Cys	ggt Gly	gaa Glu	gtc Val 2685	aat Asn	gtg Val	caa Gln	atg Met	ctg Leu 2690	Ile		ttt Phe		8115
tct Ser 2695	ttc Phe	aat Asn	gat Asp	gaa Glu	tca Ser 2700	gga Gly	gaa Glu	aat Asn	gct Ala	gag Glu 2705	aag Lys	ctt Leu	ctg Leu	cag Gln	8160
ttc Phe 2710	aag Lys	cgt Arg	tgg Trp	ttc Phe	tgg Trp 2715	tca Ser	ata Ile	gta Val	gag Glu	aag Lys 2720	atg Met	agc Ser	atg Met	aca Thr	8205
gaa Glu 2725	cga Arg	caa Gln	gat Asp	ctt Leu	gtt Val 2730	tac Tyr	ttt Phe	tgg Trp	aca Thr	tca Ser 2735			tca Ser	_	8250
cca Pro 2740	gcc Ala	agt Ser	gaa Glu	gaa Glu	gga Gly 2745	ttc Phe	cag Gln	cct Pro	atg Met	ccc Pro 2750	tca Ser	atc Ile	aca Thr	ata Ile	8295
aga Arg 2755	cca Pro	cca Pro	gat Asp	gac Asp	caa Gln 2760	cat His	ctt Leu	cct Pro	act Thr	gca Ala 2765			tgc Cys		8340
tct Ser 2770	cga Arg	ctt Leu	tac Tyr	Val	cca Pro 2775	ctc Leu	tat Tyr	tcc Ser	Ser	aaa Lys 2780			ctc Leu		8385
cag Gln 2785	aaa Lys	ttg Leu :	tta Leu	Leu	gcc Ala 2790	att : Ile :	aag Lys	acc Thr	Lys	aat Asn 2795	ttt Phe	ggt Gly	ttt Phe	gtg Val	8430
tagag	tata	a aa	agtg	tgta	ttgc	tgtg	ta a	tatt	acta	g caa	attt	tgt	agat	ttttt	t 8490
ccatt	tgtc	t ata	aaaa	gtta	aagt	ttato	gg a	agtt	aatg	c tgt	cata	ccc	ccct	ggtggi	t 8550

accttaaaga gataaaatgc agacattcct tgctgagttt atagcttaaa ggcctaagga
gcactagcaa catttggcta tattggtttg ctagtcacca acttctgggt ctaaccccag
ccaaagatga cagcagaaca acataattta cactgtgatt tatctttttg ctgaggggaa
aaaaatgtaa atgttctgaa aattcactgc tgcctttgtg gaaactgttt cagcaaaggt
tcttgtatag agggaatagg gaatttcaaa ataaaaaatt aagtatgt
<210> 64
<211> 2799
<212> PRT
<213> NM_015902 EDD, hyperplastic discs protein, DD5
<400> 64
Met Thr Ser Ile His Phe Val Val His Pro Leu Pro Gly Thr Glu Asp 1 5 15
Gln Leu Asn Asp Arg Leu Arg Glu Val Ser Glu Lys Leu Asn Lys Tyr 20 25 30
Asn Leu Asn Ser His Pro Pro Leu Asn Val Leu G1u G1n Ala Thr Ile 35 40 45
Lys Gln Cys Val Val Gly Pro Asn His Ala Ala Phe Leu Leu Glu Asp 50 55 60
Gly Arg Val Cys Arg Ile Gly Phe Ser Val Gln Pro Asp Arg Leu Glu
65 70 75 80
Leu Gly Lys Pro Asp Asn Asn Asp Gly Ser Lys Leu Asn Ser Asn Ser
85 90 95
Gly Ala Gly Arg Thr Ser Arg Pro Gly Arg Thr Ser Asp Ser Pro Trp 100 105 110

Phe Leu Ser Gly Ser Glu Thr Leu Gly Arg Leu Ala Gly Asn Thr Leu 115 120 . 125

Gly Ser Arg Trp Ser Ser Gly Val Gly Gly Ser Gly Gly Ser Ser 130 140

Gly Arg Ser Ser Ala Gly Ala Arg Asp Ser Arg Arg Gln Thr Arg Val 145 150 155 160

Ile Arg Thr Gly Arg Asp Arg Gly Ser Gly Leu Leu Gly Ser Gln Pro 165 170 175

- Gln Pro Val Ile Pro Ala Ser Val Ile Pro Glu Glu Leu Ile Ser Gln 180 185 190
- Ala Gln Val Val Leu Gln Gly Lys Ser Arg Ser Val Ile Ile Arg Glu 195 200 205
- Leu Gln Arg Thr Asn Leu Asp Val Asn Leu Ala Val Asn Asn Leu Leu 210 215 220
- Ser Arg Asp Asp Glu Asp Gly Asp Asp Gly Asp Asp Thr Ala Ser Glu 225 230 230 235
- Ser Tyr Leu Pro Gly Glu Asp Leu Met Ser Leu Leu Asp Ala Asp Ile 245 250 255
- His Ser Ala His Pro Ser Val Ile Ile Asp Ala Asp Ala Met Phe Ser 260 265 270
- Glu Asp Ile Ser Tyr Phe Gly Tyr Pro Ser Phe Arg Arg Ser Ser Leu 275 280 285
- Ser Arg Leu Gly Ser Ser Arg Val Leu Leu Leu Pro Leu Glu Arg Asp 290 295 300
- Ser Glu Leu Leu Arg Glu Arg Glu Ser Val Leu Arg Leu Arg Glu Arg 305 310 315 320
- Arg Trp Leu Asp Gly Ala Ser Phe Asp Asn Glu Arg Gly Ser Thr Ser 325 330 335
- Lys Glu Gly Glu Pro Asn Leu Asp Lys Lys Asn Thr Pro Val Gln Ser 340 345 350
- Pro Val Ser Leu Gly Glu Asp Leu Gln Trp Trp Pro Asp Lys Asp Gly 355 360 365
- Thr Lys Phe Ile Cys Ile Gly Ala Leu Tyr Ser Glu Leu Leu Ala Val 370 380
- Ser Ser Lys Gly Glu Leu Tyr Gln Trp Lys Trp Ser Glu Ser Glu Pro 385 390 395 400
- Tyr Arg Asn Ala Gln Asn Pro Ser Leu His His Pro Arg Ala Thr Phe 405 410 415
- Leu Gly Leu Thr Asn Glu Lys Ile Val Leu Leu Ser Ala Asn Ser Ile 420 425 430
- Arg Ala Thr Val Ala Thr Glu Asn Asn Lys Val Ala Thr Trp Val Asp
  435 440 445

Glu Thr Leu Ser Ser Val Ala Ser Lys Leu Glu His Thr Ala Gln Thr 450 455 460

Tyr Ser Glu Leu Gln Gly Glu Arg Ile Val Ser Leu His Cys Cys Ala 465 470 475 480

Leu Tyr Thr Cys Ala Gln Leu Glu Asn Ser Leu Tyr Trp Gly Val 485 490 495

Val Pro Phe Ser Gln Arg Lys Lys Met Leu Glu Lys Ala Arg Ala Lys 500 505 510

Asn Lys Lys Pro Lys Ser Ser Ala Gly Ile Ser Ser Met Pro Asn Ile 515 520 525

Thr Val Gly Thr Gln Val Cys Leu Arg Asn Asn Pro Leu Tyr His Ala 530 535 540

Gly Ala Val Ala Phe Ser Ile Ser Ala Gly Ile Pro Lys Val Gly Val 545 550 555 560

Leu Met Glu Ser Val Trp Asn Met Asn Asp Ser Cys Arg Phe Gln Leu 565 570 575

Arg Ser Pro Glu Ser Leu Lys Asn Met Glu Lys Ala Ser Lys Thr Thr 580 585 590

Glu Ala Lys Pro Glu Ser Lys Gln Glu Pro Val Lys Thr Glu Met Gly 595 600 605

Pro Pro Pro Ser Pro Ala Ser Thr Cys Ser Asp Ala Ser Ser Ile Ala 610 615 620

Ser Ser Ala Ser Met Pro Tyr Lys Arg Arg Arg Ser Thr Pro Ala Pro 625 630 635 640

Lys Glu Glu Glu Lys Val Asn Glu Glu Gln Trp Ser Leu Arg Glu Val 645 650 655

Val Phe Val Glu Asp Val Lys Asn Val Pro Val Gly Lys Val Leu Lys 660 665 670

Val Asp Gly Ala Tyr Val Ala Val Lys Phe Pro Gly Thr Ser Ser Asn 675 680 685

Thr Asn Cys Gln Asn Ser Ser Gly Pro Asp Ala Asp Pro Ser Ser Leu 690 695 700

Leu Gln Asp Cys Arg Leu Leu Arg Ile Asp Glu Leu Gln Val Val Lys 705 710 715 720 Thr Gly Gly Thr Pro Lys Val Pro Asp Cys Phe Gln Arg Thr Pro Lys 725 730 735

Lys Leu Cys Ile Pro Glu Lys Thr Glu Ile Leu Ala Val Asn Val Asp 740 745 750

Ser Lys Gly Val His Ala Val Leu Lys Thr Gly Asn Trp Val Arg Tyr 755 760 765

Cys Ile Phe Asp Leu Ala Thr Gly Lys Ala Glu Gln Glu Asn Asn Phe 770 780

Pro Thr Ser Ser Ile Ala Phe Leu Gly Gln Asn Glu Arg Asn Val Ala 785 790 795 800

Ile Phe Thr Ala Gly Gln Glu Ser Pro Ile Ile Leu Arg Asp Gly Asn 805 810 815

Gly Thr Ile Tyr Pro Met Ala Lys Asp Cys Met Gly Gly Ile Arg Asp 820 825 830

Pro Asp Trp Leu Asp Leu Pro Pro Ile Ser Ser Leu Gly Met Gly Val 835 840 845

His Ser Leu Ile Asn Leu Pro Ala Asn Ser Thr Ile Lys Lys Lys Ala 850 855 860

Ala Val Ile Ile Met Ala Val Glu Lys Gln Thr Leu Met Gln His Ile 865 870 875 880

Leu Arg Cys Asp Tyr Glu Ala Cys Arg Gln. Tyr Leu Met Asn Leu Glu 885 890 895

Gln Ala Val Val Leu Glu Gln Asn Leu Gln Met Leu Gln Thr Phe Ile 900 905 910

Ser His Arg Cys Asp Gly Asn Arg Asn Ile Leu His Ala Cys Val Ser 915 920 925

Val Cys Phe Pro Thr Ser Asn Lys Glu Thr Lys Glu Glu Glu Glu Ala 930 935 940

Glu Arg Ser Glu Arg Asn Thr Phe Ala Glu Arg Leu Ser Ala Val Glu 945 950 955 960

Ala Ile Ala Asn Ala Ile Ser Val Val Ser Ser Asn Gly Pro Gly Asn 965 970 975

Arg Ala Gly Ser Ser Ser Ser Arg Ser Leu Arg Leu Arg Glu Met Met 980 985 990

- Arg Arg Ser Leu Arg Ala Ala Gly Leu Gly Arg His Glu Ala Gly Ala 995 1000 1005
- Ser Ser Ser Asp His Gln Asp Pro Val Ser Pro Pro Ile Ala Pro 1010 1015 1020
- Pro Ser Trp Val Pro Asp Pro Pro Ala Met Asp Pro Asp Gly Asp 1025 1030 1035
- Ile Asp Phe Ile Leu Ala Pro Ala Val Gly Ser Leu Thr Thr Ala 1040 1045 1050
- Ala Thr Gly Thr Gly Gln Gly Pro Ser Thr Ser Thr Ile Pro Gly 1055 1060 1065
- Pro Ser Thr Glu Pro Ser Val Val Glu Ser Lys Asp Arg Lys Ala 1070 1075 1080
- Asn Ala His Phe Ile Leu Lys Leu Cys Asp Ser Val Val Leu 1085 1090 1095
- Gln Pro Tyr Leu Arg Glu Leu Leu Ser Ala Lys Asp Ala Arg Gly 1100 1105 1110
- Met Thr Pro Phe Met Ser Ala Val Ser Gly Arg Ala Tyr Pro Ala 1115 1120 , 1125
- Ala Ile Thr Ile Leu Glu Thr Ala Gln Lys Ile Ala Lys Ala Glu 1130 1135 1140
- Ile Ser Ser Ser Glu Lys Glu Glu Asp Val Phe Met Gly Met Val 1145 1155
- Cys Pro. Ser Gly Thr Asn Pro Asp Asp Ser Pro Leu Tyr Val Leu 1160 1165 1170
- Cys Cys Asn Asp Thr Cys Ser Phe Thr Trp Thr Gly Ala Glu His
- Ile Asn Gln Asp Ile Phe Glu Cys Arg Thr Cys Gly Leu Leu Glu 1190 1195 1200
- Ser Leu Cys Cys Cys Thr Glu Cys Ala Arg Val Cys His Lys Gly 1205 1210 1215
- His Asp Cys Lys Leu Lys Arg Thr Ser Pro Thr Ala Tyr Cys Asp 1220 1225 1230
- Cys Trp Glu Lys Cys Lys Cys Lys Thr Leu Ile Ala Gly Gln Lys

1235 1240 1245 Ser Ala Arg Leu Asp Leu Leu Tyr Arg Leu Leu Thr Ala Thr Asn 1250 1255 1260 Leu Val Thr Leu Pro Asn Ser Arg Gly Glu His Leu Leu Phe 1265 . 1270 Leu Val Gln Thr Val Ala Arg Gln Thr Val Glu His Cys Gln Tyr 1280 1285 Arg Pro Pro Arg Ile Arg Glu Asp Arg Asn Arg Lys Thr Ala Ser 1295 . 1300 1305 Pro Glu Asp Ser Asp Met Pro Asp His Asp Leu Glu Pro Pro Arg 1310 1315

Phe Ala Gln Leu Ala Leu Glu Arg Val Leu Gln Asp Trp Asn Ala 1325 1330 1335

Leu Lys Ser Met Ile Met Phe Gly Ser Gln Glu Asn Lys Asp Pro 1340 1350

Leu Ser Ala Ser Ser Arg Ile Gly His Leu Leu Pro Glu Glu Gln 1355 1360 1365

Val Tyr Leu Asn Gln Gln Ser Gly Thr Ile Arg Leu Asp Cys Phe 1370 1380

Thr His Cys Leu Ile Val Lys Cys Thr Ala Asp Ile Leu Leu Leu 1385 1390 1395

Asp Thr Leu Leu Gly Thr Leu Val Lys Glu Leu Gln Asn Lys Tyr 1400 1405 1410

Thr Pro Gly Arg Arg Glu Glu Ala Ile Ala Val Thr Met Arg Phe 1415 1420 1425

Leu Arg Ser Val Ala Arg Val Phe Val Ile Leu Ser Val Glu Met 1430 1440

Ala Ser Ser Lys Lys Lys Asn Asn Phe Ile Pro Gln Pro Ile Gly 1445 1450 1455

Lys Cys Lys Arg Val Phe Gln Ala Leu Leu Pro Tyr Ala Val Glu 1460 1465 1470

Glu Leu Cys Asn Val Ala Glu Ser Leu Ile Val Pro Val Arg Met 1475 1480 1485

- Gly Ile Ala Arg Pro Thr Ala Pro Phe Thr Leu Ala Ser Thr Ser 1490 1495 1500
- Ile Asp Ala Met Gln Gly Ser Glu Glu Leu Phe Ser Val Glu Pro 1505 1510 1515
- Leu Pro Pro Arg Pro Ser Ser Asp Gln Ser Ser Ser Ser Gln
  1520 1525 1530
- Ser Gln Ser Ser Tyr Ile Ile Arg Asn Pro Gln Gln Arg Arg Ile 1535 1540 1545
- Ser Gln Ser Gln Pro Val Arg Gly Arg Asp Glu Glu Gln Asp Asp 1550 1555 1560
- Ile Val Ser Ala Asp Val Glu Glu Val Glu Val Glu Gly Val
  1565 1570 1575
- Ala Gly Glu Glu Asp His His Asp Glu Gln Glu Glu His Gly Glu 1580 1585 1590
- Glu Asn Ala Glu Ala Glu Gly Gln His Asp Glu His Asp Glu Asp 1595 1600 1605
- Gly Ser Asp Met Glu Leu Asp Leu Leu Ala Ala Ala Glu Thr Glu 1610 1615 1620
- Ser Asp Ser Glu Ser Asn His Ser Asn Gln Asp Asn Ala Ser Gly 1625 1630 1635
- Arg Arg Ser Val Val Thr Ala Ala Thr Ala Gly Ser Glu Ala Gly 1640 1645 1650
- Ala Ser Ser Val Pro Ala Phe Phe Ser Glu Asp Asp Ser Gln Ser 1655  $\phantom{\bigg|}$  1660  $\phantom{\bigg|}$  1665
- Asn Asp Ser Ser Asp Ser Asp Ser Ser Ser Ser Gln Ser Asp Asp 1670 1680
- Ile Glu Gln Glu Thr Phe Met Leu Asp Glu Pro Leu Glu Arg Thr 1685 1690 1695
- Thr Asn Ser Ser His Ala Asn Gly Ala Ala Gln Ala Pro Arg Ser 1700 1705 1710
- Met Gln Trp Ala Val Arg Asn Thr Gln His Gln Arg Ala Ala Ser 1715 1720 1725
- Thr Ala Pro Ser Ser Thr Ser Thr Pro Ala Ala Ser Ser Ala Gly 1730 1735 1740

- Leu Ile Tyr Ile Asp Pro Ser Asn Leu Arg Arg Ser Gly Thr Ile 1745 1750 1755
- Ser Thr Ser Ala Ala Ala Ala Ala Ala Ala Leu Glu Ala Ser Asn 1760 1765 1770
- Ala Ser Ser Tyr Leu Thr Ser Ala Ser Ser Leu Ala Arg Ala Tyr 1775 1780 1785
- Ser Ile Val Ile Arg Gln Ile Ser Asp Leu Met Gly Leu Ile Pro 1790 1795 1800
- . Lys Tyr Asn His Leu Val Tyr Ser Gln Ile Pro Ala Ala Val Lys 1805 1810 1815
  - Leu Thr Tyr Gln Asp Ala Val Asn Leu Gln Asn Tyr Val Glu Glu 1820 1830
  - Lys Leu Ile Pro Thr Trp Asn Trp Met Val Ser Ile Met Asp Ser 1835 1840 1845
  - Thr Glu Ala Gln Leu Arg Tyr Gly Ser Ala Leu Ala Ser Ala Gly 1850 1855 1860
- Asp Pro Gly His Pro Asn His Pro Leu His Ala Ser Gln Asn Ser 1865 1870 1875
- Ala Arg Arg Glu Arg Met Thr Ala Arg Glu Glu Ala Ser Leu Arg
  1880 1885 1890
- Thr Leu Glu Gly Arg Arg Arg Ala Thr Leu Leu Ser Ala Arg Gln 1895 1900 1905
- Gly Met Met Ser Ala Arg Gly Asp Phe Leu Asn Tyr Ala Leu Ser 1910 1915 1920
- Leu Met Arg Ser His Asn Asp Glu His Ser Asp Val Leu Pro Val 1925 1930 1935
- Leu Asp Val Cys Ser Leu Lys His Val Ala Tyr Val Phe Gln Ala 1940 1945 1950
- Leu Ile Tyr Trp Ile Lys Ala Met Asn Gln Gln Thr Thr Leu Asp 1955 1960 1965
- Thr Pro Gln Leu Glu Arg Lys Arg Thr Arg Glu Leu Leu Glu Leu 1970 1980
- Gly Ile Asp Asn Glu Asp Ser Glu His Glu Asn Asp Asp Thr 1985 1990 1995

- Asn Gln Ser Ala Thr Leu Asn. Asp Lys Asp Asp Asp Ser Leu Pro 2000 2005 2010
- Ala Glu Thr Gly Gln Asn His Pro Phe Phe Arg Arg Ser Asp Ser 2015 2020 2025
- Met Thr Phe Leu Gly Cys Ile Pro Pro Asn Pro Phe Glu Val Pro 2030 2035 2040
- Leu Ala Glu Ala Ile Pro Leu Ala Asp Gln Pro His Leu Leu Gln 2045 2050 2055
- Pro Asn Ala Arg Lys Glu Asp Leu Phe Gly Arg Pro Ser Gln Gly 2060 2065 2070
- Leu Tyr Ser Ser Ser Ala Ser Ser Gly Lys Cys Leu Met Glu Val 2075 2080 2085
- Thr Val Asp Arg Asn Cys Leu Glu Val Leu Pro Thr Lys Met Ser 2090  $\phantom{\bigg|}$  2095  $\phantom{\bigg|}$  2005  $\phantom{\bigg|}$  2100  $\phantom{\bigg|}$
- Tyr Ala Ala Asn Leu Lys Asn Val Met Asn Met Gln Asn Arg Gln 2105 2110 2115
- Lys Lys Glu Glu Glu Glu Gln Pro Val Leu Pro Glu Glu Thr Glu 2120 2125 2130
- Ser Ser Lys Pro Gly Pro Ser Ala His Asp Leu Ala Ala Gln Leu 2135 2140 2145
- Lys Ser Ser Leu Leu Ala Glu Ile Gly Leu Thr Glu Ser Glu Gly 2150 2155
- Pro Pro Leu Thr Ser Phe Arg Pro Gln Cys Ser Phe Met Gly Met 2165 2170 2175
- Val Ile Ser His Asp Met Leu Leu Gly Arg Trp Arg Leu Ser Leu 2180 2185 2190
- Glu Leu Phe Gly Arg Val Phe Met Glu Asp Val Gly Ala Glu Pro 2195 2200 2205
- Gly Ser Ile Leu Thr Glu Leu Gly Gly Phe Glu Val Lys Glu Ser 2210 2215 2220
- Lys Phe Arg Arg Glu Met Glu Lys Leu Arg Asn Gln Gln Ser Arg 2225 2230 2235
- Asp Leu Ser Leu Glu Val Asp Arg Asp Arg Asp Leu Leu Ile Gln 2240 2250

- Gln Thr Met Arg Gln Leu Asn Asn His Phe Gly Arg Arg Cys Ala 2255 2260 2265
- Thr Thr Pro Met Ala Val His Arg Val Lys Val Thr Phe Lys Asp 2270 2275 2280
- Glu Pro Gly Glu Gly Ser Gly Val Ala Arg Ser Phe Tyr Thr Ala 2285 2290 2295
- Ile Ala Gln Ala Phe Leu Ser Asn Glu Lys Leu Pro Asn Leu Glu 2300 2305 2310
- Cys Ile Gln Asn Ala Asn Lys Gly Thr His Thr Ser Leu Met Gln 2315 2320 2325
- Arg Leu Arg Asn Arg Gly Glu Arg Asp Arg Glu Arg Glu Arg Glu 2330 2340
- Asp Arg Asp Arg Asp Phe Arg Arg Gln Leu Ser Ile Asp Thr Arg 2360 2365 2370
- Pro Phe Arg Pro Ala Ser Glu Gly Asn Pro Ser Asp Asp Pro Glu 2375 2380 2385
- Pro Leu Pro Ala His Arg Gln Ala Leu Gly Glu Arg Leu Tyr Pro 2390 2395 2400
- Arg Val Gln Ala Met Gln Pro Ala Phe Ala Ser Lys Ile Thr Gly 2405 2410 2415
- Met Leu Leu Glu Leu Ser Pro Ala Gln Leu Leu Leu Leu Leu Ala 2420 2425 2430
- Ser Glu Asp Ser Leu Arg Ala Arg Val Asp Glu Ala Met Glu Leu 2435 2440 2445
- Ile Ile Ala His Gly Arg Glu Asn Gly Ala Asp Ser Ile Leu Asp 2450 2455 2460
- Leu Gly Leu Val Asp Ser Ser Glu Lys Val Gln Gln Glu Asn Arg 2465 2470 2475
- Lys Arg His Gly Ser Ser Arg Ser Val Val Asp Met Asp Leu Asp 2480 2485 2490
- Asp Thr Asp Asp Gly Asp Asp Asn Ala Pro Leu Phe Tyr Gln Pro 2495 2500 2505

- Gly Lys Arg Gly Phe Tyr Thr Pro Arg Pro Gly Lys Asn Thr Glu 2510 2515 2520
- Ala Arg Leu Asn Cys Phe Arg Asn Ile Gly Arg Ile Leu Gly Leu 2525 2530 2535
- Cys Leu Leu Gln Asn Glu Leu Cys Pro Ile Thr Leu Asn Arg His 2540 2550
- Val Ile Lys Val Leu Leu Gly Arg Lys Val Asn Trp His Asp Phe 2555 2560 2565
- Ala Phe Phe Asp Pro Val Met Tyr Glu Ser Leu Arg Gln Leu Ile 2570 2580
- Leu Ala Ser Gln Ser Ser Asp Ala Asp Ala Val Phe Ser Ala Met 2585 2590 2595
- Asp Leu Ala Phe Ala Ile Asp Leu Cys Lys Glu Glu Gly Gly Gly 2600 2605 2610
- Gln Val Glu Leu Ile Pro Asn Gly Val Asn Ile Pro Val Thr Pro 2615 2620 2625
- Gln Asn Val Tyr Glu Tyr Val Arg Lys Tyr Ala Glu His Arg Met 2630 2635 2640
- Leu Val Val Ala Glu Gln Pro Leu His Ala Met Arg Lys Gly Leu 2645 2655
- Leu Asp Val Leu Pro Lys Asn Ser Leu Glu Asp Leu Thr Ala Glu 2660 2665 2670
- Asp Phe Arg Leu Leu Val Asn Gly Cys Gly Glu Val Asn Val Gln 2675
- Ala Glu Lys Leu Leu Gln Phe Lys Arg Trp Phe Trp Ser Ile Val 2705 2710 2715
- Glu Lys Met Ser Met Thr Glu Arg Gln Asp Leu Val Tyr Phe Trp 2720 2725 2730
- Thr Ser Ser Pro Ser Leu Pro Ala Ser Glu Glu Gly Phe Gln Pro 2735 2740 2745
- Met Pro Ser Ile Thr Ile Arg Pro Pro Asp Asp Gln His Leu Pro 2750 2755 2760

									•							
Thi	276		sn Tl	nr Cy	ys IJ		er 1	Arg 1	Leu '	Tyr '		Pro 2775	Leu	Tyr	Ser	
Ser	Lys 278		ln I	le Le	eu Ly		ln 1 785	Lys 1	Leu :	Leu I		Ala 2790	Ile	Гуз	Thr	
Lys	Asr 279		ne GJ	Ly Ph	ne Va	1		٠								
<21	.0>	65					•									
<21	.1>	4256	;													
<21	.2> .	DNA				•										
<21	3>	NM_C	1499	2 DA	AM1											
<22	0>															
<22		CDS						-								
<22		(126	) (	3359	)											
<22	3>															
- 40	٥.	<b>.</b> -														
	0> gagc		tgtg	aagg	aa a	ctgt	ttaa	ıc cg	gato	ccat	tgt	cacco	aga	gtgc	agagco	: 60
gcc	tttc	cag	catg	cagg	gg c	tgct	cago	g tt	tagt	caca	tca	aagaa	ata	gaac	agaatt	: 120
cag	cc a M 1	tg g et A	cc c la P	ca a ro A	ga aa rg Lj 5	ag a ys A	ga g rg G	gt g ly G	ga c ly A	rg G	gt a ly 1	att t [le S	ca t er P	tc a he I	tc ttt le Phe 15	: 170
tgc Cys	tgt Cys	ttc Phe	cga Arg	aat Asn 20	aat Asn	gat Asp	cac His	cca Pro	gaa Glu 25	atc Ile	acg Thr	tat Tyr	cgg Arg	ctg Leu 30	cga Arg	218
aat Asn	gat Asp	agc Ser	aac Asn 35	ttt Phe	gcg Ala	ctt Leu	cag Gln	acc Thr 40	atg Met	gaa Glu	cca Pro	gca Ala	ttg Leu 45	ccc Pro	atg Met	266
ccc Pro	cct Pro	gtg Val 50	gag Glu	gag Glu	ctg Leu	gat Asp	gtc Val 55	atg Met	ttc Phe	agt Ser	gaa Glu	ctg Leu 60	gtg Val	gat Asp	gaa Glu	314
ctg Leu	gac Asp 65	ctc Leu	aca Thr	gac Asp	aaa Lys	cac His 70	aga Arg	gaa Glu	gcc Ala	atg Met	ttt Phe 75	gca Ala	ctt Leu	cca Pro	gct Ala	362
gag Glu 80	aaa Lys	aaa Lys	tgg Trp	caa Gln	ata Ile 85	tac Tyr	tgt Cys	agc Ser	aag Lys	aaa Lys 90	aag Lys	gac Asp	cag Gln	gaa Glu	gaa Glu 95	410
aac Asn	aag Lys	gga Gly	gct Ala	aca Thr	agt Ser	tgg Trp	cct Pro	gaa Glu	ttc Phe	tac Tyr	att Ile	gat Asp	cag Gln	ctc Leu	aat Asn	458

tcc atg gct gct aga aaa tct ctg ctg gct tta gag aag gaa gaa gaa Ser Met Ala Ala Arg Lys Ser Leu Leu Ala Leu Glu Lys Glu Glu Glu

		٠	115					120					125			
			Ser			ata Ile		Ser					Leu			554
		Met				acc Thr 150						Asp				602
	Ile					aag Lys					Glu					650
cga Arg	ata Ile	cat His	act Thr	tct Ser 180	ctc Leu	att Ile	Gly	tgt Cys	ata Ile 185	Lys	gcg Ala	tta Leu	atg Met	aac Asn 190	aac Asn	698
						gtc Val			His							746
att Ile	gct Ala	cag Gln 210	agt Ser	ctg Leu	agc Ser	aca Thr	gag Glu 215	aac Asn	att Ile	aaa Lys	acg Thr	aag Lys 220	gtg Val	gcc Ala	gtg Val	794
						gtg Val 230										842
gtt Val 240	ctg Leu	cag Gln	gcc Ala	atg Met	ctg Leu 245	cac His	tac Tyr	cag Gln	aag Lys	tat Tyr 250	gcc Ala	agc Ser	gaa Glu	agg Arg	acc Thr 255	890
cgc Arg	ttt Phe	cag Gln	aca Thr	tta Leu 260	att	aac Asn	gac Asp	ttg Leu	gat Asp 265	aaa Lys	agc Ser	act Thr	ggg Gly	cgg Arg 270	tat Tyr	938
cga Arg	gat Asp	gaa Glu	gtg Val 275	agt Ser	ctc Leu	aag Lys	act	gcc Ala 280	atc Ile	atg Met	tcc Ser	ttc Phe	att Ile 285	aat Asn	gca Ala	986
gtg Val	ctc Leu	agc Ser 290	caa Gln	ggt Gly	gca Ala	gga Gly	gtg Val 295	gag Glu	agt Ser	ttg Leu	gac Asp	ttt Phe 300	aga Arg	ctt Leu	cat His	1034
ctt Leu	cgc Arg 305	tat Tyr	gaa Glu	ttt Phe	ctg Leu	atg Met 310	tta Leu	gga Gly	att Ile	caa Gln	cct Pro 315	gta Val	ata Ile	gat Asp	aaa Lys	1082
tta Leu 320	agg Arg	gaa Glu	cac His	gaa Glu	aat Asn 325	tca Ser	aca Thr	tta Leu	gat Asp	agg Arg 330	cat His	tta Leu	gac Asp	ttt Phe	ttt Phe 335	1130
gaa Glu	atg Met	ctc Leu	cga Arg	aat Asn 340	gaa Glu	gat Asp	gaa Glu	cta Leu	gaa Glu 345	ttt Phe	gcc Ala	aaa Lys	aga Arg	ttt Phe 350	gaa Glu	1178
ctg Leu	gtt Val	cac His	ata Ile 355	gac Asp	aca Thr	aaa Lys	agt Ser	gca Ala 360	act Thr	cag Gln	atg Met	ttt Phe	gag Glu 365	ctg Leu	acc Thr	1226
						agt Ser										1274
						atg Met										1322

	385					390				•	395	i -				•
cag Gln 400	Tyr	tgg Trp	cta Leu	cta Leu	cta Leu 405	ı Asp	aga Arg	att Ile	ata Ile	cag Gln 410	Gln	ata Ile	gtt Val	ato . Ile	cag Gln 415	1370
aat Asn	gac Asp	aaa Lys	gga Gly	cag Gln 420	Asp	cct Pro	gac Asp	tcc Ser	aca Thr 425	Pro	ttg Leu	gaa Glu	aac Asn	ttt Phe 430	aat Asn	1418
att	aag Lys	aat Asn	gtc Val 435	Val	cga Arg	atg Met	ttg Leu	gtt Val 440	Asn	gaa Glu	aat Asn	gaa Glu	gtt Val 445	Lys	cag Gln	1466
tgg Trp	aaa Lys	gaa Glu 450	caa Gln	gcg Ala	gaa Glu	aaa Lys	atg Met 455	Arg	aaa Lys	gag Glu	cac	aat Asn 460	Glu	cta Leu	caa Gln	1514
cag Gln	aaa Lys 465	ctg Leu	gaa Glu	aag Lys	aaa Lys	gaa Glu 470	Arg	gaa Glu	tgt Cys	gat Asp	gct Ala 475	Lys	act Thr	caa Gln	gag Glu	1562
aag Lys 480	Glu	gag Glu	atg Met	atg Met	cag Gln 485	acc Thr	tta Leu	aat Asn	aaa Lys	atg Met 490	Lys	gag Glu	aaa Lys	ctt Leu	gaa Glu 495	1610
aag Lys	gag Glu	act	act Thr	gag Glu 500	cat His	aag Lys	caa Gln	gtc Val	aag Lys 505	cag Gln	cag Gln	gtg Val	gcg Ala	gac Asp 510	Leu	1658
aca Thr	gca Ala	cag Gln	ctc Leu 515	His	gag Glu	ctc Leu	agc Ser	agg Arg 520	agg Arg	gcc Ala	gtc Val	tgt Cys	gct Ala 525	tca Ser	atc Ile	1706
cca Pro	ggt Gly	gga Gly 530	ccc Pro	tcg Ser	cct Pro	gga Gly	gca Ala 535	cca Pro	gga Gly	ggg Gly	ccc Pro	ttt Phe 540	cct Pro	tcc Ser	tct Ser	1754
gtg Val	cct Pro 545	gga Gly	tct Ser	ctc Leu	ctt Leu	cct Pro 550	ccc Pro	cca Pro	cca Pro	ccc Pro	cca Pro 555	cct Pro	cta Leu	cca Pro	ggt Gly	1802
260 260 399	atg Met	ctt Leu	ccc Pro	cct Pro	cca Pro 565	ccg Pro	cct Pro	ccc Pro	ctc Leu	cct Pro 570	Pro	ggt Gly	ggc	Pro	cct Pro 575	1850
cct Pro	ccc Pro	cca Pro	GJÀ aaa	cct Pro 580	cct Pro	ccc Pro	tta Leu	GJ À aaa	gca Ala 585	atc Ile	atg Met	cca Pro	cct Pro	cct Pro 590	ggt Gly	1898
gct Ala	cca Pro	atg Met	ggc Gly 595	cta Leu	gca Ala	ctg Leu	aag Lys	aag Lys 600	aaa Lys	agc Ser	att Ile	cct Pro	cag Gln 605	ccc Pro	aca Thr	1946
aat Asn	gcc Ala	ctg Leu 610	aaa Lys	tcc Ser	ttc Phe	aac Asn	tgg Trp 615	tct Ser	aaa Lys	ctg Leu	ccc Pro	gag Glu 620	aac Asn	aaa Lys	ctg Leu	1994
gaa Glu	gga Gly 625	aca Thr	gta Val	tgg Trp	acc Thr	gaa Glu 630	att Ile	gat Asp	gat Asp	aca Thr	aaa Lys 635	gtc Val	ttc Phe	aaa Lys	att Ile	2042
cta Leu 640	gat Asp	ctt Leu	gaa Glu	gac Asp	ctg Leu 645	gaa Glu	aga Arg	acc Thr	ttc Phe	tct Ser 650	gcc Ala	tat Tyr	caa Gln	aga Arg	cag Gln 655	2090
cag Gln	gat Asp	ttc Phe	ttt Phe	gtg Val	aac Asn	agt Ser	aac Asn	tcc Ser	aag Lys	cag Gln	aaa Lys	gaa Glu	gca Ala	gat Asp	gcc Ala	2138

				660	•			•	665		· .			670		
att Ile	gat Asp	gac Asp	act Thr 675	ctg Leu	agt Ser	tcc Ser	aaa Lys	ctt Leu 680	aaa Lys	gtt Val	aaa Lys	gag Glu	ctt Leu 685	tcg Ser	gtg Val	2186
att Ile	gat Asp	ggt Gly 690	cgg Arg	aga Arg	gct Ala	cag Gln	aat Asn 695	tgc Cys	aac Asn	atc Ile	ctt Leu	cta Leu 700	tcg Ser	agg Arg	ttg Leu	2234
aaa Lys	tta Leu 705	tcc Ser	aat Asn	gac Asp	gaa Glu	atc Ile 710	aaa Lys	cgg Arg	gca Ala	att Ile	cta Leu 715	aca Thr	atg Met	gac Asp	gaa Glu	2282
					aag Lys 725											2330
					att Ile											2378
-		_	_	_	gct Ala	-						_	-	_	_	2426
aat Asn					agg Arg											2474
gca Ala	gag Glu 785	cgt Arg	gtg Val	gca Ala	gaa Glu	gtg Val 790	aaa Lys	cct Pro	aaa Lys	gtg Val	gaa Glu 795	gca Ala	att Ile	cgt Arg	tct Ser	2522
800 Gly ggc	tca Ser	gaa Glu	gag Glu	gtg Val	ttt Phe 805	agg Arg	agt Ser	ggt Gly	gcc Ala	ctc Leu 810	aag Lys	cag Gln	ttg Leu	ctg Leu	gag Glu 815	2570
					gga Gly											2618
					ata Ile											2666
	_		-		aac Asn											2714
					ccc											2762
					gcg Ala 885											2810
ata Ile	agt Ser	acc Thr	ttg Leu	aga Arg 900	agt Ser	ggc Gly	ttg Leu	aaa Lys	gca Ala 905	gta Val	gag Glu	aca Thr	gag Glu	ctg Leu 910	gaa Glu	2858
					ccc Pro											2906
					aca Thr											2954

9	930	935	940	
gac ctt d Asp Leu I 945	cta gca gaa gct aaa Leu Ala Glu Ala Lys 950	gac ctg ttt act aaa Asp Leu Phe Thr Lys 955	gca gtg aag cac 3002 Ala Val Lys His	
ttt ggg ( Phe Gly ( 960	gaa gag gct ggc aaa Glu Glu Ala Gly Lys 965	ata caa cca gat gag Elle Gln Pro Asp Glu 970	ttc ttt ggc att 3050 Phe Phe Gly Ile 975	
ttt gat o	caa ttt ctt caa gct Gln Phe Leu Gln Ala 980	gtg tca gaa gcc aaa Val Ser Glu Ala Lys 985	caa gaa aac gaa 3098 Gln Glu Asn Glu 990	
aat atg a Asn Met A	aga aag aaa aag gag Arg Lys Lys Lys Glu 995	gaa gaa gaa cgt cga Glu Glu Glu Arg Arg 1000	gct cgc atg gaa 3146 Ala Arg Met Glu 1005	
Ala Gln I		t gaa agg gaa cgt aa g Glu Arg Glu Arg Ly 1015		
Ala Lys G		a agc gga gag ttt ga u Ser Gly Glu Phe As 1030		
Ser Ala I		a gtg ttt gac aaa ga u Val Phe Asp Lys As 1045		
Leu Lys A	egg aat cgc aaa cg Arg Asn Arg Lys Ar .055	t att acc aac cag at g Ile Thr Asn Gln Me 1060		
Ser Arg G	gag aga cca atc ac lu Arg Pro Ile Th .070	a aaa ctt aat ttc ta r Lys Leu Asn Phe 1075	attttcca 3369	
tgaatactt	t tttttagaaa gctca	ttagc agccctctaa agtg	actaga acgtttcatt 3429	
acactgcct	t gcaatccaaa cagtg	gcaat tttttccttc atct	gtgagt gaatgtgtga 3489	
acgtgtgta	t gtaaatgtat gtgtg	tatat attaaaaaat gtat	atagat gtctgagtgt 3549	
tgtctggag	a cctatacgta tggtt	aaaaa gatttatgtt aatg	tatgtg ctccaaaacc 3609	
tttcgtgta	t gcattcacat tgagt	gtggc tcattttctt tccc	egaacg ccatgactgt 3669	
tcagaagca	c aatactatct cctga	aagag ataagagaca ttcc	ctagat tcaaaggcaa 3729	
aacagaaga	a acaaacaaac aaaca	aacaa agcttgcaaa atat	tttatg gtttccaagc 3789	
ttgatatcc	t ttaaaattat tttca	ttgat ggaactggag ttgt	ggaaa aacatagatt 3849	
taaaatgat	t tttgataget gacat	tgtga tgttgatgta tcaca	atcagt aataggacca 3909	
gctttgaat	t totgacattg gtgtg	gggat acagtctgta aatg	ttatt gagaacatct 3969	
tgcacacaa	t ttgaattatg tagaa	gtca atcaagtttt tgtal	attta aaagttggac 4029	
atcaatttt	t teceetgatt teates	aagtt atctctgcca agtgo	etcttg ataatttett 4089	
cagattttt	g gaaaaaaaca ctata	aaat gcaatccatg cttt	tttaa agaacaacat 4149	
tgccagagta	a tgcttgttct aacaat	atag atatataaac cttaa	aaata ataaaatatc 4209	
tcacccaaga	a cttaaaggaa gaatto	totg aagggataaa gatta	4256	

. <210> 66

<211> 1078

<212> PRT

. <213> NM 014992 DAAM1

<400> 66

Met Ala Pro Arg Lys Arg Gly Gly Arg Gly Ile Ser Phe Ile Phe Cys
1 10 15

Cys Phe Arg Asn Asn Asp His Pro Glu Ile Thr Tyr Arg Leu Arg Asn 20 25 30

Asp Ser Asn Phe Ala Leu Gln Thr Met Glu Pro Ala Leu Pro Met Pro 35 40 45

Pro Val Glu Glu Leu Asp Val Met Phe:Ser Glu Leu Val Asp Glu Leu 50 55 60

Asp Leu Thr Asp Lys His Arg Glu Ala Met Phe Ala Leu Pro Ala Glu 65 70 75 80

Lys Lys Trp Gln Ile Tyr Cys Ser Lys Lys Lys Asp Gln Glu Glu Asn 85 90 95

Lys Gly Ala Thr Ser Trp Pro Glu Phe Tyr Ile Asp Gln Leu Asn Ser 100 105 110

Met Ala Ala Arg Lys Ser Leu Leu Ala Leu Glu Lys Glu Glu Glu Glu 115 120 125

Glu Arg Ser Lys Thr Ile Glu Ser Leu Lys Thr Ala Leu Arg Thr Lys 130 135 140

Pro Met Arg Phe Val Thr Arg Phe Ile Asp Leu Asp Gly Leu Ser Cys 155 150 150

Ile Leu Asn Phe Leu Lys Thr Met Asp Tyr Glu Thr Ser Glu Ser Arg 165 170 175

Ile His Thr Ser Leu Ile Gly Cys Ile Lys Ala Leu Met Asn Asn Ser 180 180 190

Gln Gly Arg Ala His Val Leu Ala His Ser Glu Ser Ile Asn Val Ile 195 200 205

Ala Gln Ser Leu Ser Thr Glu Asn Ile Lys Thr Lys Val Ala Val Leu 210 215 220 Glu Ile Leu Gly Ala Val Cys Leu Val Pro Gly Gly His Lys Lys Val 225 230 235 240

Leu Gln Ala Met Leu His Tyr Gln Lys Tyr Ala Ser Glu Arg Thr Arg 245 250 255

Phe Gln Thr Leu Ile Asn Asp Leu Asp Lys Ser Thr Gly Arg Tyr Arg 260 265 270

Asp Glu Val Ser Leu Lys Thr Ala Ile Met Ser Phe Ile Asn Ala Val 275 280 285

Leu Ser Gln Gly Ala Gly Val Glu Ser Leu Asp Phe Arg Leu His Leu 290 295 300

Arg Tyr Glu Phe Leu Met Leu Gly Ile Gln Pro Val Ile Asp Lys Leu 305 310 315 320

Arg Glu His Glu Asn Ser Thr Leu Asp Arg His Leu Asp Phe Phe Glu 325 330 335

Met Leu Arg Asn Glu Asp Glu Leu Glu Phe Ala Lys Arg Phe Glu Leu 340 345 350

Val His Ile Asp Thr Lys Ser Ala Thr Gln Met Phe Glu Leu Thr Arg 355 360 365

Lys Arg Leu Thr His Ser Glu Ala Tyr Pro His Phe Met Ser Ile Leu 370 375 380

His His Cys Leu Gln Met Pro Tyr Lys Arg Ser Gly Asn Thr Val Gln 385 390 395 400

Tyr Trp Leu Leu Asp Arg Ile Ile Gln Gln Ile Val Ile Gln Asn 405 410 415

Asp Lys Gly Gln Asp Pro Asp Ser Thr Pro Leu Glu Asn Phe Asn Ile 420 425 430

Lys Asn Val Val Arg Met Leu Val Asn Glu Asn Glu Val Lys Gln Trp
435 440 445

Lys Glu Gln Ala Glu Lys Met Arg Lys Glu His Asn Glu Leu Gln Gln 450 .455 460

Lys Leu Glu Lys Lys Glu Arg Glu Cys Asp Ala Lys Thr Gln Glu Lys 465 470 475 480

Glu Glu Met Met Gln Thr Leu Asn Lys Met Lys Glu Lys Leu Glu Lys 485 490 495 Glu Thr Thr Glu His Lys Gln Val Lys Gln Gln Val Ala Asp Leu Thr 500 510

Ala Gln Leu His Glu Leu Ser Arg Arg Ala Val Cys Ala Ser Ile Pro 515 520 525

Gly Gly Pro Ser Pro Gly Ala Pro Gly Gly Pro Phe Pro Ser Ser Val 530 535 540

Pro Gly Ser Leu Leu Pro Pro Pro Pro Pro Pro Pro Leu Pro Gly Gly 545 550 555 560

Met Leu Pro Pro Pro Pro Pro Leu Pro Pro Gly Gly Pro Pro Pro 565 570 575

Pro Pro Gly Pro Pro Pro Leu Gly Ala Ile Met Pro Pro Pro Gly Ala 580 590

Pro Met Gly Leu Ala Leu Lys Lys Lys Ser Ile Pro Gln Pro Thr Asn 595 600 605

Ala Leu Lys Ser Phe Asn Trp Ser Lys Leu Pro Glu Asn Lys Leu Glu 610 615 620

Gly Thr Val Trp Thr Glu Ile Asp Asp Thr Lys Val Phe Lys Ile Leu 625 630 635 640

Asp Leu Glu Asp Leu Glu Arg Thr Phe Ser Ala Tyr Gln Arg Gln Gln 645 650 655

Asp Phe Phe Val Asn Ser Asn Ser Lys Gln Lys Glu Ala Asp Ala Ile 660 665 670

Asp Asp Thr Leu Ser Ser Lys Leu Lys Val Lys Glu Leu Ser Val Ile 675 680 685

Asp Gly Arg Arg Ala Gln Asn Cys Asn Ile Leu Leu Ser Arg Leu Lys 690 695 700

Leu Ser Asn Asp Glu Ile Lys Arg Ala Ile Leu Thr Met Asp Glu Gln 705 710 715 720

Glu Asp Leu Pro Lys Asp Met Leu Glu Gln Leu Leu Lys Phe Val Pro 725 730 735

Glu Lys Ser Asp Ile Asp Leu Leu Glu Glu His Lys His Glu Leu Asp
740 745 750

Arg Met Ala Lys Ala Asp Arg Phe Leu Phe Glu Met Ser Arg Ile Asn

755 760 765

His Tyr Gln Gln Arg Leu Gln Ser Leu Tyr Phe Lys Lys Phe Ala 770 775 780

Glu Arg Val Ala Glu Val Lys Pro Lys Val Glu Ala Ile Arg Ser Gly 785 790 795 800

Ser Glu Glu Val Phe Arg Ser Gly Ala Leu Lys Gln Leu Leu Glu Val 805 810 815

Val Leu Ala Phe Gly Asn Tyr Met Asn Lys Gly Gln Arg Gly Asn Ala 820 825 830

Tyr Gly Phe Lys Ile Ser Ser Leu Asn Lys Ile Ala Asp Thr Lys Ser 835 840 845

Ser Ile Asp Lys Asn Ile Thr Leu Leu His Tyr Leu Ile Thr Ile Val 850 855 860

Glu Asn Lys Tyr Pro Ser Val Leu Asn Leu Asn Glu Glu Leu Arg Asp 865 870 875 880

Ile Pro Gln Ala Ala Lys Val Asn Met Thr Glu Leu Asp Lys Glu Ile 885 890 895

Ser Thr Leu Arg Ser Gly Leu Lys Ala Val Glu Thr Glu Leu Glu Tyr 900 905 910

Gln Lys Ser Gln Pro Pro Gln Pro Gly Asp Lys Phe Val Ser Val Val 915 920 925

Ser Gln Phe Ile Thr Val Ala Ser Phe Ser Phe Ser Asp Val Glu Asp 930 935 940

Leu Leu Ala Glu Ala Lys Asp Leu Phe Thr Lys Ala Val Lys His Phe 945 950 955 960

Gly Glu Glu Ala Gly Lys Ile Gln Pro Asp Glu Phe Phe Gly Ile Phe 965 970 975

Asp Gln Phe Leu Gln Ala Val Ser Glu Ala Lys Gln Glu Asn Glu Asn 980 985 990

Met Arg Lys Lys Glu Glu Glu Glu Arg Arg Ala Arg Met Glu Ala 995 1000 1005

Gln Leu Lys Glu Gln Arg Glu Arg Glu Arg Lys Met Arg Lys Ala 1010 1015 1020

Lys Glu Asn Ser Glu Glu Ser Gly Glu Phe Asp Asp Leu Val Ser

	10	25				1	030					1035				
Ala	a Le 10	u A 40	rg S	er G	ly G	lu V	al 045	Phe	Asp	Lys .		Leu 1050	Ser	Lys	Leu	
Lys	10.	g A 55	sn A	rg L	ys A	rg I.	le '	Thr	Asn	Gln 1		Thr 1065	Asp	Ser	Ser	
Arg	10°	u A 70	rg P:	ro I	le Ti		ys : 075	Leu .	Asn	Phe						
<21	.0>	67	-						. ,							
<21	1>	109	6						٠			•				
<21	2>	DNA														
<21	3>	NM_	00212	22 MI	IC cl	.ass	II I	DQ a.	Lpha	1, 1	ILA-	DQA1				
						•							•			
<22	0>															
<22	1>	CDS														
<22	2>	(1)	. (76	55)							_					
<22	3>															
										•						
atg	0> atc	cta	. aac	aaa	gct	ctg	ctg	ctg	ggg	gca	cto	gct	ctg	acc	acc	48
1	Ile	Leu	Asn	Lys 5	Ala	Leu	Leu	Leu	Gly 10	Ala	Let	Ala	ьeu	Thr 15	Thr	
gtg	atg	ago	ccc	5 tgt	gga	gat	Leu	, gac	10 att	Ala oto	act		cac	15	acc	96
gtg Val	atg Met	agc Ser	ccc Pro 20	tgt Cys	gga Gly ttg	ggt Gly	gaa Glu	gac Asp 25	10 att Ile	gtg Val	gct	gac	cac His 30	15 gtt Val	gcc Ala	96 144
gtg Val tct Ser	atg Met tgt Cys	ago Ser ggt Gly 35	ccc Pro 20 gta Val	tgt Cys aac Asn	gga Gly ttg Leu	ggt Gly tac Tyr	gaa Glu cag Gln 40	gac Asp 25 ttt Phe	att Ile tac Tyr	gtg Val ggt Gly	gct Ala ccc Pro	gac Asp tct Ser	cac His 30 ggc Gly	gtt Val cag Gln	gcc Ala tac Tyr	
gtg Val tct Ser acc Thr	atg Met tgt Cys cat His 50	ago Ser ggt Gly 35 gaa Glu	ccc Pro 20 gta Val ttt Phe	tgt Cys aac Asn gat Asp	gga Gly ttg Leu gga Gly	ggt Gly tac Tyr gat Asp 55	gaa Glu cag Gln 40 gag Glu	gac Asp 25 ttt Phe cag Gln	att Ile tac Tyr ttc Phe	gtg Val ggt Gly tac Tyr	gct Ala ccc Pro gtg Val 60	gac Asp tct Ser 45	cac His 30 ggc Gly ctg Leu	gtt Val cag Gln gag Glu	gcc Ala tac Tyr agg Arg	144
gtg Val tct Ser acc Thr	atg Met tgt Cys cat His 50 gag Glu	agc Ser ggt 35 gaa Glu act Thr	ccc Pro 20 gta Val ttt Phe gcc Ala	tgt Cys aac Asn gat Asp tgg Trp	gga Gly ttg Leu gga Gly cgg Arg 70	ggt Gly tac Tyr gat Asp 55 tgg Trp	gaaa Glu cag Gln 40 gag Glu cct Pro	gac Asp 25 ttt Phe cag Gln gag Glu	att Ile tac Tyr ttc Phe ttc Phe	gtg Val ggt Gly tac Tyr agc Ser 75	gct Ala	gac Asp tct Ser 45 gac Asp ttt Phe	cac His 30 ggc Gly ctg Leu gga Gly	gtt Val cag Gln gag Glu ggt Gly	gcc Ala tac Tyr agg Arg ttt Phe 80	144 192
gtg Val tct Ser acc Thr aag Lys 65 gac	atgt Met tgts Cys cats 50 gag Glu ccg Pro atc	agc Ser ggt Gly 35 gaa Glu act Thr cag Gln atg	ccc Pro 20 gta Val ttt Phe gcc Ala ggt Gly	tgt Cys aac Asn gat Asp tgg Trp gca Ala	gga Gly ttg Leu gga Gly cgg Arg 70 ctg Leu	ggt Gly tac Tyr gat Asp 55 tgg Trp aga Arg	gaa Glu cag Gln 40 gag Glu cct Pro	gac Asp 25 ttt Phe cag Gln gag Glu atg	att Ile tac Tyr ttc Phe ttc Phe acc	gtg Val ggt Gly tac Tyr agc Ser 75 gtg Val	gct Ala ccc gtg Val 60 aaa Lys gca Ala	gac Asp tct Ser 45 gac Asp ttt Phe aaa Lys	cac His 30 ggc Gly ctg Leu gga Gly cac His	gtt Val cag Gln gag Glu ggt Gly aac Asn 95	gcc Ala tac Tyr agg Arg ttt Phe 80 ttg Leu	144 192 240

aac Asn	acc Thr 130	ctc Leu	att Ile	tgt Cys	ctt Leu	gtg Val 135	gac Asp	aac Asn	atc Ile	ttt Phe	cct Pro 140	cct Pro	.gtg Val	gtc Val	aac Asn	432
atc Ile 145	aca Thr	tgg Trp	ctg Leu	agc Ser	aat Asn 150	ggg Gly	cag Gln	tca Ser	gtc Val	aca Thr 155	gaa Glu	ggt Gly	gtt Val	tct Ser	gag Glu 160	480
acc Thr	agc Ser	ttc Phe	ctc Leu	tcc Ser 165	aag Lys	agt Ser	gat Asp	cat His	tcc Ser 170	ttc Phe	ttc Phe	aag Lys	atc Ile	agt Ser 175	tac Tyr	528
Leu	acc Thr	ttc Phe	ctc Leu 180	cct Pro	tct Ser	gct Ala	gat Asp	gag Glu 185	att Ile	tat Tyr	gac Asp	tgc Cys	aag Lys 190	gtg Val	gag Glu	576
cac His	tgg Trp	ggc Gly 195	ctg Leu	gac Asp	cag Gln	cct Pro	ctt Leu 200	ctg Leu	aaa Lys	cac His	tgg Trp	gag Glu 205	cct Pro	gag Glu	att Ile	624
cca Pro	gcc Ala 210	cct Pro	atg Met	tca Ser	gag Glu	ctc Leu 215	aca Thr	gag Glu	act Thr	gtg Val	gtc Val 220	Cys	gcc Ala	ctg Leu	GJ À aaa	672
ttg Leu 225	tct Ser	gtg Val	ggc Gly	ctc Leu	atg Met 230	Gly	att Ile	gtg Val	gtg Val	ggc Gly 235	act Thr	gtc Val	ttc Phe	atc Ile	atc Ile 240	720
caa Gln	ggc Gly	ctg Leu	Arg	tca Ser 245	gtt Val	ggt Gly	gct Ala	tcc Ser	aga Arg 250	cac His	caa Gln	GJ A aaa	cca Pro	ttg Leu 255		765
tgaa	tccc	at c	ctgg	aagg	g aa	ggtg	cato	gcc	atct	aca	ggag	caga	ag a	.atgg	acttg	825
ctaa	atga	cc t	agca	ctat	t ct	ctgg	cccg	att	tatc	ata	tccc	tttt	ct c	ctcc	aaata	885
tttc	tcct	ct c	acct	tttc	t ct	ggga	ctta	agc	tgct	ata	tccc	ctca	ga g	ctca	caaat	945
gcct	ttac	at t	cttt	ccct	g ac	ctcc	tgat	ttt	tttt	ttc	tttt	ctca	aa t	gtta	cctac	1005
aata	catg	cc t	3 <b>3</b> 33	taag	с са	cccg	gcta	cct	aatt	cct	cagt	aacc	tc c	atct	aaaat	1065
ctcc	aagga	aa go	caat	aaat	t cc	tttt	atga	g								1096
~21 N		<b>5</b>													•	

<210> 68

<211> 255

<212> PRT

<213> NM\_002122 MHC class II DQ alpha 1, HLA-DQA1

<400> 68

Met Ile Leu Asn Lys Ala Leu Leu Leu Gly Ala Leu Ala Leu Thr Thr 1 10 15

Val Met Ser Pro Cys Gly Gly Glu Asp Ile Val Ala Asp His Val Ala 20 25 30

Ser Cys Gly Val Asn Leu Tyr Gln Phe Tyr Gly Pro Ser Gly Gln Tyr 35 40 45

Thr His Glu Phe Asp Gly Asp Glu Gln Phe Tyr Val Asp Leu Glu Arg 50 55 60

Lys Glu Thr Ala Trp Arg Trp Pro Glu Phe Ser Lys Phe Gly Gly Phe 65 70 75 80

Asp Pro Gln Gly Ala Leu Arg Asn Met Ala Val Ala Lys His Asn Leu 85 90 95

Asn Ile Met Ile Lys Arg Tyr Asn Ser Thr Ala Ala Thr Asn Glu Val 100 105 110

Pro Glu Val Thr Val Phe Ser Lys Ser Pro Val Thr Leu Gly Gln Pro 115 120 125

Asn Thr Leu Ile Cys Leu Val Asp Asn Ile Phe Pro Pro Val Val Asn 130 135 140

Ile Thr Trp Leu Ser Asn Gly Gln Ser Val Thr Glu Gly Val Ser Glu 145 155 160

Thr Ser Phe Leu Ser Lys Ser Asp His Ser Phe Phe Lys Ile Ser Tyr 165 170 175

Leu Thr Phe Leu Pro Ser Ala Asp Glu Ile Tyr Asp Cys Lys Val Glu 180 . 190

His Trp Gly Leu Asp Gln Pro Leu Leu Lys His Trp Glu Pro Glu Ile
195 200 205

Pro Ala Pro Met Ser Glu Leu Thr Glu Thr Val Val Cys Ala Leu Gly 210 225 220

Leu Ser Val Gly Leu Met Gly Ile Val Val Gly Thr Val Phe Ile Ile 225 235 235 240

Gln Gly Leu Arg Ser Val Gly Ala Ser Arg His Gln Gly Pro Leu 245 250 255

<210> 69

<211> 2820

<212> DNA

<213> NM\_003014 SFRP4, secreted frizzled-related protein 4

<220>

<221> CDS

<222> (238)..(1275)

<223>

	.ggg.	69 ttcg	cgc	cccg	aag (	gctga	agag	ct g	gege	tgct	c gt	gccc	tgtg	tgc	cagacgg	60
cgg	gagc	tccg	cgg	ccgg	acc o	cgc	ggcc	cc go	ttt	gctg	c cg	actg	gagt	ttg	ggggaag	120
aaa	actc	tcct	gcg	cccc	aga a	agati	ttct	tc ct	cgg	cgaa	g gg	acag	cgaa	aga	tgagggt	180
ggo	cagga	aaga	gaag	ggcg	ett i	ctgt	ctg	cc gg	gggto	egca	g cg	cgag	aggg	cag	tgcc	237
ato Met 1	tto Phe	c cto	tce Se	c ato r Ile 5	c cta	a gto 1 Val	g gcg	g cto a Leu	tgo Cys 10	cto	g tg ı Tr	g cto	g cad u His	c cto s Lev 15	g gcg n Ala	285
cto Lev	Gly	gto Val	Arg 20	g Gl7 c ggc	gcg Ala	p ccc	c tgo Cys	gaç Glu 25	gcq Ala	g gto val	g cgo	c ato	c cct Pro 30	ate Me	g tgc t Cys	333
cgg Arg	cac His	ato Met 35	r ccc	tgg Trp	g aac Asr	ato Ile	acç Thi	g egg : Arg	atç Met	cco Pro	aac Ası	c cac n His 45	c cto s Lev	g cad 1 His	cac His	381
agc Ser	acg Thr 50	cag Gln	gag Glu	aac Asn	gec Ala	ato Ile 55	cto Lev	gcc Ala	ato Ile	gag Glu	caç Glr 60	tao Tyi	gaç Glu	gaq Glu	g ctg 1 Leu	429
gtg Val 65	gac	gtg Val	aac Asn	tgc Cys	ago Ser 70	gcc	gtg Val	ctg Leu	cgc	ttc Phe 75	tto Phe	tto Phe	tgt Cys	gco Ala	atg Met 80	477
tac Tyr	gcg Ala	ccc Pro	att Ile	tgc Cys 85	acc Thr	ctg Leu	gag Glu	ttc Phe	ctg Leu 90	cac	gac	cct	ato Ile	aaç Lys 95	ccg Pro	525
tgc Cys	aag Lys	tcg Ser	gtg Val 100	Cys	caa Gln	cgc Arg	gcg Ala	cgc Arg 105	gac Asp	gac Asp	tgc Cys	gag Glu	ccc Pro 110	Leu	atg Met	573
aag Lys	atg Met	tac Tyr 115	aac Asn	cac His	agc Ser	tgg Trp	ccc Pro 120	gaa Glu	agc Ser	ctg Leu	gcc Ala	tgc Cys 125	Asp	gag Glu	ctg Leu	621
cct Pro	gtc Val 130	tat Tyr	gac Asp	cgt Arg	ggc Gly	gtg Val 135	tgc Cys	att Ile	tcg Ser	cct Pro	gaa Glu 140	gcc Ala	atc Ile	gtc Val	acg Thr	669
gac Asp 145	Leu	ccg Pro	Glu	gat Asp	Val	Lys	tgg Trp	ata Ile	gac Asp	atc Ile 155	aca Thr	cca Pro	gac Asp	atg Met	atg Met 160	717
gta Val	cag Gln	gaa Glu	agg Arg	cct Pro 165	ctt Leu	gat Asp	gtt Val	gac Asp	tgt Cys 170	aaa Lys	cgc Arg	cta Leu	agc Ser	ccc Pro 175	gat Asp	765
cgg Arg	tgc Cys	aag Lys	tgt Cys 180	aaa Lys	aag Lys	gtg Val	aag Lys	cca Pro 185	act Thr	ttg Leu	gca Ala	acg Thr	tat Tyr 190	ctc Leu	agc Ser	813
aaa Lys	aac Asn	tac Tyr 195	agc Ser	tat Tyr	gtt Val	att Ile	cat His 200	gcc Ala	aaa Lys	ata Ile	aaa Lys	gct Ala 205	gtg Val	cag Gln	agg Arg	861
agt	ggc	tgc	aat	gag	gtc	aca	acg	gtg	gtg	gat	gta	aaa	gag	atc	ttc.	909

Ser	Gly 210	Cys	Asn	Glu	Val	Thr 215	Thr	Val	Val	Asp	Val 220	Lys	Glu	Ile	Phe	
aag Lys 225	tcc Ser	tca Ser	tca Ser	ccc Pro	atc Ile 230	cct Pro	cga Arg	act Thr	caa Gln	gtc Val 235	ccg Pro	ctc Leu	att Ile	aca Thr	aat Asn 240	957
tct Ser	tct Ser	tgc Cys	cag Gln	tgt Cys 245	cca Pro	cac His	atc Ile	ctg Leu	ccc Pro 250	cat His	caa Gln	gat Asp	gtt Val	ctc Leu 255	atc Ile	1005
				tgg Trp												1053
gtt Val	gaa Glu	aaa Lys 275	tgg Trp	aga Arg	gat Asp	cag Gln	ctt Leu 280	agt Ser	aaa Lys	aga Arg	tcc Ser	ata Ile 285	cag Gln	tgg Trp	gaa Glu	1101
gag Glu	agg Arg 290	ctg Leu	cag Gln	gaa Glu	cag Gln	cgg Arg 295	aga Arg	aca Thr	gtt Val	cag Gln	gac Asp 300	aag Lys	aag Lys	aaa Lys	aca Thr	1149
gcc Ala 305	G1 y ggg	cgc Arg	acc Thr	agt Ser	cgt Arg 310	agt Ser	aat Asn	ccc Pro	ccc Pro	aaa Lys 315	cca Pro	aag Lys	gga Gly	aag Lys	cct Pro 320	1197
cct Pro	gct Ala	ccc Prò	aaa Lys	cca Pro 325	gcc Ala	agt Ser	ccc Pro	aag Lys	aag Lys 330	aac Asn	att Ile	aaa Lys	act Thr	agg Arg 335	agt Ser	1245
gcc Ala	cag Gln	aag Lys	aga Arg 340	aca Thr	aac Asn	ccg Pro	aaa Lys	aga Arg 345	gtg Val	tgaç	gctaa	ict a	igttt	ccaa	ıa	1295
gcgg	agac	tt c	cgac	ttcc	t ta	cagg	atga	ggc	tggg	cat	tgcc	tggg	ac a	gcct	atgta	1355
aggc	catg	tg c	ccct	tgcc	c ta	acaa	ctca	ctg	cagt	gct	cttc	atag	ac a	cato	ttgca	1415
gcat	tttt	ct t	aagg	ctat	g ct	tcag	tttt	tct	ttgt	aag	ccat	caca	ag c	cata	gtggt	1475
aggt	ttgc	cc t	ttgg	taca	g aa	ggtg	agtt	aaa	gctg	gtg	gaaa	aggc	tt a	ttgc	attgc	1535
attc	agag	ta a	cctg	tgtg	c-at	actc	taga	aga	gtag	gga	aaat	aatg	ct t	gtta	caatt	1595
cgac	ctaa	ta t	gtgc	attg	t aa	aata	aatg	cca	tatt	tca	aaca	aaac	ac g	taat	tttt	1655
taca	gtat	gt t	ttat	tacc	t tt	tgat	atct	gtt	gttg	caa	tgtt	agtg	at g	tttt	aaaat	1715
gtga	tgaa	aa t	ataa	tgtti	t tt	aaga	agga	aca	gtag	tgg	aatg	aatg	tt a	aaag	atctt	1775
tatg	tgtt	ta t	ggtc	tgca	g aa	ggat	tttt	gtg	atga	aaġ	ggga	tttt	tt g	aaaa	attag	1835
agaa	gtag	ca t	atgg	aaaat	: ta	taat	gtgt	ttt	ttta	cca	atga	cttc	ag t	ttct	gtttt	1895
tagc	taga	aa c	ttaa	aaaca	aa	aata	ataa	taa	agaa	aaa	taaa	taaa	aa g	gaga	ggcag	1955
acaa	tgtc	tg g	attc	ctgtt	tt.	ttgg	ttac	ctg	attt	cca	tgat	catg	at g	cttc	ttgtc	2015
aaca	ccct	et t	aagc	agcad	ca	gaaa	cagt	gag	tttg	tct	gtac	catt	ag g	agtt	aggta	2075
ctaai	ttagi	tt g	gcta	atgct	. ca	agtai	ttt	ata	ccca	caa	gaga	ggtai	tg t	cact	catct	2135
tacti	tecca	ag g	acat	ccacc	ct	gagaa	ataa	ttt	gacaa	agc	ttaa	aaatq	gg c	cttca	atgtg	2195
agtgo	ccaaa	at ti	ttgti	ttttc	tto	catt	caaa	tati	tttci	ttt	gccta	aaata	ac a	tgtga	agagg	2255
agtta	aaata	at a	aatgi	tacag	aga	aggaa	agt	tgag	gttco	cac (	ctct	gaaat	tg a	gaati	tactt	2315

gacagttggg	atactttaat	cagaaaaaaa	gaacttattt	gcagcatttt	atcaacaaat	2375
ttcataattg	tggacaattg	gaggcattta	ttttaaaaaa	caattttatt	ggccttttgc	2435
taacacagta	agcatgtatt	ttataaggca	ttcaataaat	gcacaacgcc	caaaggaaat	2495
aaaatcctat	ctaatcctac	tctccactac	acagaggtaa	tcactattag	tattttggca	2555
tattattctc	caggtgtttg	cttatgcact	tataaaatga	tttgaacaaa	taaaactagg	2615
aacctgtata	catgtgtttc	ataacctgcc	tcctttgctt	ggccctttat	tgagataagt	2675
tttcctgtca	agaaagcaga	aaccatctca	tttctaacag	ctgtgttata	ttccatagta	2735
tgcattactc	aacaaactgt	tgtgctattg	gatacttagg	tggtttcttc	actgacaata	2795
ctgaataaac	atctcaccgg	aattc	.•	•		2820

<210> 70

<211> 346

<212> PRT

<213> NM\_003014 SFRP4, secreted frizzled-related protein 4

<400> 70

Met Phe Leu Ser Ile Leu Val Ala Leu Cys Leu Trp Leu His Leu Ala 1 5 10 15

Leu Gly Val Arg Gly Ala Pro Cys Glu Ala Val Arg Ile Pro Met Cys
20 25 30

Arg His Met Pro Trp Asn Ile Thr Arg Met Pro Asn His Leu His His 35 40 45

Ser Thr Gln Glu Asn Ala Ile Leu Ala Ile Glu Gln Tyr Glu Glu Leu 50 55 60

Val Asp Val Asn Cys Ser Ala Val Leu Arg Phe Phe Phe Cys Ala Met 65 70 75 80

Tyr Ala Pro Ile Cys Thr Leu Glu Phe Leu His Asp Pro Ile Lys Pro 85 90 95

Cys Lys Ser Val Cys Gln Arg Ala Arg Asp Asp Cys Glu Pro Leu Met 100 105 110

Lys Met Tyr Asn His Ser Trp Pro Glu Ser Leu Ala Cys Asp Glu Leu 115 120 125

Pro Val Tyr Asp Arg Gly Val Cys Ile Ser Pro Glu Ala Ile Val Thr 130 135 140

Asp Leu Pro Glu Asp Val Lys Trp Ile Asp Ile Thr Pro Asp Met Met 145 150 155 160

Val Gln Glu Arg Pro Leu Asp Val Asp Cys Lys Arg Leu Ser Pro Asp 165 170 175

Arg Cys Lys Cys Lys Lys Val Lys Pro Thr Leu Ala Thr Tyr Leu Ser 180 185

Lys Asn Tyr Ser Tyr Val Ile His Ala Lys Ile Lys Ala Val Gln Arg

Ser Gly Cys Asn Glu Val Thr Thr Val Val Asp Val Lys Glu Ile Phe 210 215 220

Lys Ser Ser Ser Pro Ile Pro Arg Thr Gln Val Pro Leu Ile Thr Asn 225 230 230 235

Ser Ser Cys Gln Cys Pro His Ile Leu Pro His Gln Asp Val Leu Ile 245 250 255

Met Cys Tyr Glu Trp Arg Ser Arg Met Met Leu Leu Glu Asn Cys Leu 260 265 . 270

Val Glu Lys Trp Arg Asp Gln Leu Ser Lys Arg Ser Ile Gln Trp Glu 275 280 285

Glu Arg Leu Gln Glu Gln Arg Arg Thr Val Gln Asp Lys Lys Thr 290 295 300

Ala Gly Arg Thr Ser Arg Ser Asn Pro Pro Lys Pro Lys Gly Lys Pro 305 310 315 320

Pro Ala Pro Lys Pro Ala Ser Pro Lys Lys Asn Ile Lys Thr Arg Ser 325 330 335

Ala Gln Lys Arg Thr Asn Pro Lys Arg Val 340 345

<210> 71

<211> 1362

<212> DNA

<213> NM\_004039 annexin A2

<220>

<221> CDS

<222> (50)..(1066)

<223>

	_									•						
<40 cat		71 ggga	cgct	ctca	agc t	ctco	gege	ca co	gcc	caget	t tc	ette	1		tct ac Ser Tl	58
gtt Val	cac His	gaa Glu	ato Ile	cto Lei	g tgo ı Cys	aaç Lys 10	g cto Let	ago Ser	tto Lev	g gaç ı Glı	g ggt 1 Gly 15	gat Asp	cac His	tci Sei	aca Thr	106
ccc Pro 20	cca Pro	agt Ser	gca Ala	tat Tyr	ggg Gly 25	tct Ser	gto Val	aaa Lys	gco Ala	tat Tyr 30	act Thi	aac Asr	ttt Phe	gat Asp	gct Ala 35	154
gag Glu	cgg Arg	gat Asp	gct	ttg Lev 40	aac Asn	att	gaa Glu	aca Thr	gcc Ala 45	ato Ile	aaç Lys	aco Thi	aaa Lys	ggt Gly 50	gtg Val	202
gat Asp	gag Glu	gtc Val	acc Thr 55	att Ile	gtc Val	aac Asn	att Ile	ttg Leu 60	acc Thr	aac Asn	cgc Arg	ago Ser	aat Asn 65	gca Ala	cag Gln	250
aga Arg	cag Gln	gat Asp 70	att Ile	gcc Ala	ttc Phe	gcc Ala	tac Tyr 75	cag Gln	aga Arg	agg Arg	acc Thr	aaa Lys 80	aag Lys	gaa Glu	ctt Leu	298
gca Ala	tca Ser 85	gca Ala	ctg Leu	aag Lys	tca Ser	gcc Ala 90	tta Leu	tct Ser	ggc G1y	cac His	ctg Leu 95	gag Glu	acg Thr	gtg Val	att	346
ttg Leu 100	ggc	cta Leu	ttg Leu	aag Lys	aca Thr 105	cct Pro	gct Ala	cag Gln	tat Tyr	gac Asp 110	Ala	tct Ser	gag Glu	cta Leu	aaa Lys 115	394
gct Ala	tcc Ser	atg Met	aag Lys	ggg Gly 120	ctg Leu	gga Gly	acc Thr	gac Asp	gag Glu 125	gac Asp	tct Ser	ctc Leu	att Ile	gag Glu 130	atc Ile	442
atc Ile	tgc Cys	tcc Ser	aga Arg 135	acc Thr	aac Asn	cag Gln	gag Glu	ctg Leu 140	cag Gln	gaa Glu	att Ile	aac Asn	aga Arg 145	gtc Val	tac Tyr	490
aag Lys	gaa Glu	atg Met 150	tac Tyr	aag Lys	act Thr	gat Asp	ctg Leu 155	gag Glu	aag Lys	gac Asp	att Ile	att Ile 160	tcg Ser	gac Asp	aca Thr	538
tct Ser	ggt Gly 165	gac Asp	ttc Phe	cgc Arg	aag Lys	ctg Leu 170	atg Met	gtt Val	gcc Ala	ctg Leu	gca Ala 175	aag Lys	ggt Gly	aga Arg	aga Arg	586
gca Ala 180	gag Glu	gat Asp	ggc Gly	tct Ser	gtc Val 185	att Ile	gat Asp	tat Tyr	gaa Glu	ctg Leu 190	att Ile	gac Asp	caa Gln	gat Asp	gct Ala 195	634
cgg Arg	gat Asp	ctc Leu	tat Tyr	gac Asp 200	gct Ala	gga Gly	gtg Val	aag Lys	agg Arg 205	aaa Lys	gga Gly	act Thr	gat Asp	gtt Val 210	ccc Pro	682
aag Lys	tgg Trp	atc Ile	agc Ser 215	atc Ile	atg Met	acc Thr	gag Glu	cgg Arg 220	agc Ser	gtg Val	ccc Pro	cac His	ctc Leu 225	cag Gln	aaa Lys	730
gta Val	Phe	gat Asp 230	agg Arg	tac Tyr	aag Lys	Ser	tac Tyr 235	agc Ser	cct Pro	tat Tyr	gac Asp	atg Met 240	ttg Leu	gaa Glu	agc Ser	778

	atc Ile	agg Arg 245	aaa Lys	gag Glu	gtt Val	aaa Lys	gga Gly 250	gac Asp	ctg Leu	gaa Glu	aat Asn	gct Ala 255	ttc Phe	ctg Leu	aac Asn	ctg Leu	826
	gtt Val 260	cag Gln	tgc Cys	att Ile	cag Gln	aac Asn 265	aag Lys	ccc Pro	ctg Leu	tat Tyr	ttt Phe 270	gct Ala	gat Asp	cgg Arg	ctg Leu	tat Tyr 275	874
	gac Asp	tcc Ser	atg Met	aag Lys	ggc Gly 280	aag Lys	Gly ggg	acg Thr	cga Arg	gat Asp 285	aag Lys	gtc Val	ctg Leu	atc Ile	aga Arg 290	atc Ile	<b>922</b>
,	atg Met	gtc Val	tcc Ser	cgc Arg 295	agt Ser	gaa Glu	gtg Val	gac Asp	atg Met 300	Leu	aaa Lys	att Ile	agg Arg	tct Ser 305	gaa Glu	ttc Phe	970
	aag Lys	aga Arg	aag Lys 310	tac Tyr	ggc Gly	aag Lys	tcc Ser	ctg Leu 315	tac Tyr	tat Tyr	tat Tyr	atc Ile	cag Gln 320	caa Gln	gac Asp	act Thr	1018
	тАз	ggc Gly 325	gac Asp	tac Tyr	cag Gln	aaa Lys	gcg Ala 330	ctg Leu	ctg Leu	tac Tyr	Leu	tgt Cys 335	ggt Gly	gga Gly	gat Asp	gac Asp	1066
	tgaa	gccc	ga c	acgg	cctg	a gc	gtcc	agaa	atg	gtgc	tca	ccat	gctt	сс а	gcta	acagg	1126
	tcta	gaaa	ac c	agct	tgcg	a at	aaca	gtcc	ccg	tggc	cat	ccct	gtga	gg g	tgac	gttag	1186
	catt	accc	сс а	acct	catt	t ta	gttg	ccta	agc	attg	cct	ggcc	ttcc	tg t	ctag	tctct	1246
	cctg	taag	сс а	aaga	aatg	a ac	attc	caag	gag	tṭgg	aag	tgaa	gtct	at g	atgt	gaaac	1306
	actt	tgcc	tc c	tgtg	tact	g tg	tcat	aaac	aga	tgaa	taa .	actg	aatt	tg t	actt	t	1362
	<210	> 7	2														
	<211	> 3	39														
	<212	> P	RT														

<400> 72

<213> NM\_004039 annexin A2

Met Ser Thr Val His Glu Ile Leu Cys Lys Leu Ser Leu Glu Gly Asp 1 5 15

His Ser Thr Pro Pro Ser Ala Tyr Gly Ser Val Lys Ala Tyr Thr Asn 20 25 30

Phe Asp Ala Glu Arg Asp Ala Leu Asn Ile Glu Thr Ala Ile Lys Thr 35 40 45

Lys Gly Val Asp Glu Val Thr Ile Val Asn Ile Leu Thr Asn Arg Ser 50 55 60

Asn Ala Gln Arg Gln Asp Ile Ala Phe Ala Tyr Gln Arg Arg Thr Lys 65 70 75 80

Lys Glu Leu Ala Ser Ala Leu Lys Ser Ala Leu Ser Gly His Leu Glu  $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$ 

Thr Val Ile Leu Gly Leu Leu Lys Thr Pro Ala Gln Tyr Asp Ala Ser 100 105 110

Glu Leu Lys Ala Ser Met Lys Gly Leu Gly Thr Asp Glu Asp Ser Leu 115 120 125

Ile Glu Ile Ile Cys Ser Arg Thr Asn Gln Glu Leu Gln Glu Ile Asn 130 135 140

Arg Val Tyr Lys Glu Met Tyr Lys Thr Asp Leu Glu Lys Asp Ile Ile 145  $\phantom{\bigg|}150\phantom{\bigg|}155\phantom{\bigg|}$  155

Ser Asp Thr Ser Gly Asp Phe Arg Lys Leu Met Val Ala Leu Ala Lys 165 170 175

Gly Arg Arg Ala Glu Asp Gly Ser Val Ile Asp Tyr Glu Leu Ile Asp 180 185 190

Gln Asp Ala Arg Asp Leu Tyr Asp Ala Gly Val Lys Arg Lys Gly Thr
195 200 205

Asp Val Pro Lys Trp Ile Ser Ile Met Thr Glu Arg Ser Val Pro His 210 215 220

Leu Gln Lys Val Phe Asp Arg Tyr Lys Ser Tyr Ser Pro Tyr Asp Met 225 230 235 240

Leu Glu Ser Ile Arg Lys Glu Val Lys Gly Asp Leu Glu Asn Ala Phe 245 250 255

Leu Asn Leu Val Gln Cys Ile Gln Asn Lys Pro Leu Tyr Phe Ala Asp 260 265 270

Arg Leu Tyr Asp Ser Met Lys Gly Lys Gly Thr Arg Asp Lys Val Leu 275 280 285

Ile Arg Ile Met Val Ser Arg Ser Glu Val Asp Met Leu Lys Ile Arg 290 295 300

Ser Glu Phe Lys Arg Lys Tyr Gly Lys Ser Leu Tyr Tyr Tyr Ile Gln 305 310 315 320

Gln Asp Thr Lys Gly Asp Tyr Gln Lys Ala Leu Leu Tyr Leu Cys Gly 325 330 335

Gly Asp Asp

<21	.0>	73														
<21	.1>	850														
<21	2>	DNA				-		٠		•		•				
<21	3>	NM_(	00395	55 SC	ocs3						•					
<22	0>													•		
<22	1>	CDS				٠										
<22	2>	(107	7) (	(781)												
<22	3>								•							
<40 gcg		73 .cct	ctcc	gcag	icc c	cccg	rggat	g cg	gtaç	ıcggc	egc	tgtç	lcāā	aggo	cgcga	ıa 60
gca	gctg	cag	ccgc	egec	ge g	ıcaga	itcca	ıc go	tggc	tccg	tgc	gcc		gtc Val		115
cac His	agc Ser 5	aag Lys	ttt Phe	ccc Pro	gco Ala	gcc Ala 10	ggg	atg Met	ago Ser	cgc Arg	ccc Pro	ctġ Leu	gac Asp	acc Thr	agc Ser	163
ctg Leu 20	cgc Arg	cto	aag Lys	acc Thr	tto Phe 25	agc Ser	tcc Ser	aag Lys	ago	gag Glu	tac Tyr	cag Gln	ctg Lev	gtg Val	gtg Val 35	211
aac Asn	gca Ala	gtg Val	cgc	aag Lys 40	ctg Leu	cag Gln	gag Glu	agc Ser	ggc Gly 45	ttc Phe	tac Tyr	tgg Trp	ago Ser	gca Ala 50	gtg Val	259
acc Thr	Gly	ggc	gag Glu 55	gcg Ala	aac Asn	ctg Leu	ctg Leu	ctc Leu 60	agt Ser	gcc Ala	gag Glu	ccc Pro	gcc Ala 65	ggc	acc	307
ttt Phe	ctg Leu	atc Ile 70	cgc Arg	gac Asp	agc Ser	tcg Ser	gac Asp 75	cag Gln	cgc Arg	cac His	ttc Phe	ttc Phe 80	acg Thr	ctc Leu	agc Ser	355
gtc Val	aag Lys 85	acc Thr	cag Gln	tct Ser	GJ A āāā	acc Thr 90	aag Lys	aac Asn	ctg Leu	cgc Arg	atc Ile 95	cag Gln	tgt Cys	gag Glu	ggg Gly	403
ggc Gly 100	agc Ser	ttc Phe	tct Ser	ctg Leu	cag Gln 105	agc Ser	gat Asp	ccc Pro	cgg Arg	agc Ser 110	a¢g Thr	cag Gln	ccc Pro	gtg Val	ccc Pro 115	451
cgc Arg	ttc Phe	gac Asp	tgc Cys	gtg Val 120	ctc Leu	aag Lys	ctg Leu	gtg Val	tac Tyr 125	cac His	tac Tyr	atg Met	ccg Pro	ccc Pro 130	cct Pro	499
gga Gly	gcc Ala	ccc Pro	tcc Ser 135	ttc Phe	ccc Pro	tcg Ser	cca Pro	cct Pro 140	Thr	gaa Glu	ccc Pro	tcc Ser	tcc Ser 145	gag Glu	gtg Val	547
ccc Pro	gag Glu	cag Gln 150	ccg Pro	tct Ser	gcc Ala	cag Gln	cca Pro 155	ctc Leu	cct Pro	GJ A GGG	agt Ser	ccc Pro 160	ccc Pro	aga Arg	aga Arg	595
gcc	tat	tac	atc	tac	tcc	ggg	ggc	gag	aag	atc	ccc	ctg	gtg	ttg	agc	643

Ala Tyr Tyr Ile Tyr Ser Gly Gly Glu Lys Ile Pro Leu Val Leu Ser 165 170 175	
cgg ccc ctc tcc tcc aac gtg gcc act ctt cag cat ctc tgt cgg aag Arg Pro Leu Ser Ser Asn Val Ala Thr Leu Gln His Leu Cys Arg Lys 180 185 190 195	691
acc gtc aac ggc cac ctg gac tcc tat gag aaa gtc acc cag ctg ccg Thr Val Asn Gly His Leu Asp Ser Tyr Glu Lys Val Thr Gln Leu Pro 200 205 210	739
ggg ccc att cgg gag ttc ctg gac cag tac gat gcc ccg ctt Gly Pro Ile Arg Glu Phe Leu Asp Gln Tyr Asp Ala Pro Leu 215 220 225	781
taaggggtaa agggcgcaaa gggcatgggt cgggagaggg gacgcaggcc cctctcctcc	841
gtggcacat	850
<210> 74	•
<211> 225	
<212> PRT	
<213> NM_003955 SOCS3	
<400> 74	
Met Val Thr His Ser Lys Phe Pro Ala Ala Gly Met Ser Arg Pro Leu  1 10 15	
Asp Thr Ser Leu Arg Leu Lys Thr Phe Ser Ser Lys Ser Glu Tyr Gln 20 25 30	
Leu Val Val Asn Ala Val Arg Lys Leu Gln Glu Ser Gly Phe Tyr Trp 35 40 45	
Ser Ala Val Thr Gly Gly Glu Ala Asn Leu Leu Leu Ser Ala Glu Pro 50 55 60	
Ala Gly Thr Phe Leu Ile Arg Asp Ser Ser Asp Gln Arg His Phe Phe 65 70 75 80	
Thr Leu Ser Val Lys Thr Gln Ser Gly Thr Lys Asn Leu Arg Ile Gln 85 90 95	
Cys Glu Gly Gly Ser Phe Ser Leu Gln Ser Asp Pro Arg Ser Thr Gln 100 105 110	
Pro Val Pro Arg Phe Asp Cys Val Leu Lys Leu Val Tyr His Tyr Met 115 120 125	
Pro Pro Pro Gly Ala Pro Ser Phe Pro Ser Pro Pro Thr Glu Pro Ser 130 135 140	

PCT/AU2003/001166

203 Ser Glu Val Pro Glu Gln Pro Ser Ala Gln Pro Leu Pro Gly Ser Pro 145 150 Pro Arg Arg Ala Tyr Tyr Ile Tyr Ser Gly Gly Glu Lys Ile Pro Leu 165 Val Leu Ser Arg Pro Leu Ser Ser Asn Val Ala Thr Leu Gln His Leu 185 Cys Arg Lys Thr Val Asn Gly His Leu Asp Ser Tyr Glu Lys Val Thr 200 205 Gln Leu Pro Gly Pro Ile Arg Glu Phe Leu Asp Gln Tyr Asp Ala Pro 215 Leu 225 <210> 75 <211> 369 <212> DNA <213> NM\_000331 SAA1, serum amyloid A1 <220> <221> CDS <222> (1)..(366) <223> <400> 75 atg aag ctt ctc acg ggc ctg gtt ttc tgc tcc ttg gtc ctg ggt gtc 48 Met Lys Leu Leu Thr Gly Leu Val Phe Cys Ser Leu Val Leu Gly Val 10 age age ega age the the teg the ett gge gag get the gat ggg get 96 Ser Ser Arg Ser Phe Phe Ser Phe Leu Gly Glu Ala Phe Asp Gly Ala 25 cgg gac atg tgg aga gcc tac tct gac atg aga gaa gcc aat tac atc 144 Arg Asp Met Trp Arg Ala Tyr Ser Asp Met Arg Glu Ala Asn Tyr Ile ggc tca gac aaa tac ttc cat gct cgg ggg aac tat gat gct gcc aaa 192 Gly Ser Asp Lys Tyr Phe His Ala Arg Gly Asn Tyr Asp Ala Ala Lys 55 agg gga cet ggg ggt gte tgg get gea gaa geg ate age gat gee aga 240 Arg Gly Pro Gly Gly Val Trp Ala Ala Glu Ala Ile Ser Asp Ala Arg 70

gag aat atc cag aga ttc ttt ggc cat ggt gcg gag gac tcg ctg gct

Glu Asn Ile Gln Arg Phe Phe Gly His Gly Ala Glu Asp Ser Leu Ala

85

288

gat cag gct gcc aat gaa tgg ggc agg agt ggc aaa gac ccc aat cac 336 Asp Gln Ala Ala Asn Glu Trp Gly Arg Ser Gly Lys Asp Pro Asn His 100 105 110

ttc cga cct gct ggc ctg cct gag aaa tac tga Phe Arg Pro Ala Gly Leu Pro Glu Lys Tyr 115 120

369

<210> 76 ·

<211> 122

<212> PRT

<213> NM\_000331 SAA1, serum amyloid A1

<400> 76

Ser Ser Arg Ser Phe Phe Ser Phe Leu Gly Glu Ala Phe Asp Gly Ala 20 25 30

Arg Asp Met Trp Arg Ala Tyr Ser Asp Met Arg Glu Ala Asn Tyr Ile 35 40 45

Gly Ser Asp Lys Tyr Phe His Ala Arg Gly Asn Tyr Asp Ala Ala Lys 50 60

Arg Gly Pro Gly Gly Val Trp Ala Ala Glu Ala Ile Ser Asp Ala Arg 65 70 75 80

Glu Asn Ile Gln Arg Phe Phe Gly His Gly Ala Glu Asp Ser Leu Ala 85 90 95

Asp Gln Ala Ala Asn Glu Trp Gly Arg Ser Gly Lys Asp Pro Asn His 100 105 110

Phe Arg Pro Ala Gly Leu Pro Glu Lys Tyr 115 120

<210> 77

<211> 895

<212> DNA

<213> NM\_014059 RGC32

<220>

<221> CDS

<222> (147)..(497)

<223>

<40 gcg		77 cgt	cgac	cggc	gc g	gctg	gago	g ca	gcgc	:cgaa	ggg	actg	gca	gggc	tgaagt	60
gtg	cggg	aca	gcaa	gccc	cc g	aata	gccc	c gg	ctgc	cacc	tcg	cagg	acc	caag	gccacg	120
'cgc	gccg	ggc	ccag	ctga	gc c	gcct	c at Me 1	g aa t Ly	g cc s Pr	g cc o Pr	c gc o Al 5	g ga a Gl	g ga u As	c ct p Le	g tcg u Ser	173
gac Asp 10	gcg Ala	ctg Leu	tgc Cys	gag Glu	ttt Phe 15	gac Asp	gcg Ala	gtg Val	ctg Leu	gcc Ala 20	gac Asp	ttc Phe	gcg Ala	tcg Ser	ccc Pro 25	221
ttc Phe	cac His	gag Glu	cgc Arg	cac His 30	ttc Phe	cac His	tac Tyr	gag Glu	gag Glu 35	cac	ctg Leu	gag Glu	cgc Arg	atg Met 40	aag Lys	269
cgg Arg	cgc Arg	agc Ser	agc Ser 45	gcc Ala	agt Ser	gtc Val	agc Ser	gac Asp 50	agc Ser	agc Ser	ggc Gly	ttc Phe	agc Ser 55	gac Asp	tcg Ser	317
gag Glu	agt Ser	gca Ala 60	gat Asp	tca Ser	ctt Leu	tat Tyr	agg Arg 65	aac Asn	agc Ser	ttc Phe	agc Ser	ttc Phe 70	agt Ser	gat Asp	gaa Glu	365
aaa Lys	ctg Leu 75	aat Asn	tct Ser	cca Pro	aca Thr	gac Asp 80	tct Ser	acc Thr	cca Pro	gct Ala	ctt Leu 85	ctc Leu	tct Ser	gcc Ala	act Thr	413
gtc Val 90	act Thr	cct Pro	cag Gln	aaa Lys	gct Ala 95	aaa Lys	tta Leu	gga Gly	gac Asp	aca Thr 100	aaa Lys	gag Glu	cta Leu	gaa Glu	gcc Ala 105	461
ttc Phe	att Ile	gct Ala	gat Asp	ctt Leu 110	gac Asp	aaa Lys	act Thr	tta Leu	gca Ala 115	agt Ser	atg Met	tgaa	acaa	aga		507
agtt	ctgg	gt c	cttt	cato	a ta	aggg	agaa	gct	tcag	aaa	gttc	cgag	ga c	ctgo	taaaa	567
tcag	ctac	ta g	aatc	tgct	g cc	agag	ggga	caa	agac	gtg	cact	caac	ct t	ctac	caggc	627
cact	ctca	gg c	tcac	ctta	a aa	tcag	ccct	tga	tece	att	tctg	ggca	at t	taga	cagtg	687
aaac	tgac	tt t	gttt	acct	g ct	tgca	gcat	att	agaa	cag	acga	tcca	tg c	taat	attgt	747
attt	tctc	tt a	aaac	atag	c tt	tect	gtaa	ttt	aaag	tgc	tttt	atga	aa a	tatt	tgtaa	807
ttaa	ttat	at a	tagt <sup>.</sup>	tgga	a ata	agca	gtaa	gct	ttcc	cat	tata	atat	at t	tttg	tatac	867
aaat	aaaa	tt t	gaact	tgaa	c ct	cgtg	cc									895

<210> 78

<211> 117

<212> PRT

<213> NM\_014059 RGC32

<400>. 78

Met Lys Pro Pro Ala Glu Asp Leu Ser Asp Ala Leu Cys Glu Phe Asp 1 5 10 15

Ala Val Leu Ala Asp Phe Ala Ser Pro Phe His Glu Arg His Phe His 20 25 30

Tyr Glu Glu His Leu Glu Arg Met Lys Arg Arg Ser Ser Ala Ser Val 35 40 45

Ser Asp Ser Ser Gly Phe Ser Asp Ser Glu Ser Ala Asp Ser Leu Tyr 50 60

Arg Asn Ser Phe Ser Phe Ser Asp Glu Lys Leu Asn Ser Pro Thr Asp 65 70 75 80

Ser Thr Pro Ala Leu Leu Ser Ala Thr Val Thr Pro Gln Lys Ala Lys 85 90 95

Leu Gly Asp Thr Lys Glu Leu Glu Ala Phe Ile Ala Asp Leu Asp Lys 100 105 110

Thr Leu Ala Ser Met 115

<210> 79

<211> 1564

<212> DNA

<213> NM\_018004 FLJ10134

<220>

<221> CDS

<222> (314)..(1138)

<223>

<400> 79

gaagcacate tggacagetg tgcggcetec ttgcgggccg acgtcagccg agcacgtcce 60
ccacgtcete teettetege cacttattat ttattegttt teecaaagaa gegactaggg 120
acceaagttt aaaaatteet eeeeecaete aatgcgagae gtggecagat eeeatecaae 180
acacggttta atttcatgg ggetetggga teaaaagaac agaaacagca acaacaaaag 240
cceagceget gtetgattt aagetggcaa agtgggaaaa ataaagtgtt gagtaaacag 300
accaagttgg atc atg ggg aat tte aga ggt cat gee etc eet gga acc 349
Met Gly Asn Phe Arg Gly His Ala Leu Pro Gly Thr

ti Pl	tc t ne P	ne i	ttt Phe 15	att Ile	at Il	t g e G	gt d ly I	ett Seu	tgq Trp 20	g tg o Tr	g t	gt 'ys	aca Thi	a aa r Ly	ıg ad rs Sd	er	att Ile	ct Le	g aa u Ly	ag /s	397
ta T <u>y</u>	at a yr I. 3	re (	gc Cys	aaa Lys	l aa Ly:	g ca s Gi	ln I	ag ys 5	cga Arg	a ac	c t	gc gc	tai Tyi	ct Le 40	u G	gt Ly	tcc Ser	aa. Ly	a ao s Tì	a ır	445
tt Le 45	a ti eu Pl	tc t ne 1	at Yyr	cga Arg	Le	g ga u G] 50	.u 1	tt le	ttg Leu	ga Gl	g g u G	ga ly	att Ile 55	ac Th	a at r Il	ta (	gtt Val	gg Gl	c at y Me	t	493
gc Al	t ti a Le	a a u T	ct hr	Gly ggc	ato Met 65	g go E Al	t g a G	gg 1y	gag Glu	ca Gl:	g t n Pi 7	he	att Ile	cc Pr	t gg o Gl	y (	ej Saa	ecc Pro 75	c ca	t s	541
ct Le	gat uMe	g t	eu	tat Tyr 80	gac	ta Ty	t a r L	aa ys	caa Gln	gg G1: 85	у Н:	ac is	tgg Trp	aat Ası	t ca n Gl	n I	etc Seu 90	cto	gg Gl	c y	589
tg: Trj	g ca p Hi	t c s H 9	13	ttc Phe	acc Thr	at Me	g t	yr .	ttc Phe 100	tto Phe	ti Pl	tt	eja aaa	cto Lev	tt Le 10	u G	gt 1y	gtg Val	gc.	a a	637
ga Ası	t at p Il 11	е п	ta (	tgt Cys	ttc Phe	ac Th	E 1.	ic a le s	igt Ser	tca Ser	ct Le	t eu	cct Pro	gtg Val 120	. Se	c t	ta	acc Thr	aa Ly:	3	685
tta Lei 125	a at 1 Me 5	g ti t Le	tg t eu S	tca Ser	aat Asn	gc: Ala 130	a re	a t eu E	tt he	gtg Val	ga Gl	u i	gcc Ala 135	ttt	ate Ile	c t e P	tc he	tac Tyr	aad Asr 140	1	733
cac His	ac Th:	t ca r Hi	st o	ътĀ	cgg Arg 145	gaa Glu	a at 1 Me	g c	tg eu	gac Asp	at 11 15	e I	ttt Phe	gtg Val	cac His	C C	ln	ctg Leu 155	cto	Ī I	781
gtt Val	Lei	g gt 1 Va	T A	al 60	ttt Phe	cto	ac Th	a g r G	тА	ctc Leu 165	gt Va	t c	jcc Na	ttc Phe	cta	ı G.	ag lu 70	ttc Phe	ctt Leu		829
gtt Val	Arg	aa As 17	11 M	at sn	gta Val	ctt Leu	ct Le	u G	ag lu: 80	cta Leu	Lei	g c	gg	tca Ser	agt Ser 185	Le	eu :	att Ile	ctg Leu		877
ctt Leu	cag Gln 190	GI	ga yS	gc i	tgg Irp	ttc Phe	tt Pho 19	e G.	ag a	att Ile	G13	a t y P	he	gtc Val 200	ctg Leu	ta Ty	it d	ccc Pro	ccc Pro		925
agt Ser 205	gga Gly	gg G1	t co	ct o	та	tgg Trp 210	gai Asp	ct Le	ga eu M	atg Met	gat Asp	С Н - 2	is (	gaa Glu	aat Asn	at Il	t t e I	eu	ttt Phe 220		97.3
ctc Leu	acc Thr	ata Ile	a to ∋ Cy	ys E	tt he 25	tgt Cys	tgg Trp	r ca Hi	t t s I	'yr	gca Ala 230	Va	ta a	acc Thr	att Ile	gt Va	1 1	tc le 35	gtt Val		1021
gga Gly	atg Met	aat Asr	ta Ty 24	'E A	ct i	ttc Phe	att Ile	ac Th	r T	gg rp 45	ttg Leu	gt Va	tt a al I	aaa ys	tct Ser	ag Ar 25	g L	tt a eu 1	aag Lys		1069
agg Arg	ctc Leu	tgc Cys 255	Se	c t	ca d	gaa Glu	gtt Val	gg G1 26	λr	tt eu :	ctg Leu	a <i>a</i> Ly	na a /s A	sn i	gct Ala 265	gaa Glu	a c	ga q rg (	gaa Slu		1117
caá Gln	gaa Glu 270	tca Ser	ga Gl	ag uG	aa g lu G	ilu	atg Met 275	tg	act	ttga	at q	gag	ctt	cca	g tt	ttt	ct	aga			1168

taaacctttt cttttttaca ttgttcttgg ttttgtttct cgatcttttg tttggagaac	1228
agctggctaa ggatgactct aagtgtactg tttgcatttc caatttggtt aaagtatttg	1288
aatttaaata ttttcttttt agctttgaaa atattttggg tgatactttc attttgcaca	1348
tcatgcacat catggtattc aggggctaga gtgatttttt tccagattat ctaaagttgg	1408
atgcccacac tatgaaagaa atatttgttt tatttgcctt atagatatgc tcaaggttac	1468
tgggettget actatttgta acteettgae catggaatta tacttgttta tettgttget	1528
gcaatgagaa ataaatgaat gtatgtattt tggtgc	1564
<210> 80	
<211> 275	
<212> PRT	
<213> NM_018004 FLJ10134	
<400> 80	
Met Gly Asn Phe Arg Gly His Ala Leu Pro Gly Thr Phe Phe Ile	•
1 5 10 15	
Ile Gly Leu Trp Trp Cys Thr Lys Ser Ile Leu Lys Tyr Ile Cys Lys	
20 25 30	
Lys Gln Lys Arg Thr Cys Tyr Leu Gly Ser Lys Thr Leu Phe Tyr Arg	
40 45	
Leu Glu Ile Leu Glu Gly Ile Thr Ile Val Gly Met Ala Leu Thr Gly 50 55	
55 60	
Met Ala Gly Glu Gln Phe Ile Pro Gly Gly Pro His Leu Met Leu Tyr 65 70 75 80	
75 80	
Asp Tyr Lys Gln Gly His Trp Asn Gln Leu Leu Gly Trp His His Phe 85 90 95	
Thr Met Tyr Phe Phe Phe Gly Leu Leu Gly Val Ala Asp Ile Leu Cys 100 105 110	
Phe Thr Ile Ser Ser Leu Pro Val Ser Leu Thr Lys Leu Met Leu Ser 115 120 125	
Asn Ala Leu Phe Val Glu Ala Phe Ile Phe Tyr Asn His Thr His Gly 130 135 140	
·	
Arg Glu Met Leu Asp Ile Phe Val His Gln Leu Leu Val Leu Val Val	

150 155 160

												•							
	Ph	e	Let	ı Ti	ır G	Sly	Leu 165	Val	. Ala	Phe	Leu	1 Glu 170	Phe	Leu	Val	Arg	Asn 175	Asn	
	Va	1	Lev	ı Le	eu G 1	31u .80	Leu	Leu	Arg	Ser	Ser 185	Leu	Ile	Leu	Leu	Gln 190	Gly	Ser	
	Tr	p	Phe	Ph 19	e G 5	ln	Ile	Gly	Phe	Val 200	Leu	Tyr	Pro	Pro	Ser 205	Gly	Gly	Pro	
	Al	a	Trp 210	As	рL	eu i	Met	Asp	His 215	Glu	Asn	Ile	Leu	Phe 220	Leu	Thr	Ile	Cys	
	Ph 22	e :	Cys	Tr	р Н.	is '	Tyr	Ala 230	Val	Thr	Ile	Val	'Ile 235	Val	Gly	Met	Asn	Tyr 240	
	Ala	a :	Phe	11	e Ti	hr !	Trp 245	Leu	Val	Lys	Ser	Arg- 250	Leu	Lys	Arg	Leu	Cys 255	Ser	
	Sei	r (	Glu	۷a	l G1 26	ly 1 50	Leu	Leu	Lys	Asn	Ala 265	Glu	Arg	Glu	Gln	Glu 270	Ser	Glu	
	Glu	1 (	Slu	Met 275															
•	<21	.0>	٠ ٤	31															
	<21	1>	. 2	311				-											
	<21	2>		NA															
	<21	3>	N	M_0	040	04	GJB:	2, c	onne:	xin :	26								
	<22	0>												,					
	<22	1>	С	DS							٠								
	<22	2>		199	)	(876	6)						•						
	<223	3>																	,
	<400 gatt		8: aato	-	ato	jaca	aac	taa	gttg	gtt	ctgt	cttc	ac c	tgtt:	ttgg	t ga	ggtt	gtgt	60
ä	aaga	ıgt	tgg	jt c	rttt	get	cag	gaa	gaga	ttt	aago	atgo	tt g	ctta	ccca	g ac	tcag	agaa	120
•	gtct	cc	cto	jt t	ctg	rtco	tag	cta	tgtt	cct	gtgt	tgtg	tg c	attc	gtct	t tt	ccag	agca	180
ē	acc	gc	cca	ıg a	gta	gaa	ga M	tg g et A	at t sp T	gg g rp G	gc a ly T 5	hr L	tg c eu G	ag a ln T	cg a hr I	tc c le L	eu G	ly gg	231
G	gt ly	gt Va	ga 1A	sn	aaa Lys 15	ca Hi	c to s Se	ec a	cc achr Se	gc a er 1: 20	le G	ga a ly L	ag a ys I	tc to	gg ci cp Le 25	eu Tl	ec gi nr Va	c al	279
С	tc	tt	c a	tt	ttt	cg	c at	t at	g at	c ct	c g	tt g	tģ go	ct go	ca aa	ıg ga	ıg gt	g	327

Leu Phe Ile Phe Arg Ile Met Ile Leu Val Val Ala Ala Lys Glu Val 30 35 40	
tgg gga gat gag cag gcc gac ttt gtc tgc aac acc ctg cag cca ggc Trp Gly Asp Glu Gln Ala Asp Phe Val Cys Asn Thr Leu Gln Pro Gly 45 50 55	375
tgc aag aac gtg tgc tac gat cac tac ttc ccc atc tcc cac atc cgg Cys Lys Asn Val Cys Tyr Asp His Tyr Phe Pro Ile Ser His Ile Arg 60 65 70 75	423
cta tgg gcc ctg cag ctg atc ttc gtg tcc agc cca gcg ctc cta gtg Leu Trp Ala Leu Gln Leu Ile Phe Val Ser Ser Pro Ala Leu Leu Val 80 85 90	471
gcc atg cac gtg gcc tac cgg aga cat gag aag aag agg aag ttc atc Ala Met His Val Ala Tyr Arg Arg His Glu Lys Lys Arg Lys Phe Ile 95 100 105	519
aag ggg gag ata aag agt gaa ttt aag gac atc gag gag atc aaa acc Lys Gly Glu Ile Lys Ser Glu Phe Lys Asp Ile Glu Glu Ile Lys Thr 110 115 120	567
cag aag gtc cgc atc gaa ggc tcc ctg tgg tgg acc tac aca agc agc Gln Lys Val Arg Ile Glu Gly Ser Leu Trp Trp Thr Tyr Thr Ser Ser 125 130 135	615
atc ttc ttc cgg gtc atc ttc gaa gcc gcc ttc atg tac gtc ttc tat Ile Phe Phe Arg Val Ile Phe Glu Ala Ala Phe Met Tyr Val Phe Tyr 140 145 150 155	663
gtc atg tac gac ggc ttc tcc atg cag cgg ctg gtg aag tgc aac gcc Val Met Tyr Asp Gly Phe Ser Met Gln Arg Leu Val Lys Cys Asn Ala 160 165 170	711
tgg cct tgt ccc aac act gtg gac tgc ttt gtg tcc cgg ccc acg gag Trp Pro Cys Pro Asn Thr Val Asp Cys Phe Val Ser Arg Pro Thr Glu 175 180 185	759
aag act gtc ttc aca gtg ttc atg att gca gtg tct gga att tgc atc Lys Thr Val Phe Thr Val Phe Met Ile Ala Val Ser Gly Ile Cys Ile 190 195 200	807
ctg ctg aat gtc act gaa ttg tgt tat ttg cta att aga tat tgt tct Leu Leu Asn Val Thr Glu Leu Cys Tyr Leu Leu Ile Arg Tyr Cys Ser 205 210 215	855
ggg aag tca aaa aag cca gtt taacgcattg cccagttgtt agattaagaa Gly Lys Ser Lys Lys Pro Val 220 225	906
atagacagca tgagagggat gaggcaaccc gtgetcaget gtcaaggete agtegecage	966
atttcccaac acaaagatte tgacettaaa tgcaaccatt tgaaaccect gtaggeetca	1026
ggtgaaactc cagatgccac aatgagctct gctcccctaa agcctcaaaa caaaggccta	1086
attetatgee tgtettaatt ttettteact taagttagtt ceaetgagae eeeaggetgt	1146
taggggttat tggtgtaagg tactttcata ttttaaacag aggatatcgg catttgtttc	1206
tttctctgag gacaagagaa aaaagccagg ttccacagag gacacagaga aggtttgggt	1266
gtcctcctgg ggttcttttt gccaactttc cccacgttaa aggtgaacat tggttctttc	1326
atttgctttg gaagttttaa tctctaacag tggacaaagt taccagtgcc ttaaactctg	1386
ttacactttt tggaagtgaa aactttgtag tatgataggt tattttgatg taaagatgtt	1446

ctggatacca	ttatatgttc	cccctgtttc	agaggctcag	attgtaatat	gtaaatggta	1506
tgtcattcgc	tactatgatt	taatttgaaa	tatggtcttt	tggttatgaa	tactttgcag	1566
cacagctgag	agaggctgtc	tgttgtattc	attgtggtca	tagcacctaa	caacattgta	1626
gcctcaatcg	agtgagacag	actagaagtt	cctagttggc	ttatgatagc	aaatggcctc	1686
atgtcaaata	ttagatgtaa	ttttgtgtaa	gaaatacaga	ctggatgtac	caccaactac	1746
tacctgtaat	gacaggcctg	tccaacacat	ctcccttttc	catgctgtgg	tagccagcat	1806
cggaaagaac	gctgatttaa	agaggtgagc	ttgggaattt	tattgacaca	gtaccattta	1866
atggggagac	aaaaatgggg	gccaggggag	ggagaagttt	ctgtcgttaa	aaacgagttt	1926
ggaaagactg	gactctaaat	tctgttgatt	aaagatgagc	tttgtctacc	ttcaaaagtt	1986
tgtttggctt	acccccttca	gcctccaatt	ttttaagtga	aaatataact	aataacatgt	2046
gaaaagaata	gaagctaagg	tttagataaa	tattgagcag	atctatagga	agattgaacc .	2106
tgaatattgc	cattatgctt	gacatggttt	ccaaaaaatg	gtactccaca	tacttcagtg	2166
agggtaagta	ttttcctgtt	gtcaagaata	gcattgtaaa	agcattttgt	aataataaag	2226
atagcttta	atgatatgct	tgtaactaaa	ataattttgt	aatgtatcaa	atacatttaa	2286
acattaaaa	tataatctct	ataat				2311

<210> 82

<211> 226

<212> PRT

<213> NM\_004004 GJB2, connexin 26

<400> 82

Met Asp Trp Gly Thr Leu Gln Thr Ile Leu Gly Gly Val Asn Lys His 1  $\phantom{\Big|}$  10  $\phantom{\Big|}$  15

Ser Thr Ser Ile Gly Lys Ile Trp Leu Thr Val Leu Phe Ile Phe Arg 20 25 30

Ile Met Ile Leu Val Val Ala Ala Lys Glu Val Trp Gly Asp Glu Gln 35 40 45

Ala Asp Phe Val Cys Asn Thr Leu Gln Pro Gly Cys Lys Asn Val Cys 50 55 60

Tyr Asp His Tyr Phe Pro Ile Ser His Ile Arg Leu Trp Ala Leu Gln 65 70 75 80

Leu Ile Phe Val Ser Ser Pro Ala Leu Leu Val Ala Met His Val Ala 85 90 95

Tyr	Arg	Arg	His 100	Glu	Lys	Lys	Arg	Lys 105	Phe	Ile	ьўs	Gly	Glu 110		Гуз	
Ser (	Glu	Phe 115	Lys	Asp	Ile	Glu	Glu 120	Ile	Lys	Thr	Gln	Lys 125	Val	Arg	Ile	
Glu (	31y 130	Ser	Leu	Trp	Trp	Thr 135	Tyr	Thr	Ser	Ser	Ile 140	Phe	Phe	Arg	Val	
Ile I 145	?he	Glu	Ala	Ala	Phe 150	Met	Tyr	Val	Phe	Tyr 155	Val	Met	Tyr	Asp	Gly 160	
Phe S	Ser	Met	Gln	Arg 165	Leu	Val	Lys	Суз	Asn 170	Ala	Trp	Pro	Cys	Pro 175	Asn	
Thr V	al.	Asp	Cys 180	Phe	Val	Ser	Arg	Pro 185	Thr	Glu	Lys	Thr	Val 190	Phe	Thr	
Val P	he l	Met 195	Ile	Ala	Val	Ser	Gly 200	Ile	Суз	Ile	Leu	Leu 205	Àsn	Val	Thr	
Glu L 2	eu ( 10	Cys :	Tyr	Leu	Leu	Ile . 215	Arg	Tyr	Cys	Ser	Gly 220	Lys	Ser	Lys	Lys	
Pro Va 225	al				•	•										
<210>	83	}														
<211>	23	89														
<212>	DN	A								•						
<213>	NM	_002	514	NOV	l, ne	phro	blas	toma	a ove	erexp	press	sed g	jene			
<220>																
<221>	CD	s						•			•					
<222>	(7	3)	(114	3)												
<223>								. ,								
<400> 83 gggaaggcga gcagtgccaa tctacagcga agaaagtctc gtttggtaaa agcgagaggg 66									60							
gaaagc	ctga	ı gc	atg Met 1	cag Gln	agt Ser	gtg Val	cag Gln 5	agc Ser	acg Thr	agc Ser	ttt Phe	tgt Cys 10	ctc Leu	cga Arg	aag Lys	111
cag tgo Gln Cys 15	c ct 3 Le	t to u Cy	rs Le	eu Th	ec to	ne Le	g ci	tt ci	tc ca eu Hi	at ci is Le	eu Le	tg gg eu Gl	ga ca Ly Gi	ag gi In Va	tc al	159

gct gcg act cag cgc tgc cct ccc cag tgc ccg ggc cgg tgc cct gcg

Ala 30	a Ala	Thi	Gli	n Ar	g Cys 35	Pro	Pro	Glr	о Суз	9ro	Gly	Arg	J Cys	s Pro	Ala 45	
acg Thr	r CCC	p ccg	g aco	Cys 50	c gcc s Ala	ccc Pro	GI7	g gto Val	g cgc Arg 55	gcg Ala	g gto Val	Let	g gad Asp	60 60	tgc / Cys	255
tca Ser	tgo Cys	tgt Cys	Cto Let 65	g gto 1 Val	g tgt L Cys	gco Ala	cgc Arg	Glr Glr 70	g cgt n Arg	Gly	gag Glu	g ago	tgo Cys 75	c tca Ser	gat Asp	303
ctg Leu	gag Glu	Pro 80	tgo Cys	gao Asp	gag Glu	ago Ser	agt Ser 85	ggc Gly	cto Leu	tac Tyr	tgt Cys	gat Asp 90	cgc Arg	ago g Ser	gcg Ala	351
gac Asp	Pro 95	ago Ser	aac Asr	cag Glr	g act	ggc Gly 100	Ile	tgc Cys	acg Thr	gcg	gta Val 105	Glu	gga Gly	gat Asp	aac Asn	399
tgt Cys 110	Val	ttc Phe	gat Asp	Gly ggg	y gto Val 115	Ile	tac Tyr	cgc	agt Ser	gga Gly 120	Glu	aaa Lys	ttt Phe	cag Gln	cca Pro 125	447
agc Ser	tgc Cys	aaa Lys	ttc Phe	Gln 130	Суз	acc	tgc Cys	aga Arg	gat Asp 135	GJ À aàà	cag Gln	att	Gly	tgt Cys 140	gtg Val	495
ccc Pro	ÿrg	tgt Cys	cag Gln 145	Leu	gat Asp	gtg Val	cta Leu	ctg Leu 150	Pro	gag Glu	· cct Pro	aac	tgc Cys 155	Pro	gct Ala	543
cca Pro	aga Arg	aaa Lys 160	Val	gag Glu	gtg Val	cct Pro	gga Gly 165	gag Glu	tgc Cys	tgt Cys	gaa Glu	aag Lys 170	Trp	atc Ile	tgt Cys	591
ggc	cca Pro 175	gat Asp	gag Glu	gag Glu	gat Asp	tca Ser 180	ctg Leu	gga Gly	ggc Gly	ctt Leu	acc Thr 185	ctt Leu	gca Ala	gct Ala	tac Tyr	639
agg Arg 190	cca Pro	gaa Glu	gcc Ala	acc Thr	cta Leu 195	gga Gly	gta Val	gaa Glu	gtc Val	tct Ser 200	gac Asp	tca Ser	agt Ser	gtc Val	aac Asn 205	687
tgc Cys	att Ile	gaa Glu	cag Gln	acc Thr 210	aca Thr	gag Glu	tgg Trp	aca Thr	gca Ala 215	tgc Cys	tcc Ser	aag Lys	agc Ser	tgt Cys 220	ggt Gly	735
atg Met	Gly ggg	ttc Phe	tcc Ser 225	acc Thr	cgg Arg	gtc Val	acc Thr	aat Asn 230	agg Arg	aac Asn	cgt Arg	caa Gln	tgt Cys 235	gag Glu	atg Met	783
ctg Leu	aaa Lys	cag Gln 240	act Thr	cgg Arg	ctc Leu	tgc Cys	atg Met 245	gtg Val	cgg Arg	ccc Pro	tgt Cys	gaa Glu 250	caa Gln	gag Glu	cca Pro	831
gag Glu	cag Gln 255	cca Pro	aca Thr	gat Asp	aag Lys	aaa Lys 260	gga Gly	aaa Lys	aag Lys	tgt Cys	ctc Leu 265	cgc Arg	acc Thr	aag Lys	aag Lys	879
tca Ser 270	ctc Leu	aaa Lys	gcc Ala	atc Ile	cac His 275	ctg Leu	cag Gln	ttc Phe	aag Lys	aac Asn 280	tgc Cys	acc Thr	agc Ser	ctg Leu	cac His 285	927
acc Thr	tac Tyr	aag Lys	ccc Pro	agg Arg 290	ttc Phe	tgt Cys	G1y ggg	gtc Val	tgc Cys 295	agt Ser	gat Asp	ggc Gly	cgc Arg	tgc Cys 300	tgc Cys	975
act	ccc	cac	aat	acc	aaa	acc	atc	cag	gca	gag	ttt	cag	tgc	tcc	cca	1023

Thr Pro His Asn Thr Lys Thr Ile Gln Ala Glu Phe Gln Cys Ser Pro 305 310 315	
ggg caa ata gtc aag aag cca gtg atg gtc att ggg acc tgc acc tgt Gly Gln Ile Val Lys Lys Pro Val Met Val Ile Gly Thr Cys Thr Cys 320 325 330	1071
cac acc aac tgt cct aag aac aat gag gcc ttc ctc cag gag ctg gag His Thr Asn Cys Pro Lys Asn Asn Glu Ala Phe Leu Gln Glu Leu Glu 335 340 345	1119
ctg aag act acc aga ggg aaa atg taacctatca ctcaagaagc acacctacag Leu Lys Thr Thr Arg Gly Lys Met 350 355	1173
agcacctgta gctgctgcgc cacccaccat caaaggaata taagaaaagt aatgaagaat	1233
cacgatttca teettgaate etatgtattt teetaatgtg ateatatgag gacettteat	1293
atctgtcttt tatttaacaa aaaatgtaat taactgtaaa cttggaatca aggtaagctc	1353
aggatatggc ttaggaatga cttactttcc tgtggtttta ttacaaatgc aaatttctat	1413
aaatttaaga aaacaagtat ataatttact ttgtagactg tttcacattg cactcatcat	1473
attttgttgt gcactagtgc aattccaaga aaatatcact gtaatgagtc agtgaagtct	1533
agaatcatac ttaacatttc attgtacaag tattacaacc atatattgag gttcattggg	1593
aagattetet attggeteee tttttgggta aaccagetet gaactteeaa geteeaaate	1653
caaggaaaca tgcagctctt caacatgaca tccagagatg actattactt ttctgtttag	1713
ttttacacta ggaaacgtgt tgtatctaca gtaatgaaat gtttactaag tggactggtg	1773
tcataaactt tctccattta agacacattg actcctttcc aatagaaaga aactaaacag	1833
aaaactccca atacaaagat gactggtccc tcatagccct cagacattta tatattggaa	1893
gctgctgagg cccccaagtt ttttaattaa gcagaaacag catattagca gggattctct	1953
catctaactg atgagtaaac tgaggcccaa agcacttgct tacatcctct gatagctgtt	2013
tcaaaigtgc attttgtgga attttgagaa aaatagagca aaatcaacat gactggtggt	2073
gagagaccac acattttatg agagtttgga attattgtag acatgcccaa aacttatcct	2133
tgggccataa ttatgaaaac tcatgatcaa gatatatgtg tatacataca tgtatctggt	2193
ttgtcaggct acaaggtagg ctgcaaaatt aaatctagac attctttaa tgccaccaca	2253
cgtgttccgc ttctctcttt taaagtattt ataaaaatat aaattgtaca ttttgtaaaa	2313
tattatgttt gatttctcta cttgtcatat cactaaataa acacgatttt attgctgaaa	2373
aaaaaaaaa aaaaaa	2389

<210> 84

<211> 357

<212> PRT

<213> NM\_002514 NOV1, nephroblastoma overexpressed gene

<400> 84

Met Gln Ser Val Gln Ser Thr Ser Phe Cys Leu Arg Lys Gln Cys Leu 1 5 10 15

Cys Leu Thr Phe Leu Leu Leu His Leu Leu Gly Gln Val Ala Ala Thr 20 25 30 .

Gln Arg Cys Pro Pro Gln Cys Pro Gly Arg Cys Pro Ala Thr Pro Pro 35 40 45

Thr Cys Ala Pro Gly Val Arg Ala Val Leu Asp Gly Cys Ser Cys Cys 50 55 60

Leu Val Cys Ala Arg Gln Arg Gly Glu Ser Cys Ser Asp Leu Glu Pro 65 70 75 80

Cys Asp Glu Ser Ser Gly Leu Tyr Cys Asp Arg Ser Ala Asp Pro Ser 85 90 95

Asn Gln Thr Gly Ile Cys Thr Ala Val Glu Gly Asp Asn Cys Val Phe 100 105 110

Asp Gly Val Ile Tyr Arg Ser Gly Glu Lys Phe Gln Pro Ser Cys Lys 115 120 125

Phe Gln Cys Thr Cys Arg Asp Gly Gln Ile Gly Cys Val Pro Arg Cys 130 135 140

Gln Leu Asp Val Leu Leu Pro Glu Pro Asn Cys Pro Ala Pro Arg Lys 145 150 155 160

Val Glu Val Pro Gly Glu Cys Cys Glu Lys Trp Ile Cys Gly Pro Asp 165 170 175

Glu Glu Asp Ser Leu Gly Gly Leu Thr Leu Ala Ala Tyr Arg Pro Glu 180 185 190

Ala Thr Leu Gly Val Glu Val Ser Asp Ser Ser Val Asn Cys Ile Glu 195 200 205

Gln Thr Thr Glu Trp Thr Ala Cys Ser Lys Ser Cys Gly Met Gly Phe 210 215 220

Ser Thr Arg Val Thr Asn Arg Asn Arg Gln Cys Glu Met Leu Lys Gln 225 230 235 240

Thr Arg Leu Cys Met Val Arg Pro Cys Glu Gln Glu Pro Glu Gln Pro
245 250 255

Thr Asp Lys Lys Gly Lys Lys Cys Leu Arg Thr Lys Lys Ser Leu Lys 260 265 270

Ala Ile His Leu Gln Phe Lys Asn Cys Thr Ser Leu His Thr Tyr Lys 275 280 285

Pro Arg Phe Cys Gly Val Cys Ser Asp Gly Arg Cys Cys Thr Pro His 290 295 300

As n Thr Lys Thr Ile Gln Ala Glu Phe Gln Cys Ser Pro Gly Gln Ile 305 310 315 320

Val Lys Lys Pro Val Met Val Ile Gly Thr Cys Thr Cys His Thr Asn 325 330 335

Cys Pro Lys Asn Asn Glu Ala Phe Leu Gln Glu Leu Glu Leu Lys Thr 340 345 350

Thr Arg Gly Lys Met 355

International application No.

PCT/AU03/01166

Α.		CLASSIFICATION OF SUBJECT M	ATTE	R		
Int. Cl.	7;	C12Q 1/68, G01N 33/53				
Accord	ling to l	International Patent Classification (IPC)	or to be	oth national classification and IPC		
В.		FIELDS SEARCHED				
Minimu SEE B		mentation searched (classification system foll	owed b	y classification symbols)		
			to the	extent that such documents are included in the fields sea	rched	
SEE B		·		£ dat has a dayle and have and		
DGEN	E, EM	IBL, GENBANK: SEQ ID NOS 1-7,	9, 11,	of data base and, where practicable, search terms used) 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35,		
46, 48, diagno		2, 53, 55, 57, 59, 60- 63, 65, 67, 69, 7	1-75,	77, 79, 81, 83 WPIDS: ovarian, cancer, tumor	ır, detect,	
C.		DOCUMENTS CONSIDERED TO BE RE	LEVA	NT		
Categ	ory*	Citation of document, with indication,	where	appropriate, of the relevant passages	Relevant to claim No.	
X		Weitzel JN et al (1994) Gynecologichanges associated with ovarian car See table 3 and pages 249 and 50 w	ncer" :		1-5, 7, 9-12, 16-20, 22, 24- 27, 31-43, 62- 69	
×		DGENE Abstract Accession No AI SCIENCES, INC.) 3 January 2002 See abstract and claim 1, SEQ ID N				
	X Fi	urther documents are listed in the con	tinuat	tion of Box C X See patent family an	nex	
"A" o	document which is relevance earlier a after the	pplication or patent but published on or international filing date	"T"	later document published after the international filing and not in conflict with the application but cited to un or theory underlying the invention document of particular relevance; the claimed invention considered novel or cannot be considered to involve a when the document is taken alone	derstand the principle on cannot be in inventive step	
"L" document which may throw doubts on priority "Y" document of particular relevance; the claimed invention cannot be claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious a person skilled in the art					ment is combined	
reason (as specified)  "O" document referring to an oral disclosure, use,  "&" document member of the same patent family exhibition or other means						
"P" d	locumer	nt published prior to the international filing later than the priority date claimed	-	•	•	
Date of t	he actua	al completion of the international search		Date of mailing of the international search report	0.6 NOV 2003	
23 Octo		003 ng address of the ISA/AU		Authorized officer		
		PATENT OFFICE		remotized officer		
		/ODEN ACT 2606, AUSTRALIA pct@ipaustralia.gov.au		TERRY MOORE		
		02) 6285 3929		Telephone No : (02) 6283 2632		

PCT/AU03/01166

C (Continua	tion 1). DOCUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
х	Hough CD et al (2000) Cancer Research 60, 6281-7 "Large-scale serial analysis of gene expression reveals genes differentially expressed in ovarian cancer" See table 3: mal2, claudin 3(relevant to SEQ ID NOs 11, 15)	1-5, 7, 10-12 16-20, 22, 25 27, 31-43, 60
PX	WO 02 071928 A2 (MILLENNIUM PHARMACEUTICALS, INC) 19 September 200 See table 1 pg 20 KIAA0869: pg 19 FLJ22252: pg 21 SLP1, pgs 21, 24, 25 PAX8: pg 18 ANXA2 (applicant's SEQ ID NOs 13, 17, 25, 43 and 71/72) & DGENE Accession No ABS76418 (applicant's SEQ ID NO 17)	1
PX	WO 02 102235 A2 (EOS BIOTECHNOLOGY, INC) 27 December 2002 See pg 187 KIAA0869: pg 184 claudin 7: pgs 111, 162 and 180 KIAA0101: pg 120 KIAA1481, pgs 152, 174, 184, 202 paired box gene 8: pgs 155, 161 methylene tetrahydrofolate dehydrogenase: pg 176 DD5: pg 125 SOCS3: pg 261 serum amyloid A1 (applicant's SEQ ID NOs 13, 21, 23, 37, 43, 55, 63, 73/74 and 75)	1-5, 7, 10-12 16-20, 22, 25 27, 31-43, 45 47-53, 60-65
PX	DGENE Abstract Accession No ACA66405 & US 2003/0004102 A1 (ASHKENAZI AJ et al) 2 January 2003 See abstract and claim 2, figure 203 (applicant's SEQ ID NO 19)	1-5, 7, 10-12 16-20, 22, 25 27, 31-43, 60
x	Scheurle D et al "Cancer gene discovery using digital differential display" (2000)(online), (retrieved on 16 October 2003) Retrieved from Internet <url:http: cancergenes.htm="" cmbb="" publications="" www.fau.edu=""> See "up-regulation of known genes: claudin 7, secreted frizzled rel. pro. 4 (applicant's SEQ ID NOs 21 and 69)</url:http:>	1-5, 7, 10-12 16-20, 22, 25 27, 31-43, 45 47-53, 60-69
<b>X</b>	DGENE Abstract Accession No AAS56533 & WO 01 70796 A2 (CORIXA CORPORATION) 27 September 2001 See abstract and claim 1, page 165 KIAA0101 (applicant's SEQ ID NO 23)	1-5, 7, 10-12 16-20, 22, 25 27, 31-43, 60
x	Medline Abstract 11906548 Shigemasa K et al (2001) Int J Gynecol Cancer 11(6), 454-61 "Expression of the protease inhibitor antileukoprotease and the serine protease stratum corneum chymotryptic enzyme (SCCE) is coordinated in ovarian tumours" See abstract, relevant to applicant's SEQ ID NO 25	

International application No.
PCT/AU03/01166

	PCT/AU03/6	11100
C (Continua	tion 2) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	Laval S et al (1994) Cell Growth Differ 5(11), 1173-83 "Isolation and characterization of an epithelial-specific receptor tyrosine kinase from an ovarian cancer cell line"	
х	See whole document, relevant to applicant's SEQ ID NO 27	1-5, 7, 9-11, 16-20, 22, 25- 27, 31-42, 62- 69
	WO 03 068054 A2 (THE GOVERNMENT OF THE UNITED STATES OF AMERICA as represented by THE SECRETARY, DEPARTMENT OF HEALTH SERVICES) 21 August 2003	
PX	See in particular table 1, pg 22/60 EDDR1: pg 32/60 TOP2A: 29/60 PAX8 and 33/60 SFRP4 (applicant's SEQ ID NOs 27, 29, 43 and 69)	1-5, 7, 10-12, 16-20, 22, 25- 27, 31-43, 45, 47, 51-53, 62- 69
	Koshiyama M et al (2001) Anticancer Res 21, 2925-32 "Immunohistochemical expression of topoisomerase II alpha (TopII alpha) and multidrug resistance-associated protein (MRP), plus chemosensitivity testing, as chemotherapeutic indices of ovarian endometrial carcinomas"	
Х	See whole document, relevant to applicant's SEQ ID NO 29	1-5, 7, 10-12, 16-20, 22, 26, 27, 31-40, 42, 43, 62-69
X	Gotlieb WH et al (2001) Gynecol Oncol 82(1), 99-104 "Topoisomerase II immunostaining as a prognostic marker for survival in ovarian cancer" See whole document, relevant to applicant's SEQ ID NO 29	1-5, 7, 10-12, 16-20, 22, 26, 27, 31-40, 42, 43, 62-69
	Costa MJ et al (2000) Int J Gynecol Pathol 19(3), 248-57 "Topoisomerase II alpha: prognostic predictor and cell cycle marker in surface epithelial neoplasms of the ovary and peritoneum"	43, 02-09
х	See whole document, relevant to applicant's SEQ ID NO 29	1-5, 7, 10-12, 16-20, 22, 25- 27, 31-43, 62- 69
·	DGENE Abstract Accession No ABQ54317 & WO 02 00677 A1 (HUMAN GENOME SCIENCES, INC.) 3 January 2002	
X	See abstract and claim 1, SEQ ID NO 197 (applicant's SEQ ID NO 31 reverse complement)	1-5, 7, 10-12, 16-20, 22, 25- 27, 31-43, 60-
	DGENE Abstract Accession No ACA03948 & US 2002/0192678 A1 (CHEN H-M) 19 December 2002	-
PX	See abstract and example 13, pages 98-99 (applicant's SEQ ID NO 33)	1-5, 7, 10-12, 16-20, 22, 24- 27, 31-43, 62- 69

International application No. PCT/AU03/01166

16-20, 22, 25-27, 31-43, 60-

1-5, 7, 10-12, 16-20, 22, 25-27, 31-43, 60-

1-5, 7, 10-12, 15-20, 22, 25-27, 30-43, 60-

1-5, 7, 10-12, 15-20, 22, 25-27, 30-43, 60-

69

DOCUMENTS CONSIDERED TO BE RELEVANT C (Continuation 3) Relevant to Category\* Citation of document, with indication, where appropriate, of the relevant passages claim No. Medline Abstract 11925933 Sakamoto M et al (2001) Hum Cell 14(4), 305-15 "Analysis of gene expression profiles associated with cisplatin resistance in human ovarian cancer cell lines and tissues using cDNA microarray" Х 1-5, 7, 10-12, See whole document, relevant to applicant's SEQ ID NO 35 16-20, 22, 24-27, 31-43, 62-69 DGENE Abstract Accession No ABQ54876 & WO 02 00677 A1 (HUMAN GENOME SCIENCES, INC.) 3 January 2002 Х See abstract and claim 1, SEQ ID NO 756 (applicant's SEQ ID NO 39) 1-5, 7, 10-12, 16-20, 22, 25-27, 31-43, 60-69 DGENE Abstract Accession No ABL83226 & WO 01 92581 A2 (CORIXA CORPORATION) 6 December 2001 Х See abstract and claim 1, SEQ ID NO 6204 (applicant's SEQ ID NO 41) 1-5, 7, 10-12, 16-20, 22, 25-27, 31-43, 60-Genbank Accession No AK075046 3 September 2002 Isogai T et al Homo sapiens cDNA FLJ90565, clone OVARC1001336 X See "features" (applicant's SEQ ID NO 50) 1-5, 7, 16-20, 22, 31-40, 62-67 DGENE Abstract Accession No AAH83197 & WO 01 51513 A2 (CORIXA CORPORATION) 19 July 2001 Х See abstract and claim 5, SEQ ID NO 821 (applicant's SEQ ID NO 52) 1-5, 7, 10-12,

DGENE Abstract Accession No ABL90699 & WO 01 90304 A2 (HUMAN GENOME

See abstract and claim 4, SEQ ID NO 1261 (applicant's SEQ ID NO 53)

GESELLSCHAFT F R GENOMFORSHUNG) 21 October 1999

See abstract, claim 3 and page 226 (applicant's SEQ ID NO 57)

PHARMACEUTICALS) 13 December 2001

DGENE Abstract Accession No AAZ77553 & WO 99 53040 A2 (METAGEN

DGENE Abstract Accession No ABL67667 & WO 01 94629 A2 (AVALON

See abstract and claim 1, SEQ ID NO 6004 (applicant's SEQ ID NOs 59, 60)

SCIENCES) 29 November 2001

Х

X

X

International application No.
PCT/AU03/01166

C (Continua	tion 4) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	DGENE Abstract Accession No ABL61830 & WO 01 94629 A2 (AVALON PHARMACEUTICALS) 13 December 2001	
Х	See abstract and claim 1, SEQ ID NO 167 (applicant's SEQ ID NOs 61, 62)	1-5, 7, 10-12 15-20, 22, 24 27, 30-43, 62
	DGENE Abstract Accession No AAV64871 & WO 98 48010 A1 (GARVAN INSTITUTE OF MEDICAL RESEARCH) 29 October 1998	
х	See Abstract and figure 3B, example and pg 22 line 34 of WO 98 48010 (applicant's SEQ ID NO 63)	1-5, 7, 10-1: 16-20, 22, 2 27, 31-43, 4 47, 51-53, 6
	DGENE Abstract Accession No ABV34360 & WO 01 60860 A2 (MILLENIUM PREDICTIVE MEDICINE, INC) 23 August 2001	
<b>x</b> .	See Abstract, claim 1, pg 19 and pg 7222 (applicant's SEQ ID NO 65)	1-5, 7, 10-12 16-20, 22, 2 27, 31-43, 4 47, 51-53, 6
	DGENE Abstract Accession No ABQ54583 & WO 02 00677 A1 (HUMAN GENOMI SCIENCES, INC.) 3 January 2002	
х	See in particular abstract and claim 1, SEQ ID NO 463 (applicant's SEQ ID NO 67)	1-5, 7, 10-12 16-20, 22, 2 27, 31-43, 4 47, 51-53, 6
	DGENE Abstract Accession No ABL62031 & WO 01 94629 A2 (AVALON PHARMACEUTICALS) 13 December 2001	
Х	See abstract and claim 1, SEQ ID NO 368 (applicant's SEQ ID NOs 73, 74)	1-5, 7, 10-12 16-20, 22, 2 27, 31-43, 4 47-53, 60-69
	DGENE Abstract Accession No AAI99253 & WO 01 55313 A2 (HUMAN GENOME SCIENCES, INC) 2 August 2001	İ
X	See abstract, example 2 and SEQ ID NO 1017 (applicant's SEQ ID NO 77)	1.5, 7, 10-12 16-20, 22, 2 27, 31-43, 4 47, 51-53, 66
	DGENE Abstract Accession No ABL64081 & WO 01 94629 A2 (AVALON PHARMACEUTICALS) 13 December 2001	
х	See abstract and claim 1, SEQ ID NO 2418 (applicant's SEQ ID NO 79)	1-5, 7, 10-12 16-20, 22, 25 27, 31-43, 45 47-53, 60-69

International application No.
PCT/AU03/01166

	PC17AU03/0							
C (Continuation 5) DOCUMENTS CONSIDERED TO BE RELEVANT								
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.						
	DGENE Abstract Accession No ABL66499 & WO 01 94629 A2 (AVALON PHARMACEUTICALS) 13 December 2001							
Х	See abstract and claim 1, SEQ ID NO 4836 (applicant's SEQ ID NO 81)	1-5, 7, 10-12 16-20, 22, 24 27, 31-43, 45 47, 51-53, 60						
	·	69						
	· · · · · · · · · · · · · · · · · · ·							
	·							

International application No. PCT/AU03/01166

Box I	Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)
This inter	national search report has not been established in respect of certain claims under Article 17(2)(a) for the following
1.	Claims Nos:
	because they relate to subject matter not required to be searched by this Authority, namely:
2.	X Claims Nos: 70-73
	because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
	The claims are not limited to the technical features of the invention as disclosed in, and supported by, the specification. The claims relate to methods of assessing the promoter regions of specific genes with respect to hypermethylation and ovarian cancer. However the specification only discloses sequences derived from cDNA clones and does not disclose any promoter sequences.
3.	Claims Nos :
	because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)
Box II	Observations where unity of invention is lacking (Continuation of item 3 of first sheet)
This Inter	national Searching Authority found multiple inventions in this international application, as follows:
See si	upplementary sheet
	•
	•
1.	As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ·	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:2-5, 7, 9, 15, 19, 20, 22, 24, 30, 45 and 47-50 in full and claims 1, 10-12, 16-18, 25-27, 31-43, 51-53 and 60-69 as they relate to table 3.
4.	No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark o	on Protest The additional search fees were accompanied by the applicant's protest.
	X No protest accompanied the payment of additional search fees.

International application No.

PCT/AU03/01166

#### Supplemental Box

(To be used when the space in any of Boxes I to VIII is not sufficient)

#### Continuation of Box No: II

The application claims more than one invention. Rule 13.1 of the PCT states the principle that an International Application should relate to only one invention or, if there is more than one invention, that the inclusion of those inventions in one International Application is only permitted if all inventions are so linked as to form a single general inventive concept.

Rule 13.2 of the PCT defines the method for determining whether the requirement of unity of invention is satisfied in respect of a group of inventions claimed in an International application. Unity of invention exists only when there is a technical relationship among the claimed inventions involving one or more of the same or corresponding "special technical features." The expression "special technical features" is defined in Rule 13.2 as meaning those technical features that define a contribution which each of the inventions, considered as a whole, makes over the prior art. The determination is made on the contents of the claims as interpreted in light of the description and drawings (if any).

The claims and the description relate to methods for the diagnosis, prediction and treatment of ovarian cancer, using gene and peptide sequences whose expression is altered in ovarian cancer. In particular the claims define methods relating to the use of over 693 specific gene and/or peptide sequences.

Although all of the claims share the common feature that they all relate to methods or apparatus that involve the use of specific gene or peptide sequences whose expression is altered in ovarian cancer, this feature is known (see the documents listed below). As such this feature cannot be regarded as a special technical feature and cannot confer unity on the inventions relating to the use of specific gene or peptide sequences for the detection or treatment of ovarian cancer.

- D1 US 5 912 142
- D2 US 6 268 165
- D3 WO 01 21653
- D4 Bayani et al (2002) Cancer Research 62, 3466-76

Furthermore, there is nothing in the specification to suggest that the genes can be further divided into groups, where sequences within a group share a common special technical feature. Although the specification discloses that some of the genes can be classified into further, narrower groups, these further divisions also cannot be regarded as special technical features. These divisions correspond to groups of genes that are down-regulated or up-regulated in cancers and genes that are associated with specific types of ovarian cancers, such as epithelial or mucinous ovarian cancer. However these groups are known, as are methods of using genes specifically associated with these narrower groups for the treatment and detection of specific subsets of ovarian cancers. See in particular D4

Although there is a lack of unity the ISA has searched, as a service to the applicant, five sets of ten sequences for five search fees.

These sequences comprise SEQ ID NOS 1-7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 46, 48, 50, 52, 53, 55, 57, 59, 60-63, 65, 67, 69, 71-75, 77, 79, 81, 83.

Information on patent family members

International application No. PCT/AU03/01166

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

	Document Cited in Search Report			Pate	ant Family Member	•	
wo	0200677	AU	29508/01	AU	30958/01	AU	36459/01
		AU	36460/01	AU	36461/01	AU	36462/01
		AU	36463/01	AU	36464/01	AU	36465/01
		AU	36466/01	ΔU	37943/01	AU	37944/01
٠	,	AU	37947/01	UΑ	37949/01	AU	37950/01
		AU	37951/01	AU	37952/01	AU	37953/01
		AU	37954/01	ΑU	37955/01	AU	37957/01
		AU	37958/01	UΑ	38585/01	ΑU	39726/01
		AU	39727/01	AU	39728/01	AU	41402/01
		AU	41403/01	AU	41404/01	AU	41405/01
		AU	41406/01	AU	41407/01	AU	41408/01
		AU	41409/01	AU	41410/01	AU	41411/01
	•	AU	41412/01	AU	41413/01	AU	41414/01
		AU	41415/01	AU	41416/01	AU	41417/01
		AU	41418/01	AU	41419/01	AU	43134/01
		ΑU	43135/01	AU	43136/01	AU	43137/01
		ÁU	45262/01	AU	45354/01	AU	47190/01
		AU	47191/01	AU	49052/01	AU	49053/01
		AU	49054/01	ΑU	50767/01	AU	50768/01
		AU	50769/01	AU	50770/01	AU	50771/01
		AU	50772/01	AU	52878/01	AÜ	52879/01
		AU	55162/01	AU.	57912/01	AU	60969/01
		ΑÚ	62899/01	AU	66787/01	CA	2392398
		CA	2392422	CA	2392428	CA	2392438
		CA	2392450	CA	2392751	CA	2392757
		CA	2393002	CA	2393616	CA	2393618
		CA	2393652	CA	2393912	CA	2393941
	•	CA	2393954	CA	2394022	CA	2394039
		CA	2394841	CA	2395178	CA	2395295
		CA	2395398	CA	2395403	CA	2395654
		CA	2395666	CA	2395671	ÇA	2395676
		CA	2395693	CA	2395699	CA	2395724

Information on patent family members

l c	A 2395729	CA	2395734	CA	2395738	
C.		CA	2395794	CA	2395811	
C.	A 2395815	CA	2395816	CA	2395827	
C	A 2395838	CA	2395849、	CA	2395857	
C	A 2395858	CA	2395872	· CA	2395885	
C	A 2395889	CA	2396719	CA	2397407	
C	A 2397502	CA	2397839	CA	2398227	
C	A 2398275	CA	2398411	CA	2398877	
· E	P 1251863	EP	1252176	· EP	1252185	
E	P 1252289	EP	1252290	EP	1252297	
E	2 1252302	EP	1252303	EP	1252312	
E	2 1252326	EP	1252337	EP	1254147	
E	P 1254148	EP	1254150	EP ·	1254151	
E	P 1254152	EP	1254153	EP	1254154	
El	P 1254157	EP	1254165	EP	1254170	
E	2 1254171	EP	1254172	EP	1254173	
E	2 1254217	EP	1254218	EP	1254219	
El	2 1254228	EP	1254242	EP	1254248	
EI	P 1254272	EP	1255766	EP	1255767	.
EI	1255768	EP	1255776	EP	1255777	
E	1255778	EP	1255817	EP	1255864	
E	1255869	EP	1259525	EP	1259526	İ
E	1259528	EP	1259531	EP	1259540	
E	1259642	EP	1261380	EP .	1261618	- 1
EF	1261619	EP	1261633	EP	1261634	
EF	1261635	EP	1261637	EP	1261703	
EF	1261742	EP	1261745	EP	1263944	
· EF	1265910	EP	1268527	US	2002042096	
, US	2002042386	US	2002061521	US	2002077270	=
US	2002086330	US	2002086353	US	2002086811	
US	2002086820	US	2002086821	US	2002086822	1
. US	2002086823	US	2002090615	US	2002090672	
US	2002090673	US	2002090674	US	2002094953	
US	2002102638	US	2002119919	.US	2002120103	İ
US	2002132753	US	2002132767	US	2002147140	
US	2002151009	US	2002151479	US	2002160493	1
US	2002161208	US	2002164685	US	2002165137	
					•	•

Information on patent family members

_							
١		US	2002168711	US	2002173454	US	2003013649
		US	2003039993	US	2003039994	US .	2003044890
	•	US	2003044904	US	2003044905	US	2003044907
		US	2003049650	US	2003049652	US	2003049703
		US	2003050231	US	2003054368	US	2003054373
		US	2003054375	US	2003054377	US	2003054379
		US	2003054420	US	2003059875	US	2003059908
		US .	2003077602	US	2003077606	US	2003077703
		US	2003077704	US	2003077808	US	2003082681
		US	2003082758	US	2003092102	US	2003092611
		US	2003092615	US	2003096346	US	2003108907
۱		US	2003125246	US	2003139327	US	2003157508
		US	2003175739	US	2003190707	wo	0154472
١	•	wo	0154473	wo	0154474	wo	0154708
	•	wo	0154733	wo	0155162	wo	0155163
		wo	0155164	WO-	0155167	wo	0155168
		wo	0155173	wo	0155200	wo	0155201
İ		wo	0155202	WO	0155203	wo	0155204
		wo	0155205	wo	0155206	wo	0155207
		wo	0155208	wo	0155300	WO	0155301
		wo	0155302	wo	0155303	WO	0155304
		wo	0155305	wo	0155306	WO	0155307
		wo	0155308	wo	0155309	wo	0155310
ı		wo	0155311	wo	0155312	wo	0155313
ı		wo	0155314	wo	0155315	WO	0155316
		wo	0155317	wo	0155318	WO	0155319
1		wo	0155320	wo	0155321	WO	0155322
ł		wo	0155323	WO	0155324	WO	0155325
	•	wo	0155326	wo	0155327	WO	0155328
		wo	0155329	wo	0155343	WO	0155350
	•	wo	0155355	wo	0155364	wo	0155367
		wo	0155368	wo	0155387	wo	0155388
	•	wo	0155430	wo	0155440	WO	0155441
		wo	0155447	wo	0155448	wo	0155449
		wo	0157182	wo	0159063	wo	0159064
		wo	0162789	WO	0164703		· · · · · · · · · · · · · · · · · · ·
	WO 02071928	US	2003087250				

Information on patent family members

wo	02102235	ΑU	69022/00	CA	2381699	EP	1204764
		US	2002015970	US	2002019330	US	2003124579
		US	2003152926	wo	0111086	WO	02079492
	·	WO	02086443	wo	02098358	WO	03003906
_		wo	03025138	wo	03042661		
US	2003004102	AU	10703/99	AU	15324/99	AU	22122/99
		CA	2305385	CA	2309358	CA	2321677
		CA	2347835	CA	2353775	EP	1025227
		EP	1032667	EP	1135485	EP	1141284
		EP	1141289	EP	1159419	EP	1159422
ļ		EP	1173563	EP	1220905	EP	1263948
		US	6410708	US	2002072067	US	2002072092
		US	2002072496	US	2002072497	US	2002103125
		US	2002119118	US	2002119130	US	2002123463
		US	2002127576	· US	2002127584	US	2002132252
		US	2002132253	US	2002142961	US	2002156006
		US	2002160384	US	2002161199	US	2002169284
		US	2002177164	US	2002177165	US	2002177188
		US	2002177553	US	2002182638	US	2002182673
		US	2002183493	US	2002183494	US	2002183505
		US	2002192209	US	2002192706	US	2002192752
		US	2002193299	US	2002193300	US	2002197615
		US	2002197674	US	2002198148	US	2002198149
		US	2003003531	US	2003003546	US	2003004311
1 -		US	2003008297	US	2003008352	US	2003008353
		US	2003009012	US	2003009013	US	2003013855
		US	2003017476	US	2003017540	US	2003017541
		US	2003017542	· US	2003017543	US	2003017544
		US	2003017563	US	2003017981	US	2003017982
		US	2003018168	US	2003022187	US ·	2003022239
		US	2003022293	US	2003022294	US	2003022295
		US	2003022296	US	2003022297	US	2003022298
		US	2003022300	US	2003022301	US	2003022328
		US	2003022331	US	2003027162	US	2003027163
		us	2003027212	US	2003027262	US	2003027263
		US	2003027264	US	2003027265	US	2003027266
		US	2003027267	US	2003027269	US	2003027270
•							

Information on patent family members

					TIC	200202222
	US	2003027271	US	2003027272	US	2003027273
	US	2003027274	US	2003027275	US	2003027276
	US	2003027277	US	2003027278	US	2003027279
	US	2003027280	US	2003027281	US	2003027324
·	US	2003027754	US	2003027985	US	2003027986
	US	2003027992	US	2003027993	US	2003032023
	US	2003032057	US	2003032062	US	2003032063
	US	2003032101	US	2003032102	US	2003032103
	US	2003032104	US	2003032105	US	2003032106
	US	2003032107	US	2003032108	US	2003032109
	US	2003032110	US	2003032111	US	2003032112
	US	2003032113	US	2003032114	US	2003032115
	US	2003032116	US	2003032117	US	2003032118
	US	2003032119	US	2003032120	US	2003032121
	US	2003032122	US	2003032123	US	2003032124
	US	2003032125	US	2003032126	US	2003032127
	US	2003032128	US	2003032129	US	2003032130
	US	2003032131	US	2003032132	US	2003032133
	US	2003032134	US	2003032135	US	2003032136
	US	2003032137	US	2003032138	US	2003032139
	US	2003032140	US	2003032155	US	2003032156
	US	2003034993	US	2003036117	US	2003036118
	US	2003036120	US	2003036121	US	2003036122
	US	2003036123	US	2003036124	US	2003036125
	US	2003036126	US	2003036127	US	2003036128
	US	2003036129	US	2003036130	US	2003036131
	US	2003036132	US	2003036133	US	2003036134
	US	2003036135	US	2003036136	US	2003036137
·	US	2003036138	US	2003036139	US	2003036140
	US	2003036141	US	2003036142	US	2003036143
	US	2003036144	US	2003036145	US	2003036146
	US	2003036147	US	2003036148	US	2003036149
	US	2003036150	US	2003036151	US	2003036152
	US	2003036153	US	2003036154	US	2003036155
	US	2003036156	US	2003036157	US	2003036158
	US	2003036159	US	2003036160	US	2003036161
	US	2003036162	US	2003036164	US	2003036165
· ,						•

Information on patent family members

US	2003036179	US	2003036180	US	2003036634	
. US	2003038826	US	2003038827	US	2003038828	
US	2003039648	US	2003040013	US	2003040014	
US	2003040054	US	2003040055	US	2003040056	
US	2003040057	US	2003040058	US	2003040059	
US	2003040060	US	2003040061	US	2003040062	
US	2003040063	US	2003040064	US	2003040065	
US	2003040066	US	2003040067	US	2003040068	
US	2003040069	US	2003040070	US	2003040071	i
US	2003040072	US	2003040073	US	2003040074	
US	2003040075	US	2003040076	US	2003040077	
US	2003040078	US	2003040473	US	2003043176	
US	2003044806	US	2003044841	US	2003044844	
US	2003044902	US	2003044916	US	2003044917	
US	2003044918	US	2003044919	US	2003044920	
US	2003044921	US	2003044922	US	2003044923	
US	2003044924	US	2003044925	US	2003044926	
US	2003044927	US	2003044928	US	2003044929	
US	2003044930	US	2003044931	US	2003044932	
US	2003044945	US	2003045462	US	2003045463	
US	2003045684	US	2003049633	US	2003049638	
US	2003049681	ÜS	2003049682	US	2003049684	
US	2003049735	US	2003049738	US	2003049741	
US	2003049756	US	2003049816	US	2003049817	
US	2003050239	US	2003050240	US	2003050241	
US	2003050457	US	2003050462	US	2003054359	
US	2003054403	US	2003054404	· US	2003054405	
US	2003054406	US	2003054442	US	2003054454	
US	2003054455	US	2003054456	US	2003054458	
US	2003054459	US	2003054460	US	2003054468	
US	2003054516	US	2003054517	US	2003054986	
US	2003054987	· US	2003055216	. US	2003055222	
US	2003059780	US	2003059782	US	2003059783	
US	2003059831	US	2003059832	US	2003059833	
US	2003059876	US	2003059909	US	2003060406	
US	2003060407	US	2003060600	US	2003060601	
US	2003060602	US	2003064062	US	· 2003064375	

Information on patent family members

. US	2003064407	US	2003064440	US	2003064441	
· US	2003064925	US	2003065142	US	2003065143	
US	2003068793	US	2003068794	US	2003068795	
· US	2003068796	US	2003068797	US	2003068798	
US	2003069178	US	2003069179	US	2003069394	
US	2003069403	US	2003071835	US	2003072745	
US	2003073090	US	2003073129	US	2003073130	
US	2003073131	US	2003073169	US	2003073170	
US	2003073171	US	2003073172	US	2003073173	
US	2003073174	US	2003073177	US	2003073208	
US	2003073210	US	2003073211	US	2003073212	
US	2003073213	US	2003073214	US	2003073215	
US	2003073216	US	2003073624	ŲS	2003073809	
US.	2003073810	US	2003073813	US ·	2003073821	
US	2003077593	US	2003077594	US	2003077659	Ì
US	2003077700	US	2003077718	' US	2003077732	
US	2003077733	. US	2003077776	US	2003077777	
US	2003077778	US	2003077779	US	2003077780	
US	2003077781	US	2003077782	US	2003077783	
US	2003077784	US	2003077785	US	2003077786	
US	2003077787	US	2003077788	US	2003077789	
US	2003077790	US	2003077791	US	2003077792	
US	2003078377	US	2003082199	US	2003082546	
US	2003082626	US	2003082627	US	2003082628	
US	2003082689	US	2003082715	US	2003082716	
US	2003082717	US	2003082759	`US <sub>.</sub>	2003082760	
US	2003082761	US	2003082762	US	2003082763	
US	2003082764	US	2003082765	US	2003082766	
. US	2003082767	US	2003083248	US	2003083461	
US	2003083462	US	2003087304	US	2003087305	
US	2003087344	US	2003087345	US	2003087365	
US	2003087367	US	2003087373	US	2003087374	
US	2003092061	US	2003092104	US	2003092147	
US	2003092883	US	2003096353	US	2003096386	
US	2003096744	US	2003096954	US	2003096955	
US	2003099625	US	2003100061	US	2003100087	
US	2003104536	US	2003104538	US	2003104539	

Information on patent family members

•				1 01/1100	
US	2003104540	US	2003104541	US	2003104542
US	2003104543	US	2003104998	US	2003109672
US	2003113793	US	2003113795	UŞ	2003114652
US	2003119001	US	2003119055	US	2003119097
US	2003119105	US	2003119106	US	2003120053
US	2003124531	. US	2003124661	US	2003125535
us	2003129650	US	2003129695	US	2003130181
US	2003130182	US	2003130483	US	2003130490
US	2003130491	US	2003134284	US	2003134785
US	2003135034	US	2003138882	US	2003138898
US	2003139328	US	2003139329	US	2003148373
US	2003148375	US	2003148376	US	2003153000
US	2003157635	US	2003166104	.US	2003166105
· US	2003166106	US	2003166107	US	2003166108
US	2003166109	. US	2003166110	US	2003166111
US	2003166112	US	2003166113	US	2003166114
US	2003166115	US	2003166117	· US	2003166118
US	2003166119	US	2003166120	US	2003166121
US	2003166122	US	2003166123	US	2003166124
US	2003166125	US	2003166126	US	2003166127
US	2003166128	US	2003166129	US	2003166130
US	2003166131	US	2003166133	US	2003166134
US	2003166135	US	2003166188	US	2003166848
US	2003170718	US	2003170721	US	2003175882
US	2003180311	US	2003180792	US	2003180793
US	2003180794	US	2003180795	US	2003180796
US	2003180836	US	2003180837	US	2003180838
US	2003180839	US .	2003180840	US	2003180841
US	2003180842	US	2003180843	US	2003180844
US	2003180845	US	2003180846	US	2003180847
US	2003180848	US	2003180849	US	2003180850
· US	2003180851	US	2003180852	US	2003180853
US	2003180854	US	2003180855	US	2003180856
US	2003180857	US	2003180858	US	2003180859
US	- 2003180860	US	2003180861	US	2003180862
US	2003180863	US	2003180867	US	2003180904 .
US	2003180905	US	2003180906	US	2003180907

Information on patent family members

1	US	2003180908	US	2003180909	US	2003180910
. 1	US	2003180911	US	2003180912	US	2003180913
1	US	2003180914	US	2003180915	US	2003180916
1	US	2003180917	US	2003180918	US	2003180919
1	US	2003180920	US	2003180921	US	2003180922
1	US	2003180923	US	2003181636	US	2003181637
. 1	US	2003181638	US	2003181639	US	2003181640
1	US	2003181641	US	2003181642	US	2003181643
1	US	2003181644	US	2003181645	US	2003181646
1	US	2003181647	US	2003181648	US	2003181649
1	US	2003181650	US	2003181651	US	2003181652
1	US	2003181653	US	2003181654	US	2003181655
	US	2003181656	US	2003181657	US	2003181666
· 1	US	2003181667	US	2003181668	US	2003181669
1	US	2003181670	US	2003181671	US	2003181672
	US	2003181673	US	2003181674	US	2003181675
ı	US	2003181676	US	2003181677	US	2003181678
· · · · · · · · · · · · · · · · · · ·	US	2003181679	US	2003181680	US	2003181681
1	US	2003181682	US	2003181683	US	2003181684
ļ	US	2003181685	US	2003181696	US	2003181697
Ī	US	2003181698	US	2003181699	US	2003181700
Ţ	US	2003181701	US	2003181702	US	2003181703
τ	US	2003185841	US	2003186318	US	2003186319
ί	US	2003186365	. US	2003186368	US	2003186407
·	US	2003186408	US	2003187189	US	2003187190
_ τ	US	2003187191	US	2003187192	US	2003187193
τ	US	2003187195	US	2003187196	US	2003187197
τ	US	2003187228	US	2003187229	US	2003187230
τ	US	2003187239	US	2003187241	US	2003187242
. τ	JS	2003190321	US	2003190667	US	2003190668
<u>,</u> τ	JS	2003190669	US	2003190698	US	2003190699
τ	JS	2003190701	US	2003190703	US	2003190716
τ	JS	2003190717	US	2003190718	US ·	2003190719
τ	JS	2003190720	US .	2003190721	US	2003190722
τ	JS	2003190723	US	2003190724	US	2003190725
τ	JS	2003190726	US	2003190727	US	2003190728
τ	JS	2003190729	US	2003190730	US	2003190731

Information on patent family members

							5/01100	
1	•	US	2003191281	US	2003191282	US	2003191283	
		· US	2003191284	US	2003191287	US <sub>.</sub>	2003191288	
		US	2003191289	US	2003191290	US	2003191299	
		. US	2003194744	US	2003194780	US	2003194781	
		US	2003194791	US	2003194792	US	2003194793	
		US	2003194794	US	2003195148	US	2003195333	1
	,	US	2003195334	US	2003195344	· US	2003195345	
		US	2003195347	WO	0032221	WO	0037638	
	•	wo	0037640	WO	0053752	WO	0053753	
		WO	0053755	wo	0053756	WO	0053757	
		WO	0053758					
wo	<sup>-</sup> 0170976	AU	49317/01	US	2002048759	US	2003059781	
wo	03068054							
US	2002192678							
wo	0192581	AU	65296/01	CA	2411404	EP	1356092	
		US	2002132237					
wo	0151513	AU	36470/01					
wo	0190304	AU	29508/01	AU	30958/01	AU	36459/01	
		AU	36460/01	AU	36461/01	AU	36462/01	
		AU	36463/01	AU	36464/01	AU	36465/01	
		AU	36466/01	AU	37943/01	AU	37944/01	
		AU	37947/01	AU	37949/01	AU	37950/01	
	•	AU	37951/01	AU	37952/01	AU	37953/01	
		AU	37954/01	AU	37955/01	AU	37957/01	
ļ		AU	37958/01	AU	38585/01	AU	39726/01	
1		AU	39727/01	AU	39728/01	AU	41402/01	
		AU	41403/01	AU	41404/01	AU	41405/01	
		AU	41406/01	AU	41407/01	AU	41408/01	
		AU	41409/01	AU	41410/01	AU	41411/01	
		AU	41412/01	AU	41413/01	AU	41414/01	
		AU	41415/01	AU	41416/01	AU	41417/01	
		AU	41418/01	ΔŬ	41419/01	AU	43134/01	
		AU	43135/01	AU	43136/01	AU	43137/01	
		AU	45262/01	AU	45354/01	AU	47190/01	
		AU	47191/01	AU	49052/01	AU	49053/01	
		AU	49054/01	AU	50767/01	AU	50768/01	
		AU	50769/01	AU	50770/01	AU	50771/01	

Information on patent family members

			<del></del>	TCIAGO	
AU	50772/01	AU	52878/01	AU	52879/01
AU	55162/01	AU	57912/01	AU	60969/01
AU	62899/01	AU	74888/01	CA	2392398
CA	2392422	CA	2392428	CA	2392438
CA	2392450	CA	2392751	CA	2392757
CA	2393002	CA	2393616	CA	2393618
CA	2393652	CA	2393912	CA	2393941
CA	2393954	CA	2394022	CA	2394039
CA	2394841	CA	2395178	CA	2395295
CA	2395398	CA	2395403	CA	2395654
CA	2395666	CA	2395671	CA `	2395676
CA	2395693	CA	2395699	CA	2395724
CA	2395729	CA	2395734	CA	2395738
CA	2395787	CA	2395794	CA	2395811
CA	2395815	CA	2395816	CA	2395827
CA	2395838	CA	2395849	CA	2395857
CA	2395858	CA	2395872	CA	2395885
CA	2395889	CA	2396719	CA	2397407
CA	2397502	CA	2397839	CA	2398227
CA	2398275	CA	2398411	CA	2398877
EP	1251863	EP	1252176	EP	1252185
EP .	1252289	EP	1252290	EP -	1252297
EP	1252302	EP	1252303	EP	1252312
EP	1252326	EP	1252337	EP	1254147
EP	1254148	EP	1254150	EP .	1254151
EP .	1254152	EP	1254153	EP	1254154
EP	1254157	EP	1254165	EP	1254170
EP	1254171 .	EP	1254172	EP	1254173
EP	1254217	EP	1254218	EP	1254219
EP	1254228	EP	1254242	EP	1254248
EP	1254272	EP	1255766	EP	1255767
EP	1255768	EP	1255776	EP	1255777
EP	1255778	EP	1255817	EP	1255864
EP	1255869	EP	1259525	EP	1259526
EP	1259528	EP	1259531	EP	1259540
EP	1259642	EP	1261380	EP	1261618
EP	1261619	EP	1261633	EP	1261634

Information on patent family members

EP	1261635	EP	1261637	EP	1261703
EP	1261742	EP	1261745	EP	1263944
EP	1265910	EP	1268527	US	2002042096
US	2002042386	US	2002061521	US	2002077270
US	2002086330	US	2002086353	US	2002086811
US	2002086820	US	2002086821	US	2002086822
US	2002086823	US	2002090615	US	2002090672
US	2002090673	US	2002090674	US	2002094953
US	2002102638	US	2002119919	US	2002120103
. US	2002132753	US	2002132767	US	2002147140
US	2002151009	US	2002151479	US	2002160493
US	2002161208	US	2002164685	US	2002165137
US	2002168711	US	2002173454	US	2003013649
US	2003039993	US	2003039994	US	2003044890
US	2003044904	US	2003044905	US	2003044907
. US	2003049650	US	2003049652	US	2003049703
US	2003050231	US	2003054368	US	2003054373
US	2003054375	US	2003054377	US	2003054379
US ·	2003054420	US	2003059875	US	2003059908
us	2003077602	US	2003077606	US	2003077703
US	2003077704	US	2003077808	US	2003082681
US	2003082758	US	2003092102	US	2003092611
US	2003092615	US	2003096346	US	2003108907
US	2003125246	US	2003139327	US	2003157508
US	2003175739	US	2003190707	WO	0154472
wo	0154473	wo	0154474	WO	0154708
wo	0154733	wo	0155162	wo	0155163
wo	0155164	wo	0155167	wo	0155168
OW.	0155173	wo	0155200	wo	0155201
· wo	0155202	wo	0155203	wo	0155204
wo	0155205	wo	0155206	wo	0155207
wo	0155208	wo	0155300	WO	0155301
' wo	0155302	wo	0155303	wo	0155304
wo	0155305	wo	0155306	WO	0155307
wo	0155308	wo	0155309	WO	0155310
wo	0155311	WO	0155312	WO	0155313
wo	0155314	wo	0155315	WO	0155316

٠,	INTERNATIONAL SEARCH REPORT  Information on patent family members					International application No. PCT/AU03/01166	
		wo	0155317	wo	0155318	wo	0155319
		wo	0155320	wo	0155321	WO	0155322
		wo	0155323	wo	0155324	WO	0155325
		wo	0155326	wo	0155327	WO	0155328
		wo	0155329	wo	0155343	wo	0155350
		wo	0155355	wo	0155364	wo	0155367
		wo	0155368	wo	0155387	wo	0155388
		wo	0155430	,wo	0155440	wo	0155441
		wo	0155447	wo	0155448	· WO	0155449
		wo	0157182	wo	0159063	WO	0159064
	•	wo	0162789	wo	0164703		
wo	9953040	DE	19817557	EP	1073727	EP	1291422
		US	2003105315			·	
wo	0194629	AU	64559/01	CA	2411601	US	2002081301
		US	2002102531	US	2002102532	US	2002110821
		US	2002115057.	บร	2002115085	US	2002150877
		US	2002165180	US	2003165839	•	
wo	9848010	AU	70139/98	EP	0979277	. US	2002192160
wo	0160860	AU	41541/01				
wo	0155313	AU	29508/01	AU	30958/01	AU	36445/01
		AU	36459/01	AU	36460/01	AU.	36461/01
		AU	36462/01	AU	36463/01	AU	36464/01
		AU	36465/01	AU	36466/01	AU	37943/01
		AU	37944/01	ΑU	37947/01	ΑÜ	37949/01
		AU	37950/01	AU	37951/01	AU	37952/01
		ΑŬ	37953/01	AU	37954/01	AU	37955/01
		ΑŬ	37957/01	AU	37958/01	AU	38585/01
		AU	39726/01	AU	39727/01	AU	39728/01
		AU	41402/01	ΑU	41403/01	AU	41404/01
		AU	41405/01	AU	41406/01	AU	41407/01
		AU	41408/01	AU	41409/01	AU	41410/01
		AU	41411/01	AU	41412/01	ΑŬ	41413/01
		. AU	41414/01	ΑU	41415/01	AU	41416/01
		AU	41417/01	ΑU	41418/01	AU	41419/01
		AU	43134/01	AU	43135/01	ΑŬ	43136/01

45262/01

47191/01

ΑU

AU

ΑU

AU

45354/01

49052/01

AU

AU

43137/01

47190/01

Information on patent family members

AU	49053/01	AU	49054/01	AU	50767/01
AU	50768/01	AU.	50.769/01	AU	50770/01
AU	50771/01	AU	50772/01	AU	52878/01
. AU	52879/01	· AU	52917/01	AU	53323/01
AU	55162/01	ΑU	57912/01	AU	60969/01
AU	62899/01	AU	66787/01	AU	71714/01
. AU	74888/01	CA	2392398	CA	2392422
CA	2392428	CA	2392438	CA	2392450
. CA	2392751	CA	2392757	CA	2393002
CA	2393616	CA	2393618	CA	2393652
CA	2393912	CA	2393941	CA	2393954
CA	2394022	CA	2394039	CA	2394841
CA	2395178	CA	2395295	CA	2395398
CA	2395403	CA	2395654	CA	2395666
CA	2395671	CA	2395676	CA	2395693
CA	2395699	CA	2395724	CA	2395729
· CA	2395734	CA	2395738	CA	2395787
CA	2395794	CA	2395811	CA	2395815
CA	2395816	CA	2395827	CA.	2395838
CA	2395849	. CA	2395857	CA	2395858
CA	2395872	CA	2395885	CA	2395889
CA	2396719	CA	2397407	CA	2397502
CA	2397839	CA	2398227	CA	2398275
CA	2398411	CA	2398877	CA	2400638
CA	2403508	CA	2406649	EP	1251863
EP	1252176	EP	1252185	EP	1252289
EP	1252290	EP	1252297	EP	1252302
EP	1252303	EP	1252312	EP	1252326
EP	1252337	EP	1254147	EP	1254148
. EP	1254150	EP	1254151	EP	1254152
EP	1254153	EP	1254154	EP	1254157
EP	1254165	EP	1254170	EP	1254171
EP	1254172	EP	1254173	EP	1254217
EP	1254218	EP	1254219	EP	1254228
EP	1254242	EP	1254248	EP	1254272
EP	1255766	EP	1255767	EP	1255768
EP	1255776	EP	1255777	EP	1255778
					· ·

Information on patent family members

	_ <del></del> _					
EP	1255817		EP	1255864	EP	1255869
EP	1259525		EP	1259526	EP	1259528
EP	1259531		EP	1259540	EP	1259642
EP	1261380		EP	1261618	EP	1261619
EP	1261633		EP	1261634	EP	1261635
EP	1261637		EP	1261703	EP	1261742
EP	1261745		EP	1263944	EP	1265910
EP	1268527		EP	1268541	EP	1287015
EP	1301524		US	2002042096	US	2002042386
US	2002061521		US	2002077270	US	2002086330
US	2002086353		US	2002086811	US	2002086820
US	2002086821		US	2002086822	US	2002086823
. US	2002090615		US	2002090672	US	2002090673
US	2002090674		US	2002094953	US	2002102638
US	2002106780		US	2002119919	US	2002120103
US	2002132753		US	2002132767	US	2002147140
. US	2002151009		US	2002151479	US	2002160493
US	2002161208		US	2002164685	US	2002165137
US	2002168711		US	2002173454	US	2003013649
US	2003039993		US	2003039994	. US	2003044890
US	2003044904		US	2003044905	US	2003044907
US	2003049650		US	2003049652	US	2003049703
US	2003050231	•	US	2003054368	US.	2003054373
US	2003054375		US	2003054377	US	2003054379
US	2003054420		US	2003059875	US	2003059908
US	2003077602		US	2003077606	US	2003077703
US	2003077704	•	US	2003077808	US	2003082681
US .	2003082758		US	2003092102	US	2003092611
US	2003092615		US	2003096346	US	2003108907
US	2003125246	)	US	2003139327	US	2003157508
. US	2003175739		US	2003190707	wo	0154472
WO	0154473		wo	0154474	wo	0154708
WO	0154733		wo	0155162	WO	0155163
, WO	0155164		wo	0155167	wo	0155168
WO	0155173		wo	0155200	wo	0155201
WO	0155202		wo	0155203	WO	0155204
wo	0155205		wo	0155206	wo	0155207

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

# **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

# IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.